

SEED-BORNE MYCOFLORA OF *CAPSICUM ANNUUM* IMPORTED FROM INDIA

SHARFUN-NAHAR, MUHAMMAD MUSHTAQ* AND IQBAL H. PATHAN

*Department of Plants Protection,
Ministry of Food, Agriculture and Livestock,
Govt. of Pakistan, Jinnah Avenue, Malir Halt, Karachi, Pakistan.*

Abstract

Using standard blotter and deep-freezing techniques, seed-borne mycoflora of 40 samples from consignments of *Capsicum annuum* L. (red chillies var. Dhora, imported from India) was studied. Of the 47 fungal species *Absidia corymbifera*, *Acremonium fusidioides*, *Aspergillus tamarii*, *Blakeslea* sp., *Cephalophora irregularis*, *Cladosporium accacicola*, *Scopulariopsis* sp., *Streptomyces* sp., *Tritirachium* sp., and *Ulocladium tuberculatum* have been not reported before from seeds as well as pericarp of *C. annuum*.

Introduction

Capsicum (*Capsicum annuum* L.) is an important crop and extensively cultivated throughout the tropics and Southern countries such as Bangladesh, India, Pakistan and Sri Lanka. Nutritionally, it contains significant amounts of vitamins A and C. The dried ripe chillies are used as condiment for culinary purposes and seasoning. In medicine, it is used as powerful stimulant and carminative and also to counter irritant. In the present study, mycoflora associated with seeds and fruits of *Capsicum annuum* imported from India has been examined and compared with seed-borne diseases (Khan *et al.*, 1974a&b; Ghafoor & Khan, 1976; Mirza & Qureshi, 1978; Richardson, 1979,1981, 1983; Nasreen *et al.*, 1988, 1992; Hashmi, 1989; Hashmi & Hasal, 1989; Mushtaq & Hashmi, 1997; Abbas *et al.*, 2004).

Materials and Methods

Forty samples of *C. annuum* (red chillies var. Dhora) imported from India were collected from the Department of Plant Protection, Govt. of Pakistan, Karachi and intercepted at Plant Quarantine Office, Lahore. Seeds and pericarp were separately analyzed for the presence of fungi by standard blotter (Anon., 1976) and deep-freezing (Limonard, 1968) methods. Four hundred seeds and 75 pieces of pericarp for each sample were plated on 3 layered moistened blotter discs in 9 cm glass Petriplates @ 25 seeds and 15 pieces of pericarp per plate. Plates were incubated for 7 days at 22±1°C in Eyela La 1000 low temperature incubator. Incubated seeds were examined under microscope. In deep-freezing method, seeds and pericarp were incubated at 22±1°C for 24 hrs., followed by incubation at -20±1°C for 24 hrs., and then at 22±1°C for 5 days. In both methods growing fungi were isolated and purified on potato dextrose agar (PDA), corn meal agar (CMA), and speziellier nahrstoffarmer agar (SNA). The fungi isolated were identified after reference to Barnet & Hunter (1972), Booth (1971), Domsch *et al.*, (1980), Joffe (1986), Neergaard (1977), Nelson *et al.*, (1983), Nirenberg (1990), Pascoe (1990a,b), Raper & Fennel (1965) and Singh *et al.*, (1991). Data of seed-borne mycoflora and pericarp was statistically analyzed by using computer-based software SPSS version 10.

*Corresponding Author, Department of Botany, Adamjee Govt. Sc. College, Karachi-74800, Pakistan
E-mail: mmushtaq72@yahoo.com

Results and Discussion

A total number of 47 species of fungi belonging to 30 genera by standard blotter method as compared to 16 genera and 23 species by deep-freezing method were isolated from seed samples of *Capsicum annuum* (Table 1). Similarly from pericarp, 32 species belonging to 23 genera by standard blotter method and 17 species belonging to 14 genera were isolated by deep-freezing method (Table 2). The isolated fungi were identified as: *Absidia corymbifera*, *Acremonium fusidioides*, *Acremonium* sp., *Alternaria alternata*, *Aspergillus candidus*, *A. flavus*, *A. niger*, *A. ochraceus*, *A. sulphureus*, *A. tamarii*, *A. terreus*, *A. versicolor*, *Aspergillus* spp., *Bipolaris australiensis*, *B. spicifera*, *Blakeslea* sp., *Cephalophora irregularis*, *Cephalosporium* sp., *Chaetomium bostrychodes*, *C. globosum*, *Cladobotryum varium*, *Cladosporium accacicola*, *Cladosporium* spp., *Colletotrichum capsici*, *Curvularia lunata*, *C. pallescens*, *Cylindrocarpon* sp., *Drechslera* spp., *Exserohilum holmii*, *E. rostratum*, *Emericella* sp., *Epicoccum* sp., *Fusarium chlamydosporum*, *F. moniliforme*, *F. pallidoroseum*, *F. proliferatum*, *F. solani*, *F. sporotrichioides*, *F. subglutinans*, *Macrophomina phaseolina*, *Memnoniella echinata*, *Myrothecium* sp., *Paecilomyces* sp., *Penicillium* spp., *Phoma* sp., *Rhizoctonia solani*, *Rhizopus stolonifer*, *Scopulariopsis* sp., *Stachybotrys atra*, *Streptomyces* sp., *Trichoderma harzianum*, *Tritirachium* sp., *Ulocladium tuberculatum* and *Verticillium albo-atrum*.

Aspergillus ochraceus, *Bipolaris australiensis*, *Chaetomium bostrychodes*, *Fusarium chlamydosporum*, *F. sporotrichioides* and *Ulocladium tuberculatum* appear to be new reports only from seeds, whereas, *Acremonium fusidioides*, *Acremonium* sp., *Aspergillus terreus*, *Cladobotryum varium*, *Cladosporium acacicola* and *Emericella* sp., from pericarp of *C. annuum*. On the other hand *Absidia corymbifera* sp., *Aspergillus tamarii*, *Blakeslea* sp., *Cephalophora irregularis*, *Scopulariopsis* sp., *Streptomyces* sp., and *Tritirachium* sp., have been isolated for the first time from seeds as well as pericarp of *C. annuum*. It may be mentioned that species of *Alternaria*, *Colletotrichum*, *Fusarium* and *Phoma* have been reported by Hashmi (1990) from samples of capsicum imported from India.

Significant differences were found between occurrence of fungi in blotter and deep-freezing methods by Bonferroni test that was further confirmed by ANOVA (Table 3 & 4). Occurrence of fungal species was more frequent in blotter test as compared to deep-freezing in both seeds and pericarps. However, for the detection of slow growing and parasitic fungi deep-freezing method appeared more suitable, where the dead embryo of seed provides nourishment, moreover, growth of fast growing saprophytic fungi is checked due to an interrupting deep-freezing period of 24 hrs. Fast growing saprophytic fungi like, *Aspergillus* spp., *Cladosporium* spp., *Penicillium* and *Rhizopus* spp., may be troublesome particularly in the detection of slow growing parasites present internally (Tempe, 1970). *Alternaria alternata*, *Aspergillus flavus*, *A. niger*, *Chaetomium bostrychodes*, *Drechslera tetramera*, *Fusarium moniliforme*, *F. pallidoroseum*, *Paecilomyces* sp., and *Rhizopus stolonifer* were predominantly isolated from seeds of capsicum by both blotter and deep-freezing methods. Likewise, *Absidia corymbifera*, *Alternaria alternata*, *Aspergillus flavus*, *A. niger*, *Colletotrichum capsici*, *Fusarium pallidoroseum*, *F. solani*, *Rhizopus stolonifer*, *Streptomyces* sp., and *Verticillium albo-atrum* were predominantly isolated from pericarp.

Table 1. Seed-borne mycoflora of *Capsicum annuum*.

No.	Mycoflora	Standard blotter method			Deep freezing method		
		¹ Occ. %	Inf. Range	² Mean± ³ SE	¹ Occ. %	Inf. Range	² Mean± ³ SE
1.	<i>Absidia corymbifera</i>	7.50	4.5-10.75	0.65±0.38	---	---	---
2.	<i>Alternaria alternata</i>	42.50	0.5-36.5	4.85±1.42	80.00	1.50-40.00	9.19±1.35
3.	<i>Aspergillus candidus</i>	5.0	2.00	0.10±0.06	---	---	---
4.	<i>A. flavus</i>	62.50	2.00-45.00	9.11±1.82	77.50	3.00-80.00	27.15±4.04
5.	<i>A. niger</i>	52.50	1.00-4.75	1.56±0.30	57.50	2.50-25.00	24.78±3.81
6.	<i>*A. ochraceus</i>	2.50	2.00	0.05±0.04	10.00	0.25-3.00	0.16±0.09
7.	<i>A. sulphureus</i>	5.00	4.50-7.00	0.28±0.20	---	---	---
8.	<i>*A. tamaritii</i>	2.50	1.75	0.04±0.04	---	---	---
9.	<i>A. terreus</i>	10.00	0.50-2.00	0.11±0.05	---	---	---
10.	<i>A. versicolor</i>	7.50	0.50-2.00	0.10±0.06	---	---	---
11.	<i>*Bipolaris australiensis</i>	2.50	1.75	0.04±0.04	5.00	0.25-0.75	0.02±0.01
12.	<i>*Blakeslea</i> sp.	2.50	0.25	0.006±0.006	---	---	---
13.	<i>*Cephalophora irregularis</i>	12.50	1.25-6.25	0.48±0.22	5.00	0.75-1.25	0.05±0.03
14.	<i>Cephalosporium</i> sp.	5.00	1.00-5.50	0.16±0.13	12.50	4.50-9.25	0.89±0.38
15.	<i>*Chaetomium bostrychodes</i>	40.00	0.50-7.00	0.93±0.25	7.50	1.50-6.50	0.31±0.19
16.	<i>C. globosum</i>	20.00	0.50-5.75	0.31±0.15	---	---	---
17.	<i>Cladosporium</i> sp.	10.00	0.50-5.00	0.29±0.16	25.00	0.50-6.25	0.71±0.23
18.	<i>Colletotrichum capsici</i>	12.50	11.25-30.25	2.68±1.17	20.00	5.50-10.25	1.54±0.49
19.	<i>Curvularia lunata</i>	12.50	0.50-3.00	0.20±0.10	30.00	0.25-8.25	0.98±0.31
20.	<i>C. pallescens</i>	17.50	0.50-4.50	0.27±0.13	---	---	---
21.	<i>Cylindrocarpum</i> sp.	2.50	8.00	0.52±0.37	---	---	---
22.	<i>D. holmii</i>	2.50	0.50	0.01±0.01	---	---	---
23.	<i>D. tetramera</i>	22.50	0.50-3.00	0.25±0.09	25.00	1.50-7.25	0.93±0.29
24.	<i>Emicella</i> sp.	5.00	2.50-3.00	0.13±0.09	---	---	---

Table 1. (Cont'd.)

No. Mycoflora	Standard blotter method			Deep freezing method		
	¹ Occ.%	Inf. Range	² Mean± ³ SE	¹ Occ.%	Inf. Range	² Mean± ³ SE
25. <i>Epicoccum</i> sp.	2.50	1.00	0.02±0.02	---	---	---
26. * <i>Exerohilum halodes</i>	22.50	0.50-32.50	1.02±0.80	---	---	---
27. * <i>Fusarium chlamydosporum</i>	2.50	0.75	0.01±0.01	---	---	---
28. <i>F. moniliforme</i>	50.00	0.50-9.00	1.14±0.29	45.00	3.50-15.50	3.63±0.72
29. <i>F. pallidoroseum</i>	15.00	0.50-8.75	0.81±0.39	37.50	0.50-8.50	1.66±0.41
30. <i>F. proliferatum</i>	25.00	0.50-7.00	0.45±0.18	32.50	0.50-10.50	1.76±0.47
31. <i>F. solani</i>	20.00	0.50-6.25	0.39±0.18	25.00	0.50-3.50	0.60±0.18
32. * <i>F. sporotrichioides</i>	5.00	7.50-11.00	0.46±0.32	5.00	0.75	0.03±0.02
33. <i>F. subglutinans</i>	5.00	1.00-3.00	0.10±0.07	---	---	---
34. <i>Macrophomina phaseolina</i>	12.50	0.50-2.50	0.17±0.08	---	---	---
35. <i>Memnoniella echinata</i>	10.00	0.50-1.25	0.09±0.04	---	---	---
36. <i>Myrothecium</i> sp.	10.00	0.25-0.75	0.05±0.02	5.00	0.25-1.50	0.04±0.03
37. <i>Paecilomyces</i> sp.	67.50	2.00-3.25	0.13±0.09	---	---	---
38. <i>Penicillium</i> spp.	7.50	0.25-40.00	3.60±1.05	22.50	0.50-3.75	0.55±0.18
39. <i>Phoma</i> sp.	20.00	0.50-2.00	0.23±0.08	---	---	---
40. <i>Rhizoctonia solani</i>	17.50	2.00-10.00	0.82±0.34	---	---	---
41. <i>Rhizopus stolonifer</i>	70.00	2.00-30.50	9.25±1.52	50.00	2.50-40.50	5.97±1.28
42. * <i>Scopulariopsis</i> sp.	10.00	1.00-3.00	0.21±0.11	---	---	---
43. <i>Stachybotrys atra</i>	5.00	0.50-3.00	0.08±0.07	25.00	0.50-10.50	1.17±0.40
44. <i>Trichoderma harzianum</i>	12.50	0.25-2.00	0.11±0.05	25.00	0.25-6.50	0.46±0.19
45. * <i>Tritirachium</i> sp.	5.00	1.75-2.25	0.10±0.06	---	---	---
46. * <i>Ulocladium tuberculatatum</i>	2.50	5.50	0.13±0.13	---	---	---
47. <i>Verticillium albo-atrum</i>	7.50	8.75-12.25	0.79±0.44	10.00	2.00-6.50	0.41±0.21

1= Occurrence %, 2= Mean infection %, 3= Standard error.

*New reports of fungi isolated from *Capsicum annuum* seeds.

Table 2. Fungi associated with pericarp of *Capsicum annuum*.

No.	Mycoflora	Standard blotter method			Deep freezing method		
		Occ. %	Inf. Range	² Mean [±] SE	Occ. %	Inf. Range	² Mean [±] SE
1	<i>Absidia corymbifera</i>	33.33	1.25-22.50	15.18±6.91	33.33	4.55-12.61	10.75±4.37
2	<i>Alternaria alternata</i>	33.33	0.50-3.25	2.95±1.62	33.33	0.75-3.21	2.15±0.62
3	<i>Acremonium fusidoides</i>	20.00	0.25-2.25	1.33±0.73	---	---	---
4	<i>Acremonium</i> sp.	26.60	0.50-2.75	1.78±0.83	---	---	---
5	<i>Aspergillus flavus</i>	100.00	55.75-100.00	89.21±3.35	100.00	45.21-100.00	75.55±7.78
6	<i>A. niger</i>	100.00	44.50-100.00	69.13±9.71	93.33	13.40-89.21	34.22±8.00
7	<i>A. ochraceus</i>	6.66	0.88	0.88±0.85	---	---	---
8	<i>A. tamarit</i>	6.66	0.12	0.12±0.11	---	---	---
9	<i>A. terreus</i>	6.66	0.18	0.18±0.18	---	---	---
10	<i>Aspergillus</i> spp.	33.33	2.50-28.50	16.08±6.89	---	---	---
11	<i>*Blakeslea</i> sp.	20.00	4.75-9.20	7.65±4.08	33.33	1.21-6.52	5.51±2.54
12	<i>*Cephalophora irregularis</i>	6.66	1.87	1.87±1.54	20.00	1.11-2.14	1.31±0.89
13	<i>*Cladobotryum varium</i>	6.66	0.08	0.08±0.08	---	---	---
14	<i>*Cladosporium accacicola</i>	6.66	0.18	0.18±0.18	---	---	---
15	<i>Cladosporium</i> spp.	13.33	5.68-5.20	0.72±0.72	20.00	0.20-0.90	0.62±0.37
16	<i>Chaetomium globosum</i>	13.33	0.70-6.80	0.9±0.70	13.33	1.20-4.50	1.25±0.83
17	<i>Colletotrichum capsici</i>	33.33	17.56-70.38	11.73±6.41	60.00	7.50-20.60	11.95±5.89
18	<i>Curvularia lunata</i>	6.66	0.08	0.08±0.08	13.33	0.41-0.89	0.26±0.19
19	<i>Drechslera</i> spp.	6.66	0.08	0.08±0.08	---	---	---
20	<i>*Emericella</i> sp.	6.66	0.62	0.62±0.41	---	---	---
21	<i>Fusarium moniliforme</i>	20.00	3.10-3.90	0.72±0.46	20.00	0.80-4.20	1.15±0.63
22	<i>F. pallidoroseum</i>	26.66	7.13-28.50	5.7±3.13	20.00	2.10-4.60	1.17±0.73
23	<i>F. solani</i>	40.00	2.75-4.25	1.5±0.61	40.00	3.20-7.10	1.25±0.54
24	<i>Memnoniella echinata</i>	6.66	0.27	0.27±0.26	---	---	---
25	<i>Penicillium</i> spp.	26.66	1.10-3.20	0.75±0.50	13.33	1.44-2.61	0.27±0.53
26	<i>Rhizopus stolonifer</i>	93.33	10.20-45.60	28.8±5.55	80.00	14.66-48.20	30.85±7.78
27	<i>*Scopulariopsis</i> sp.	26.66	02.40-11.20	6.95±4.68	---	---	---
28	<i>Stachybotrys atra</i>	20.00	0.40-0.08	0.62±0.35	---	---	---
29	<i>*Streptomyces</i> sp.	60.00	10.61-22.30	15.45±5.46	60.00	12.68-25.61	14.75±5.35
30	<i>Trichoderma harzianum</i>	6.66	0.35	0.35±0.63	---	---	---
31	<i>*Tritirachium</i> sp.	6.66	1.07	1.07±1.03	---	---	---
32	<i>Verticillium albo-atrum</i>	26.66	2.62-7.10	4.63±3.03	33.33	3.81-7.35	3.82±1.93

1= Occurrence %, 2= Mean infection %, 3= Standard error. *New reports of fungi isolated from pericarp of *Capsicum annuum*.

Table 3. Analysis of variance of major seed-borne fungi of *Capsicum annum*.

Source	Sum of squares	df	Mean square	F	Probability
Methods (A)	306.9	1	306.9	1052.867	0.001
Fungi (B)	1643.678	9	182.631	627.732	0.001
Samples (C)	946.131	19	49.796	171.158	0.001
A*B	775.865	9	86.207	296.309	0.001
A*C	892.206	19	46.958	161.403	0.001
B*C	4977.553	171	29.108	100.051	0.001
A*B*C	5329.716	171	31.168	107.129	0.001
Error	116.375	400	0.291		

Table 4. Analysis of variance of major fungi isolated from pericarp of *Capsicum annum*.

Source	Sum of squares	df	Mean square	F	Probability
Methods (A)	315.451	1	283.378	976.139	0.001
Fungi (B)	1452.301	10	178.537	595.396	0.001
Samples (C)	112.326	15	38.387	155.138	0.001
A*B	678.112	10	76.105	269.893	0.001
A*C	732.415	15	38.283	127.111	0.001
B*C	3849.678	150	23.819	89.013	0.001
A*B*C	5115.392	150	27.137	98.153	0.001
Error	96.497	400	116		

Quality of seeds and fruits of capsicum were destroyed due to infestation of these fungi that produce mycotoxins causing health hazards in human beings and animals (Hiscocks, 1965). Hashmi (1990) reported anhydrofusarubin, bostrycoidin, fusarubin, moniliformin, vomitoxin (deoxynivalenol), zearalenol and zearalenone mycotoxins from *F. equiseti*, *F. moniliforme*, *F. oxysporum*, *F. pallidorozeum* and *F. solani* isolated from *Capsicum annum*. These mycotoxins are hazardous to human and animal health and mostly carcinogens, mutagens and teratogens (Marasas *et al.*, 1984). Of the various fungi isolated from seeds and pericarp of capsicum, *Aspergillus flavus* was most predominant, known to produce aflatoxin B1, B2, G1 and G2 in food and feed stuff, whereas, *A. ochraceus* produces ochratoxins (Pohland & Wood, 1987). Similarly species of *Alternaria* produce large number of secondary metabolites (idiolites) but most of them have not been evaluated toxicologically. Considering the occurrence of pathogenic and toxigenic fungal species it is proposed that the import of red chillies must be prohibited to prevent entrance of exotic fungi in country as well as to prevent health hazard in national interest.

References

- Abbas, S.Q., S. Shahzad and A. Ghaffar. 2004. *The Fungi of Karachi*. Department of Botany, University of Karachi, Pakistan (In Press).
- Anonymous, T. 1976. International Rules for Seed Testing. *Proc. Intl. Seed Test. Assoc.*, 31:1-52.
- Barnett, H.L. and B.B. Hunter. 1972. *Illustrated Genera of Imperfect Fungi*. Burgess Co. Minneapolis Minnesota, 241 pp.
- Booth, C. 1971. *The Genus Fusarium*. Commonwealth Mycol. Inst. Kew Surrey, England, 237 pp.
- Domsch, K.H., H.W. Gams and T.H. Anderson. 1980. *Compendium of Soil Fungi*. Vol. I Academic Press, N.Y. 1089 pp.

- Ghafoor, A. and S.A.J. Khan. 1976. *List of diseases of economic plants in Pakistan*. Ministry of Food and Agriculture, Islamabad, Pakistan, 86pp.
- Hashmi, A.A. and K.A. Hasal. 1989. *Pests and Pesticides in Pakistan*. Pakistan Agriculture Research Council, Islamabad. 171 pp.
- Hashmi, M.H. 1989. Seed-borne mycoflora of *Capsicum annuum* L. *Pak. J. Bot.*, 21(2): 302-308.
- Hashmi, M.H. and U. Thrane. 1990. Mycotoxins and other secondary metabolites in species of *Fusarium* isolated from seeds of capsicum, coriander and fenugreek. *Pak. J. Bot.*, 22:106-116.
- Hiscocks, E.S. 1965. The importance of moulds in the deterioration of tropical food and feedstuffs. pp 15-26. In: *Mycotoxins in foodstuffs*. G.N. Wogan (Ed.): MIT Press, Cambridge, Massachusetts.
- Joffe, A.Z. 1986. *Fusarium species: Their biology and toxicology*. John Willey and Sons, Inc. 588 pp.
- Khan, S.A.J., S.B. Mathur and P. Neergaard. 1974a. Survey of new seed-borne organism of different important crop seeds from Pakistan. *Abst. Pak. Sci. Conf. D.*, 53-55.
- Khan, S.A.J., S.B. Mathur and P. Neergaard. 1974b. Survey on new seed-borne organism of Pakistan. *Seed Sci. Technol.*, 32: 477-479.
- Limonard, 1968. Ecological aspects of seed health testing. *Proc. Intl. Seed Test. Assoc.*, 33:71-73.
- Marasas, W.F.O., P.E. Nelson and T.A. Toussoun. 1984. *Toxigenic Fusarium species, identity and mycotoxicology*. The Pennsylvania State Univ. Press, University Park and London, 328 pp.
- Mirza, J.H. and M.S.A. Qureshi. 1978. *Fungi of Pakistan*. Deptt of Plant Pathology, University of Agriculture, Pakistan, 31 pp.
- Mushtaq, M and M.H. Hashmi. 1997. Fungi associated with wilt diseases of capsicum in Sindh, Pakistan. *Pak. J. Bot.*, 29: 217-222.
- Nasreen Sultana, S.A.J. Khan and A.K. Khanzada. 1988. Studies on seed-borne mycoflora of chillies and control of fruit rot diseases. *Pak. J. Sci. Ind. Res.*, 31(5): 365-368.
- Nasreen Sultana, A.K. Khanzada and M.A. Aslam. 1992. A new cause of fruit rot of chillies in Pakistan. *Pak. J.Sci. Ind. Res.*, 35(11): 461-462.
- Neergaard, P. 1977. *Seed Pathology*. Vol. I. The MacMillan Press Ltd., London, 389 pp.
- Nelson, P.E., T.A. Toussoun and W.F.O. Marasas. 1983. *Fusarium spp. An Illustrated Manual of Identification*. The Pennsylvania State Univ. Pennsylvania, 203 pp.
- Nirenberg, H.J. 1990. Recent advances in the taxonomy of *Fusarium*. *Stud. Mycol.*, 32: 91-101.
- Pascoe, I.G. 1990a. *Fusarium* morphology I: Identification and characterization of a third conidium type the mesoconidium. *Mycotaxon.*, 37: 121-160.
- Pascoe, I.G. 1990b. *Fusarium* morphology II: Experiments on growing conditions and dispersal of mesoconidia. *Mycotaxon.*, 37: 161-172.
- Pohland, A.E. and G.E. Wood. 1987. Occurrence of Mycotoxins in Foods. pp 35-64. In: *Mycotoxins in Food*. (Eds.): P. Krogh, Academic Press, Inc.
- Raper, K.E. and D.I. Fennel. 1965. *The Genus Aspergillus*. The William and Wilkins Co. Baltimore. 686 pp.
- Richardson, M.J. 1979. *An Annotated List of Seed-borne Diseases*. Int. Seed. Test. Assoc. Zurich, Switzerland, 320 pp.
- Richardson, M.J. 1981. *An Annotated List of Seed-borne Diseases*. Suppl. 1. Int. Seed. Test. Assoc. Zurich, Switzerland, 78 pp.
- Richardson, M.J. 1983. *An Annotated List of Seed-borne Diseases*. Suppl. 2. Int. Seed. Test. Assoc. Zurich, Switzerland, 108 pp.
- Singh, K.J.C., V. Firsavad, U. Thrane and D.B. Mathur. 1991. *An Illustrated Manual for Identification of some seed-borne Aspergillii, Fusaria and Penicilli and their Mycotoxins*. Danish Govt. Inst. Seed Pathol., for Dev. Countries. Ryvangs Alle, 78; DK 2900 Hellerup, Denmark, 133 pp.
- Tempe, J. 1970. Routine methods for determining the health condition of seeds in the seed testing station. *Proc. Int. Seed Test. Assoc.*, 35: 7-15.