

EVALUATION OF THE HETEROTIC AND HETEROBELTIOTIC POTENTIAL OF WHEAT GENOTYPES FOR IMPROVED YIELD

INAMULLAH, HABIB AHMAD*, FIDA MOHAMMAD¹, SIRAJ-UD-DIN²,
GHULAM HASSAN¹ AND RAHMANI GUL¹

Department of Botany, Hazara University, Mansehra, Pakistan

¹*Department of Plant Breeding and Genetics NWFP Agriculture University Peshawar, Pakistan*

²*Department of Botany, Islamia College Peshawar, Pakistan*

Abstract

Heterotic and heterobeltiotic performance of 28 F₁'s obtained through crossing 8 commercial varieties, were evaluated for days to heading, days to maturity, tillers per plant, flag leaf area, plant height, spike length, grains per spike, 1000 grains weight, harvest index and yield per plant. The highest heterotic and heterobeltiotic interactions were recorded in GZ x Der (-5.35 and -4.98%) for days to heading, Tat x SQ (-2.16 and -1.21%) for days to maturity, GZ x Tat (14.92 and 7.46%) for tillers/ plant, SQ x ICP (17.14 and 14.01%) for flag leaf area, Tat x Sar (-3.29 and -2.89%) for plant height, SQ x Der (19.11 and 16.21%) for spike length, Sar x Der (23.14 and 17.60%) for grains per spike, Tkb x Der (28.42 and 28.0%) for 1000 grain weight, SQ x ICP (24.69 and 21.68%) for harvest index and Tat x SQ (56.25 and 26.87%) for yield per plant, respectively. The results revealed that the hybrid combinations Tat x SQ and SQ x ICP could be recommended for improved yield and enhanced biological production of wheat, respectively.

Introduction

To get maximum yield associated with best quality combinations are the aims of the breeding programs. The use of heterosis for getting high yield with improved quality has been largely used in cross-pollinated crops. In self-pollinated crops evidences are available to confirm the potential use of heterosis (Haq & Laila, 1991), suggesting the easiest ways of the possibility of commercial exploitation of genetic potential of wheat crops. A number of explanations could be placed in this regard; *the theory of dominant linked genes* appears to be the most acceptable, both in the concept and utilization of hybrid vigor in self-pollinated crops (Singh *et al.*, 1989; Saleem & Hussain, 1988).

Heterotic studies can also be used for getting information about the increase or decrease of F₁s over their mid parent and better parent (heterobeltiosis). Its use for elaborating the general and specific combining ability, in the selection process is also confirmed.

Keeping in view the general rule of breeding, the higher the heterosis and heritability, the simpler the selection process and greater the response to selection. A diallel of bread wheat was employed for determination of the out yielding effects of wheat hybrids for 10 agronomic traits and their possible exploitation for commercial use.

*Corresponding author's email: drhahmad@gmail.com

Materials and Methods

Eight commercial varieties of Bread wheat (*Triticum aestivum*), viz. Ghaznavi-98 (GZ), Fakhre Sarhad (FS), Tatara (Tat), Takbeer (Tkb), SQ-92 (SQ), Sar-3 (Sar), ICP-3 (ICP) and Der-98 (Der) and 28 F1's got from crossing the parents were planted in a Randomized Complete Block Design, with three replication. Each entry was planted in two rows of 2.5-meter length, with a plant-to-plant and row-to-row distance of 15 and 30 cm, respectively. Standard agronomic practices were followed from sowing till harvest. Data were recorded on 10 agronomic parameters i.e., days to heading, days to maturity, flag leaf area, tillers per plant, plant height, spike length, grains per spike, 1000-grain weight, harvest index and yield per plant.

The values of mid parent heterosis and better parent heterosis (heterobeltiosis) were calculated according to the scheme outlined by Matzingar *et al.*, (1962) as given in the formulae:

$$\text{Heterosis \%} = \frac{F1 - MP}{MP} \times 100$$

$$\text{Heterobeltiosis \%} = \frac{F1 - BP}{BP} \times 100$$

F1, MP and BP in the formulae denote the performance of hybrid, average performance of parents and performance of better parent, respectively.

The data recorded for different characters were analyzed for variance in accordance with the technique of Steel & Torrie (1984). The characters showing significant differences were subjected to heterosis calculation.

Results and Discussion

Analysis of variance, means for the parents and hybrids are given in Tables 1, 2 and 3, respectively. Table 1 indicates highly significant differences among the genotypes for all the traits studied. Character wise discussion of the results is summarized below.

1. Days to heading: Early heading is desirable due to the fact that earlier heading provides sufficient time for grain formation and filling processes of the grain. The negative heterosis for days to heading is therefore useful. Negative heterosis was recorded for 23 out of the 28 crosses (Table 4). Maximum decrease over the mid parent was recorded by the cross GZ x Der (-5.35) followed by FS x Der (-4.46) and Sar x Der (-4.28), whereas maximum negative heterosis over better parent was recorded by GZ x Der (-4.98) followed by FS x Der (-4.46) and Tkb x Der (-3.19). The results are in conformity with the findings of Wu *et al.*, (2001), Sadeque *et al.*, (1991) and Murai (1998) that reveals the importance of heterotic studies for inducing earliness in wheat.

2. Days to maturity: Genotypes with early maturing habits are generally wanted; negative heterosis for days to maturity is therefore a useful parameter. 20 out of 28 crosses revealed negative heterosis (Table 4). Maximum negative mid parent heterosis was recorded for the cross Tat x SQ (-2.16) followed by GZ x Der (-1.61), whereas maximum negative better parent heterosis was shown by the same cross Tat x SQ (-1.21) followed by GZ x Der (-1.04). The cross SQ x Der (+2.29) recorded maximum positive mid parent and better parent heterosis. Mahajan & Nagarajan (2001) and Simon (1989) also observed that heterotic studies could be effectively used for incorporating early maturity in wheat.

Table 1. Analysis of variance for the characters studied.

S. No.	Characters	Mean squares		
		Replications df (2)	Genotypes df (63)	Error df (126)
1.	Days to heading	55.79	20.63**	3.28
2.	Days to maturity	128.92	7.30**	3.57
3.	Tillers per plant	17.07	6.31**	3.61
4.	Flag leaf area	115.38	22.23**	12.96
5.	Plant height	1545.17	67.21**	31.14
6.	Spike length	6.81	1.93**	0.72
7.	Grains per spike	219.64	110.85**	54.30
8.	1000 grain weight	105.59	41.90**	24.22
9.	Harvest index	200.38	35.51**	24.46
10.	Yield per plant	2.42	25.60**	15.57

** P<0.01

3. Tillers per plant: Number of productive tillers directly contributes to plant yield; positive heterosis for tillers per plant is therefore desirable in wheat. Heterotic studies revealed that 13 out of 28 crosses were positive for heterotic effect over their respective mid parent (Table 4). Maximum positive mid parent heterosis was exhibited by the cross GZ x Tat (+14.92) followed by SQ x Sar (+10.89) and Sar x Der (+10.39). Maximum heterobeltiosis was shown by GZ x Tat (+7.46), followed by SQ x Sar (+6.96). Yu *et al.*, (1997), Walia *et al.*, (1993) and Sadique *et al.*, (1991) are also of the opinion that mid parent and better parent heterosis for tillers per plant could be obtained in wheat. Hence the crosses confirming the maximum heterosis GZ x Tat and SQ x Sar could further be employed for getting improved yield in bread wheat.

4. Flag leaf area: Flag leaf area is an effective yield related trait. A larger flag leaf helps to synthesize photosynthates in greater quantities, which are translocated to grains increasing their weight. Positive heterosis for flag leaf area is thus desirable (Kratochvil & Sammons 1990; Khan *et al.*, 1995; Mahmood & Chaudhry, 2000). Heterotic studies for flag leaf area revealed that 16 out of 28 crosses showed positive heterosis (Table 4). Maximum positive mid parent heterosis was recorded by SQ x ICP (+17.14) followed by Tat x ICP (+14.32) and FS x SQ (13.56), whereas maximum positive better parent heterosis was recorded for SQ x ICP (+14.03) followed by FS x Sar (+13.56).

5. Plant height: Plants with greater height are likely to lodge quite often. Tall plants require more energy to translocate solutes to the grain and have lower grain weight. Short stature wheat is therefore preferred and negative heterosis is desirable. Maximum negative heterosis was shown by Tat x Sar (-3.29) followed by FS x Tat (-0.38) and GZ x Tat (-0.37) whereas maximum negative heterobeltiosis was revealed by the hybrid Tat x Sar (-2.89). Sadeque *et al.*, (1991) reported negative heterosis for plant height, whereas positive heterosis for plant height has been recorded by Mahajan *et al.*, (1999), Khan & Bajwa (1989). The result showed that heterotic interaction improves genetic diversity and provides ample chances to select the desired combinations.

Table 2. Calculated means of selected agronomic traits for the parents used.

Parents	Days to heading	Days to maturity	Tillers plant ⁻¹	Flag leaf area (cm ²)	Plant height (cm)	Spike length (cm)	Grains spike ⁻¹	1000-grain weight	Harvest index (%)	Yield plant ⁻¹ (g)
Ghaznavi-98	128.00	165.33	13.33	43.07	81.00	12.66	78.73	32.70	30.66	17.32
Fakhre Sarhad	127.00	163.66	12.13	41.52	92.66	12.06	71.86	37.94	31.33	17.08
Tatara	123.66	163.66	10.33	44.42	97.40	12.26	75.13	45.33	32.33	18.12
Takbeer	125.33	160.00	12.46	36.33	83.33	11.26	70.53	32.94	33.33	20.83
SQ-92	118.66	160.00	11.60	41.01	93.13	12.33	60.80	42.65	27.66	11.30
Sarc-3	121.66	161.33	12.56	41.27	96.60	13.86	61.33	38.14	32.66	18.98
ICP-3	121.66	160.33	12.13	38.82	92.26	12.93	61.53	37.70	26.33	17.51
Dera-98	127.00	164.00	13.40	42.86	97.46	11.73	67.40	32.76	23.66	18.13

Table 3. Recorded means of agronomic traits for the selected 28 F₁'s.

Crosses	Days to heading	Days to maturity	Tillers per plant	Flag leaf area (cm ²)	Plant height (cm)	Spike length (cm)	Grains per spike	1000 grain weight (g)	Harvest index (%)	Yield plant ⁻¹ (g)
GZ x FS	127.66	163.00	13.00	41.27	88.13	11.86	71.06	37.39	30.00	20.86
GZ x Tat	124.66	163.00	13.60	42.27	88.86	12.46	76.40	39.85	27.66	18.06
GZx Tkb	123.66	161.00	12.00	35.98	86.20	12.26	72.53	38.39	31.33	13.17
GZ x SQ	121.00	161.00	13.00	37.67	92.96	12.00	65.46	41.75	28.00	17.50
GZ x Sar	123.00	161.33	10.66	43.63	89.66	12.96	72.20	41.96	31.00	20.86
GZ x ICP	121.33	162.66	10.40	39.27	91.66	14.06	73.20	41.38	27.66	17.86
GZ x Der	120.66	162.00	14.40	41.63	93.26	12.86	76.93	37.45	30.00	19.62
FS x Tat	125.66	162.33	12.33	43.46	94.66	12.33	77.06	41.15	32.66	19.71
FS x Tkb	125.33	163.66	9.60	42.44	91.06	11.80	67.26	37.39	37.33	20.62
FS x SQ	123.66	161.33	9.06	46.87	96.33	13.20	67.73	45.23	28.66	13.45
FS x Sar	125.66	162.00	11.20	47.07	94.73	14.06	67.26	38.79	33.00	18.19
FS x ICP	125.00	162.00	12.73	37.10	94.66	12.80	666.63	38.20	32.33	15.77
FS x Der	121.33	163.00	12.46	42.68	98.26	13.73	76.86	40.0	27.00	13.47
Tat x Tkb	122.33	161.33	10.86	45.08	94.20	12.93	78.13	49.20	34.66	15.57
Tat x SQ	117.66	161.66	10.86	41.70	104.80	13.26	67.46	46.50	33.00	22.99
Tat x Sar	122.00	161.66	10.53	42.66	94.80	12.60	74.13	42.24	34.00	17.64
Tat x ICP	122.00	162.66	11.73	47.58	96.66	12.96	84.00	46.47	29.33	17.95
Tat x Der	121.66	159.33	9.53	45.99	98.13	12.66	77.93	43.74	38.33	15.89
Tkb x SQ	129.66	161.66	12.26	36.25	94.33	12.60	67.46	42.82	34.33	16.37
Tkb x Sar	122.33	161.33	13.20	40.20	92.53	11.53	75.33	42.46	38.66	18.00
Tkb x ICP	122.66	161.66	11.86	40.88	92.66	13.00	71.33	40.76	35.33	19.53
Tkb x Der	121.33	163.66	13.00	42.09	93.30	11.93	73.46	42.19	34.00	15.83
SQ x Sar	119.33	160.00	13.40	41.53	96.00	13.40	56.93	44.72	32.00	19.99
SQ x ICP	120.00	161.33	10.53	46.76	95.00	14.33	63.20	43.61	33.66	16.20
SQ x Der	119.33	163.66	9.63	43.05	96.86	12.20	71.06	39.56	27.33	20.27
Sar x ICP	120.00	160.00	12.80	41.60	98.80	14.06	69.33	47.78	30.66	16.15
Sar x Der	119.00	160.33	14.33	39.15	103.46	14.60	79.26	42.48	29.66	19.13
ICP x Der	121.66	163.00	12.20	37.31	105.60	14.00	61.80	39.89	25.00	18.35

Table 4. Heterotic and Heterobeltiotic values for important traits of wheat in the 28 hybrids.

Crosses	Days to heading		Flag leaf area (cm ²)		Tillers Plant ¹		Days to maturity		Plant height (cm)	
	%MP	%BP	%MP	%BP	%MP	%BP	%MP	%BP	%MP	%BP
GZ X FS	0.13	0.52	-2.41	-4.17	2.09	-2.50	-0.91	-0.40	1.49	8.80*
GZ X Tat	-0.92	0.80	-3.37	-4.84	14.92	7.46	-0.91	-0.40	-0.37	9.71*
GZ X Tkb	-2.36**	-1.32	-9.57	-16.65**	-6.97	-10.00	-1.02	0.62	4.90	6.41
GZ X SQ	-1.89*	1.96*	-10.38*	-12.52*	4.27	-2.50	-1.02	0.62	6.66	14.65**
GZ X Sar	-1.46	1.09	3.46	1.30	-17.63*	-20.00*	-1.22	0.22	0.97	10.69*
GZ X ICP	-2.80**	-0.27	-4.07	-8.81	-18.32*	-22.00*	-0.10	1.45	5.80	13.16**
GZ X Der	-5.35**	-4.98**	-3.11	-3.34	7.73	2.00	-1.61*	-1.04	4.52	15.14**
FS X Tat	0.26	1.61	1.14	-2.15	9.79	1.64	-0.81	-0.81	-0.38	2.15
FS X Tkb	-0.66	0.00	9.03	2.21	-21.95*	-22.99*	1.13	2.29**	3.48	9.28*
FS X SQ	0.67	4.21**	13.56*	12.86*	-23.59*	-25.27**	-0.30	0.83	3.69	3.95
FS X Sar	1.07	3.28**	13.71*	13.56	-9.31	-10.87	-0.30	0.41	0.10	2.23
FS X ICP	0.53	2.73	-7.65	-10.65	4.94	4.94	0.00	1.03	1.29	1.51
FS X Der	-4.46**	-4.46**	1.14	-0.43	-2.34	-6.96	-0.50	-0.40	3.36	6.04
Tat X Tkb	-1.74	-1.07	11.66*	1.50	-4.67	-12.83	-0.30	0.83	4.24	13.04**
Tat X SQ	-2.88**	-0.84	-2.38	-6.12	-0.91	-6.32	-2.16**	-1.21	10.00**	12.52**
Tat X Sar	-0.54	0.27	-0.42	-3.95	-8.00	-16.18	-0.51	0.20	-3.29	-2.89
Tat X ICP	-0.54	0.27	14.32**	7.11	4.45	-3.27	-0.20	0.83	1.93	4.76
Tat X Der	-2.92**	-1.61	5.39	3.54	-19.66**	-28.85**	-0.71	-0.61	0.71	0.75
Tkb X SQ	-1.91*	0.84	-6.26	-11.61*	1.93	-1.60	-0.84	-0.41	6.91	13.20**
Tkb X Sar	-0.94	0.54	3.60	-2.59	4.45	5.03	0.62	1.04	2.85	11.04*
Tkb X ICP	-0.67	0.82	8.79	5.31	-3.52	-4.81	0.72	0.83	5.54	11.20*
Tkb X Der	-3.83**	-3.19**	6.28	-1.81	0.51	-2.98	1.02	2.29**	3.09	11.84**
SQ X Sar	-0.69	0.56	0.94	0.63	10.89	6.63	-0.41	0.00	1.19	3.07
SQ X ICP	-0.13	1.12	17.14**	14.01**	-11.23	-13.18	0.72	0.83	3.34	3.82
SQ X Der	-2.84**	0.56	2.85	0.44	-22.93*	-28.10**	1.02	2.29**	1.64	4.00
Sar X ICP	-1.36	-1.36	3.89	0.80	3.64	1.85	-0.51	-0.20	4.62	7.08
Sar X Der	-4.28**	-2.19*	-6.92	-8.66	10.39	6.96	-1.43*	-0.61	6.63	7.10*
ICP X Der	-2.94**	-0.82	-0.65	-12.96**	-4.43	-8.95	0.51	1.66*	11.31**	14.45**

6. Spike length: In case of spike length 22 and 19 out of the 28 crosses showed positive heterosis over mid parent and better parent respectively (Table 5). Maximum mid parent and better parent heterosis values were +19.11 and +16.21 respectively, for the hybrid SQ x Der which was followed by FS x Der (+15.40) and (+13.81), respectively. The difference in response of genotypes towards heterosis is a studied phenomena e.g., Moiscu *et al.*, (1984) and Thakur *et al.*, (1991) have reported positive mid parent heterosis, while Sadeque *et al.*, (1991) have recorded negative heterosis for spike length in wheat genotypes.

7. Grains per spike: Grains per spike directly determine the yield potential of a genotype. Analysis of the data revealed that 17 out of 28 crosses were positive mid parent heterotic and 8 out of 28 crosses were positive better parent heterotic in interaction. Maximum positive mid parent heterosis was recorded for Sar x Der (+23.14) followed by Tat x ICP (+22.92) and Sar x ICP (+12.85), whereas positive better parent heterosis was showed by Sar x Der (+17.60) followed by Sar x ICP (+12.67) and Tat x ICP (+11.80). These results could be verified from the findings of Saleem & Hussain (1988), Khan & Bajwa (1989) and Tiwari & Chakraborty (1992) who observed mid and better parent heterosis for grains per spike in wheat.

8. 1000 grain weight: Data on means for 1000-grain weight revealed that it was highest for the hybrid Tat x Tkb (49.20 g) and Sar x ICP (47.78 g), as shown in Table 3. For the 1000-grain weight 27 out of 28 crosses displayed positive heterosis (Table 5). Maximum positive mid parent heterosis revealed by the hybrid Tkb x Der (+28.42) followed by Sar x ICP (+26.0) and Tat x Tkb (+25.72) whereas maximum positive better parent heterosis was observed for the cross Tkb x Der (+28.0) followed by Sar x ICP (+25.27) and GZ x Tkb (+16.56). Positive heterosis for 1000-grain weight have also been reported by Pickett (1993), Saleem & Hussain (1988). The predominant heterotic interaction with respect to 1000-grain weight, in all the hybrids showed the effectiveness of heterosis for increased grain yield.

9. Harvest index: The highest mean value for harvest index was recorded (Table 3) for Tkb x Sar (38.6) and FS x Tkb (37.3). Heterotic studies revealed that 8 out of 28 crosses were positive by heterotic over mid parent value for harvest index (Table 5). Maximum positive heterosis for harvest index over mid parent was displayed by SQ x ICP (+24.7%), Tkb x Der (+19.3%) and Tkb x ICP (+18.4%). The maximum positive heterosis over better parent displayed by the cross SQ x ICP (+21.6%) was however followed by Tkb x Sar (+16.01%) and FS x Tkb (+12.0%). Wu *et al.*, (2001) and Murai (1998) have also reported positive heterosis for harvest index.

10. Yield per plant: The highest mean value was recorded for the cross Tat x SQ (22.9g) followed by GZ x FS (20.8g) per plant (Table 3). Heterotic studies revealed that 18 out of 28 crosses were positive by heterotic interaction for yield per plant (Table 5). Maximum positive heterosis for yield per plant over mid parent was displayed by Tat x SQ (+56.25) followed by SQ x Der (+37.68) and SQ x Sar (+32.04). The maximum positive heterosis over better parent was displayed by Tat x SQ (+26.87) followed by GZ x FS (+20.39) and SQ x Der (+11.76). Singh *et al.*, (2004), Afia *et al.*, (2000) and Prasad *et al.*, (1988) have reported mid parent and better parent heterosis for yield per plant.

Table 5. Heterotic and Heterobeltiotic values for important traits of wheat in the 28 hybrids

Crosses	Spike length (cm)		Grains per spike		1000 grain weight (g)		Harvest index (%)		Yield plant ¹ (g)	
	%MP	%BP	%MP	%BP	%MP	%BP	%MP	%BP	%MP	%BP
GZ X FS	-4.04	-6.31	-5.62	-9.73	5.85	-1.44	-3.22	-4.52	21.24	20.39
GZ X Tat	0.00	-1.57	-0.68	-2.87	2.14	-12.08	-12.16	-14.43	1.89	-0.34
GZ X Tkb	2.50	-3.15	-8.91	-9.93	16.97	16.56	-2.08	-6.00	-30.93*	-36.74**
GZ X SQ	-4.00	-5.25	-6.16	-16.85**	10.82	-2.09	-4.0	-8.69	22.23	1.00
GZ X Sar	-2.26	-6.49	3.09	-8.29	18.46*	10.01	-2.10	-5.10	14.92	9.92
GZ X ICP	9.89*	8.76*	4.37	-7.02	17.54*	9.75	-2.92	-9.78	2.53	1.97
GZ X Der	5.46	1.57	5.29	-2.82	14.41	14.31	10.42	-2.17	10.46	8.17
FS X Tat	1.36	0.54	4.85	2.58	-1.16	-9.22	2.16	1.03	11.98	8.77
FS X Tkb	1.14	-2.20	-11.72*	-16.47**	5.52	-1.4	15.46	12.00	8.79	-0.99
FS X SQ	8.19	7.02	2.11	-5.75	12.25	6.05	2.82	-8.51	-5.22	-21.24
FS X Sar	8.48*	1.44	1.00	-6.40	1.98	1.70	3.12	1.02	0.89	-4.14
FS X ICP	2.40	-1.03	-0.10	-7.28	1.01	0.70	12.13	3.19	-8.82	-9.95
FS X Der	15.40**	13.81**	10.38	6.95	13.14	5.42	-1.81	-13.82	-23.49	-25.71
Tat X Tkb	9.91*	5.43	0.38	-2.98	25.72**	8.53	5.58	4.00	-20.05	25.25*
Tat X SQ	7.85	7.56	-0.73	-10.20	5.70	2.58	10.0	2.06	56.25**	26.87**
Tat X Sar	-3.57	-9.13*	8.64	-1.33	1.19	-6.82	4.60	4.08	-4.80	-7.02
Tat X ICP	2.91	0.25	22.92**	11.80*	11.94	2.52	-0.00	-9.27	0.72	-0.95
Tat X Der	5.55	3.26	9.35	3.72	12.01	-3.51	8.33	-6.18	-12.33	-12.36
Tkb X SQ	6.77	2.16	-4.52	-16.22**	13.30	0.40	11.47	2.00	1.87	-21.14
Tkb X Sar	-8.22*	-16.82**	6.20	-6.45	19.98*	11.79	17.17	16.00	-9.53	-13.55
Tkb X ICP	7.43	0.41	0.42	-11.42*	15.41	8.12	18.43	6.00	1.87	-6.22
Tkb X Der	3.76	1.70	-0.67	-8.77	28.42**	28.00**	19.29	2.00	-18.75	-24.0*
SQ X Sar	6.87	0.96	-6.76	-7.17	10.70	4.86	6.07	-2.04	32.04*	5.35
SQ X ICP	6.06	3.60	3.32	2.70	8.56	2.26	24.69**	21.68*	12.43	-7.49
SQ X Der	19.11**	16.21**	10.86	5.44	5.0	-7.16	6.49	-1.20	37.68*	11.76
Sar X ICP	4.97	1.44	12.85	12.67	26.00**	25.27**	3.95	-6.12	-11.49	-14.91
Sar X Der	14.06**	5.28	23.14**	17.60**	19.81**	11.35	5.32	-9.18	3.09	0.80
ICP X Der	13.51**	8.24	-4.13	-8.30	13.21	5.79	0.00	-5.06	2.97	1.21

MP= Mid Parent, BP= Better Parent value

References

- Afiah, S.A.N., N.A. Mohammad and M.M. Saleem. 2000. Statistical genetic parameters, heritability and graphical analysis in 8 X 8 wheat diallel crosses under saline conditions. *Annals of Agricultural Science Cairo*, 45: 257-280.
- Haq, I. and Laila. 1991. Diallel analysis of grain and other agronomic traits in durum wheat. *RACHIS, Barley and wheat Newsletter*, 10: 8-12.
- Khan, N.I and M.A. Bajwa. 1989. Potential of hybrid wheat in Punjab. *Sarhad J. Agric.*, 5: 381-386.
- Khan, N.U., M.S. Swati, Gul Hassan and Q. Nawaz. 1995. Heterosis exhibited by some morphological traits of diallel crosses in wheat (*Triticum aestivum* L.). *Sarhad J. of Agri.*, 11: 485-489.
- Kratochvil, R.J. and D.J. Sammons. 1990. A comparison of soft red winter wheat F-2 populations, their F1 hybrids and parents. *J. Prod. Agric.*, 3: 363-367.
- Mahajan, V. and S. Nagarajan. 2001. Hybrid wheat evaluation by raised bed fixed plot drill. *ICAR News (A Science and technology news letter)*, 7: 19.
- Mahajan, V., S. Nagarajan, M. Srivastav, V. Kumar and Ganga Rao. 1999. Commercial heterosis in wheat- an overview. *Rachis*, 18: 13-16.
- Mahmood, N. and M.A. Chaudhry. 2000. Inheritance of flag leaf in bread wheat genotypes. *Wheat Information Service*, 90: 7-12.
- Moiscu, A., G. Nedelea and V. Sonea. 1984. Heterosis in components of wheat yield. *Agronomie*, 19: 73-80.
- Murai, K. 1998. F₁ seed production efficiency by photoperiod sensitive, cytoplasmic male sterility and performance of F1 hybrid lines in wheat. *Breed. Sci.*, 48: 35-40.
- Pickett, A.A. 1993. Hybrid wheat: results and problems. *Adv. Plant Breed., Suppl. J. Plant Breed.*, 15: 1-259.
- Prasad, K.D., M.F. Haque and D.K. Ganguli. 1998. Heterosis studies for yield and its components in bread wheat (*Triticum aestivum* L.). *Indian J. of Gen. and Pl. Br.*, 58: 97-100.
- Sadeque, Z., A. Bhowmik and M.S. Ali. 1991. Estimates of heterosis in wheat (*Triticum aestivum* L.). *Annals of Bangladesh Agriculture*, 1: 75-79.
- Saleem, M and S. Hussain. 1988. Estimation of heterosis in yield components of wheat. *Pak. J. Agric. Res.*, 9: 1-5.
- Simon, M.R. 1989. Hybrid vigour and combining ability in wheat (*Triticum aestivum* L.) for yield and its components, tillering, height and heading date. *Plant breeding Abst.*, 060-07348.
- Singh, H., S.N. Sharma and R.S. Sain. 2004. Heterosis studies for yield and its components in bread wheat over environments. *Hereditas*, 141: 106-114.
- Singh, I. and S.K. Sharma. 1989. Heterosis in relation to general and specific combining ability effects in wheat. *Indian. J. Agric. Res.*, 23: 163-168.
- Steel, R.G.D. and J.H. Torrie. 1984. *Principles and procedures of statistics*. Mc Graw Hill Book Co.
- Thakur, K.S., T. Dawa, P. Plaha and G.S. Sethi. 1991. Heterosis in spring x winter wheat crosses. *SABRO Journal of Plant Breeding and Genetics*, 23: 15-19.
- Tiwari, V. and S.K. Chakraborty. 1992. Heterosis studies in bread wheat (*Triticum aestivum* L.). *Environ. Ecol.*, 10: 197-202.
- Walia, D.P., T. Dawa, P. Plaha and H.K. Chaudhary. 1993. Gene action and heterosis in bread wheat. *Crop Improvement Society of India*, pp. 84-85.
- Wu, L.M., Z.F. NI, Z.K. Wang, Z. Lin and Q.X. Sun. 2001. Relationship between differential expression patterns of multigene families and heterosis in a wheat diallel crosses. *Yi Chuan Xue Bao*, 28: 256-266.
- Yu, Y.J., Q.S. Chen, J.Y. Yin and Q.S. Chen. 1997. A study on the use of wheat heterosis in the Beijing area - an analysis of heterosis in certain important wheat varieties and lines. *Beijing Agricultural Sciences*, 15: 13-15.

(Received for publication 17 March 2006)