

## ACCUMULATION OF CADMIUM IN THREE SUNFLOWER (*HELIANTHUS ANNUUS* L.) CULTIVARS

JIN ZOU<sup>1</sup>, PU XU<sup>1</sup>, XI LU<sup>1</sup>, WUSHENG JIANG<sup>2</sup> AND DONGHUA LIU<sup>1\*</sup>

<sup>1</sup>*Department of Biology, College of Chemistry and Life Sciences, Tianjin Normal University, Tianjin 300387, China*

<sup>2</sup>*Library, Tianjin Normal University, Tianjin 300387, P.R. China*

### Abstract

The effects of different concentrations of Cadmium on the growth of sunflower (*Helianthus annuus* L.) and its accumulation in roots, stem and leaves were investigated using inductively coupled plasma atomic emission spectrometry (ICP-AES). The concentrations of Cadmium chloride ranged from  $10^{-5}$  M to  $10^{-3}$  M. Seedlings of three sunflower cultivars viz., No. 665, RH118 and QFS14 exposed to  $10^{-3}$  M Cd exhibited substantial growth reduction and all of them died 10 days after treatment application. Growth of roots and shoots was inhibited at concentrations of  $10^{-5}$  M and  $10^{-4}$  M Cd during the entire experiment (20 days). Cadmium accumulation in roots stems and leaves increased significantly ( $p < 0.05$ ) with increasing Cd concentration. Cadmium was concentrated mainly in the roots, and variable amounts of Cd were also transported to stem and leaves. Among the three cultivars, RH118 produced more roots and higher biomass than No. 665 and had a greater ability to accumulate Cd when compared with QFS14.

### Introduction

Cadmium (Cd) a trace heavy metal is considered a hazardous pollutant due to its high toxicity and solubility in water (Lockwood, 1976). At low concentration it has stimulatory effect on root growth of *Allium sativum* (Liu *et al.*, 2003/2004), but at higher concentration it is toxic and directly or indirectly inhibits physiological processes such as respiration, photosynthesis, cell division, plant-water relationships, N metabolism and mineral nutrition, resulting in poor growth and low biomass (Barceló & Poschenrieder, 1990; Sanità & Gabbrielli, 1999; Liu *et al.*, 2003/2004). Cadmium can be easily taken up by plants, transferred through food chains to impart adverse effects on human health (Nordberg, 2003; Wagner, 1994).

Emphasis has become more prevalent towards the problems of Cd pollution with the development of modern industry and agriculture. Anthropogenic activities, such as mining, industry, agriculture and waste disposal has increased since the beginning of the 20<sup>th</sup> century (Alloway, 1995). Most conventional remediation approaches do not provide an acceptable solution to treating contamination caused by metal pollution. Alternatively, phytoremediation provides an attractive strategy, being low cost and environmentally sustainable (McGrath *et al.*, 2002; Salt *et al.*, 1998). The idea of using rare plants which hyperaccumulate metals to selectively remove and recycle excessive soil metals was introduced by Chaney (1983). Salt *et al.* (1996) also considered phytoremediation as an emerging technology using selected and engineered metal-accumulating plants for environmental clean-up. Metal-accumulating plants can accumulate unusually high concentrations of heavy metals in both roots and shoots from polluted soil and waters (Kumar *et al.*, 1995; Dushenkov *et al.*, 1995). A few terrestrial plant species e.g. *Thlaspi* sp., (Lombi *et al.*, 2000) and *Arabidopsis halleri* (Bert *et al.*, 2002) were reported to accumulate high concentrations of Cd, higher than the hyperaccumulation criterion, 0.01% of shoot dry matter (Baker *et al.*, 2000).

---

E-mail: donghua@mail.zlnet.com.cn

Sunflower (*Helianthus annuus*), a fast-growing crop, has a reasonable tolerance to heavy metals. It has been used for rhizofiltration because it has a high root uptake of metals but shows low efficiency in their translocation from root to shoot (Saxena *et al.*, 1999; Kamnev & Van der Lelie, 2000; Lin *et al.*, 2003).

Few reports on cadmium accumulation by *H. annuus* are available (Madejón *et al.*, 2003; Pena *et al.*, 2006). The aim of this investigation was to study the effects of different Cd concentrations on growth of sunflower. In addition, the accumulation of Cd by sunflower roots, stem and leaves was measured using inductively coupled plasma atomic emission spectrometry (ICP-AES).

## Materials and Methods

**Plant cultivation and treatment application:** Seeds of three sunflower (*Helianthus annuus* L.) cultivars No. 665, RH118 and QFS14 used in the present investigation were provided by Qinfeng Agricultural Science Company in Yangling, Shanxi province, P.R. China.

Healthy and equal-sized seeds were chosen from each cultivar, soaked in tap water for 12 h and germinated in Petri dishes in the dark at a constant temperature of 25°C for 48 h. Seeds were allowed to grow in vermiculite in a climate chamber with a day/night period 14/10 h, a day/night temperature and humidity regime of 20°C/25°C and 55/75% relative humidity (RH), respectively. After 7 days 20 sunflower seedlings were selected, fixed on a polystyrol-plate in a pot containing 2L of 1/2 modified Hoagland nutrient solution (Stephan & Prochazka, 1989) for 10 days, respectively. The Hoagland's solution consisted of 5 mM Ca(NO<sub>3</sub>)<sub>2</sub>, 5 mM KNO<sub>3</sub>, 1 mM KH<sub>2</sub>PO<sub>4</sub>, 50 µM H<sub>3</sub>BO<sub>3</sub>, 1 mM MgSO<sub>4</sub>, 4.5 µM MnCl<sub>2</sub>, 3.8 µM ZnSO<sub>4</sub>, 0.3 µM CuSO<sub>4</sub>, 0.1 µM (NH<sub>4</sub>)<sub>6</sub>Mo<sub>7</sub>O<sub>24</sub> and 10 µM FeEDTA adjusted to pH 5.5. Seeds were next treated with different concentrations of Cd ranging from 10<sup>-5</sup> M to 10<sup>-3</sup> M for 20 days. Cadmium was provided as Cadmium chloride. The Cd solutions were prepared in deionized water and were added to the full strength Hoagland nutrition solution. The full strength Hoagland solution without Cd was used for the control plants. The nutrient solutions were continuously aerated and changed regularly every 5 days until the seedlings were harvested.

**Macroscopic observations:** The lengths of roots and stems were measured after 5, 10, 15 and 20 d, respectively. Ten seedlings from each treatment were harvested based on uniformity in size and colour after 20 days of incubation. The seedlings were removed from solution and washed thoroughly with running tap water for 30 min. and then with deionized water to remove traces of nutrients and Cd ions from root surfaces. The samples were dried for 3 days at 45°C, for 1 day at 80°C, and for 12 h at 105°C.

**Estimation of total Cd:** All dried plant samples were prepared using a wet-digestion method (Piper, 1942). Tissue Cd concentrations were analyzed using inductively coupled plasma atomic emission spectrometry (ICP-AES, LEEMAN LABS INC., New Hampshire, USA) as described by Duan (2003).

**Statistical analysis:** Data were expressed as mean±standard error (SE) and were analyzed through analysis of variance (ANOVA) using Sigma statistical software (Jandel Scientific Corporation, USA). Test of equality of averages using a *t*-test was applied equally. The statistical significance was set at *p*<0.05.

**Table 1. Cadmium concentration ( $\mu\text{g/g DW}$ ) and distributive changes in roots, stem and leaves of *Helianthus annuus L.***

Cultivars	Treatment (M)	Total amount (%)	Root (%)	Stem (%)	Leaf (%)
No. 665	Control	15.8 (100)	5.0 $\pm$ 0.73a (33.1)	4.6 $\pm$ 0.61a (35.0)	4.2 $\pm$ 0.66a (31.9)
	10 <sup>-5</sup>	762.1 (100)	623.2 $\pm$ 5.7b (81.8)	73.8 $\pm$ 2.42b (9.7)	65.1 $\pm$ 2.73b (8.5)
	10 <sup>-4</sup>	2742.7 (100)	1878.4 $\pm$ 168.33c (68.5)	605.1 $\pm$ 4.45c (22.1)	259.2 $\pm$ 21.16c (9.4)
	10 <sup>-3</sup>	No data	No data	No data	No data
RH118	Control	3.8 (100)	2.3 $\pm$ 0.19a (59.6)	0.4 $\pm$ 0.08a (11.5)	1.1 $\pm$ 0.15a (28.9)
	10 <sup>-5</sup>	693.9 (100)	544.7 $\pm$ 13.12b (78.5)	84.6 $\pm$ 1.49b (12.2)	64.6 $\pm$ 1.61b (9.3)
	10 <sup>-4</sup>	2278.5 (100)	1315.4 $\pm$ 13.40c (57.7)	617.6 $\pm$ 2.61c (27.1)	345.5 $\pm$ 6.02c (15.2)
	10 <sup>-3</sup>	No data	No data	No data	No data
QFS14	Control	0.8 (100)	0.5 $\pm$ 0.06a (67.6)	0.1 $\pm$ 0.10a (18.1)	0.1 $\pm$ 0.13a (14.3)
	10 <sup>-5</sup>	469.9 (100)	316.3 $\pm$ 2.10b (67.3)	71.2 $\pm$ 0.65b (15.1)	82.5 $\pm$ 2.81b (17.6)
	10 <sup>-4</sup>	2030.0 (100)	1150.1 $\pm$ 29.76c (56.7)	511.1 $\pm$ 11.60c (25.2)	368.8 $\pm$ 2.79c (18.1)
	10 <sup>-3</sup>	No data	No data	No data	No data

Values followed by same letters are not significantly different at ( $p > 0.05$ ). Means  $\pm$  SE,  $n = 3$ . No data: all seedlings died at 10<sup>-3</sup> M Cd. Values in parentheses are percent of control

## Results

**Macroscopic symptoms:** The effects of Cd on root growth of cultivars varied with Cd concentration (Fig. 1 A-C). Cadmium inhibited root growth during the entire experiment, when compared with control seedlings. Root growth in sunflower cultivar No. 665 was inhibited significantly, and root length decreased with increasing Cd concentration and duration of treatment (Fig. 1 A). Seedlings exposed to 10<sup>-3</sup> M Cd solution exhibited substantial growth reduction, and root growth even stopped; they appeared thinner and were sparsely branched. The roots became yellow and were slightly decomposed after 2 days of treatment. After 5 days, the roots appeared yellow brown and some were broken. Root growth in the other two cultivars was approximately the same as that of No. 665 (Fig. 1B and C). Biomass of roots was in the order: QFS14 > RH118 > No. 665. The root system of QFS14 appeared qualitatively strongest.

The effects of Cd on shoot growth varied with different Cd concentrations (Fig. 2 A-C). Shoot length in the three cultivars progressively decreased with increasing Cd concentration and duration of treatment. The least shoot growth was in seedlings treated with 10<sup>-3</sup> M Cd. Biomass of shoots was in the order: QFS14 > RH118 > No. 665.

Control seedlings had 5 pairs of leaves when they were harvested. The plants of No. 665 treated with Cd did not grow as large, robust or healthy as the control during the experiment. With increasing Cd concentration, this phenomenon appeared progressively obvious. After 10 days the young laminae exposed to 10<sup>-5</sup> M and 10<sup>-4</sup> M Cd solutions appeared yellow. After 15 days chlorosis was evident in all young laminae. Toxic symptoms began in 10<sup>-3</sup> M Cd 2 days after treatment and after 5 days chlorosis and necrosis appeared. The shoots wilted and leaf margins and leaf apices also wilted. When compared to No. 665, the toxic effects of Cd on laminae of the other two cultivars were less severe.




Fig. 1. A-C Effect of different Cd concentrations and duration of treatment on root length of sunflower cultivars (Vertical bars denote SE, n=10). Columns denoted by a different letter indicate significantly different values at  $p < 0.05$  in one-way ANOVA

Fig. 2. A-C Effects of different Cd concentrations and duration of treatment on shoot length of sunflower cultivars (Vertical bars denote SE, n=10). Columns denoted by a different letter indicate significantly different values at  $P < 0.05$  in one-way ANOVA

At  $10^{-3}$  M Cd, the root and shoot length significantly decreased and the seedlings died after 10 days (Figs. 1 and 2). These phenomena indicated sensitivity of all root and shoot growth to higher concentrations of Cd ( $10^{-3}$  M).

**Accumulation of Cd:** The accumulation of Cd in the roots, stem and leaves of the three cultivars increased significantly ( $p < 0.05$ ) with increasing Cd concentration. Cd ions accumulated primarily in roots (Table 1). The Cd concentration in roots of No. 665 was higher than that in RH118 and QFS14 at  $10^{-4}$  M Cd. Cadmium was also transported to

stem and leaves (Table 1). The Cd shoot/root ratios increased with increasing Cd concentration. Levels of Cd in the three cultivars treated with  $10^{-5}$  M and  $10^{-4}$  M Cd were in the order: roots > stem > leaves, with the exception of roots > leaves > stem at  $10^{-5}$  M Cd in QFS14. In contrast, No. 665 showed the best capacity to absorb and take up cadmium ions from solution, while the shoot/root ratios of QFS14 were higher than the other two cultivars (QFS14 > RH118 > No. 665).

### Discussion

Cadmium pollution has become a major environmental problem with the development of modern industry and agriculture. Salt *et al.*, (1995, 1996) proposed the use of metal-accumulating plants (e.g., *Brassica juncea*, *Thlaspi caerulescens*, *Zea mays* and *Helianthus annuus*) to remove toxic metals. According to the accepted shoot concentration defining hyperaccumulation being 0.01% (w/w) for Cadmium (Baker *et al.*, 2000), the three cultivars of *H. annuus* can be considered metal hyperaccumulators (Table 1). Among three cultivars, Cd concentrations in the stem and leaves of RH118 were 149.2  $\mu\text{g/g}$  DW (at  $10^{-5}$  M Cd) and 963.1  $\mu\text{g/g}$  DW (at  $10^{-4}$  M Cd) respectively, suggesting a greater ability to accumulate Cd when compared with QFS14. This cultivar, produced more roots and a higher biomass than No. 665, makes it a better choice for remediation of Cd contaminated soils.

The results in the present investigation provide basic information necessary for further development of phytoremediation methods and confirm previous studies (Lin *et al.*, 2003; Madejón *et al.*, 2003; Pena *et al.*, 2006 and Soudek *et al.*, 2006). Localization of Cd in plants depends on the total concentration of contaminant in soil or water. The results from this investigation showed that the Cd content in roots increased with increasing solution concentrations of Cd. Cadmium primarily accumulated in roots, and small amounts were transferred to shoots in the seedlings treated with  $10^{-5}$  M Cd. The reported differences in root uptake and shoot accumulation might be explained by the fact that one of the normal functions of roots is to selectively remove ions from the soil solution (Salt *et al.*, 1997). Cadmium-tolerant plants must be able to prevent the absorption of excess Cd or detoxify the Cd after it has been absorbed.

A few unfavourable changes in the appearance of the Cd-treated seedlings of *H. annuus* in the present investigation were observed, for example browning roots, the decreased number of roots, and growth inhibition of roots and shoots (Figs. 1 and 2). Chlorotic phenomena in seedlings treated with Cd in this investigation were also found, which agrees with the findings of Das *et al.*, (1997).

It is likely that Cd tolerance mechanisms may differ depending on the species. More work still needs to be done to determine the mechanism of high uptake and accumulation of Cd by *H. annuus*. In our investigation, sunflower accumulated substantial amounts of Cd at  $10^{-4}$  M Cd, but seedling growth was severely decreased. Therefore, it is important to select a suitable species with capabilities for high uptake and accumulation of Cd without the plant undergoing severe damage.

### Acknowledgements

This project was supported by the National Natural Science Foundation of China. The authors thank Mr. Xuchuan Duan for analyzing samples of ICP-AES and support, and the anonymous referees for helpful comments.

## References

- Alloway, B.J. 1995. *Heavy metals in soils*. Blackie Academic & Professionals. New York, 386.
- Baker, A.J.M., S.P. McGrath, R.D. Reeves, and J.A.C. Smith. 2000. Metal hyperaccumulator plants: a review of the ecology and physiology of a biological resource for phytoremediation of metal-polluted soils. In: *Phytoremediation of Contaminated Soil and Water*. pp. 85-107. (Eds.): N. Terry and G. Bañuelos, Lewis Publishers, Boca Raton, Germany.
- Barceló, J. and C. Poschenrieder. 1990. Plant water relations as affected by heavy metal stress: a review, *J. Plant Nutr.*, 13: 1-37.
- Bert, V., I. Bonnin, P. Saumitou-Laprade and D. de Laguérie Petit. 2002. Do *Arabidopsis halleri* from nonmetallicolous populations accumulate zinc and cadmium more effectively than those from metallicolous populations. *New Phytol.*, 155: 47-57.
- Chaney, R.L. 1983. Plant uptake of inorganic waste constituents. In: *Land Treatment of Hazardous Wastes*. pp. 50-76. (Eds.): J.F. Parr, P.B. Marsh and J.M. Kla, Noyes Data Corp., Park Ridge, N.J.
- Das, P., S. Samantaray and G.R. Rout. 1997. Studies on cadmium toxicity in plants: a review, *Environ. Pollut.*, 98: 29-36.
- Duan, X.C. 2003. The study on the trace elements in the free growing vine of harvested tuber dioscoreas. *J. Guangxi Normal Univ.*, 21: 122-123.
- Dushenkov, V., P.B.A.N. Kumar, H. Motto and I. Raskin. 1995. Rhizofiltration: the use of plants to remove heavy metals from aqueous streams. *Environ. Sci. Technol.*, 29: 1239-1245.
- Kamnev, A.A. and D. Van der Lelie. 2000. Chemical and biological parameters as tools to evaluate and improve heavy metal phytoremediation. *Biosci Rep.*, 20: 239-258.
- Kumar, P.B.A.N., V. Dushenkov, H. Motto and I. Raskin. 1995. Phytoextraction—the use of plants to remove heavy metals from soils. *Environ. Sci. Technol.*, 29: 1232-1238.
- Lin, J.X., W.S. Jiang and D.H. Liu. 2003. Accumulation of copper by roots, hypocotyls, cotyledons and leaves of sunflower (*Helianthus annuus* L.). *Bioresource Technol.*, 86: 151-155.
- Liu, D.H., W.S. Jiang and X.Z. Gao. 2003/2004. Effects of cadmium on root growth, cell division and nucleoli in root tip cells of garlic. *Biol. Plantarum*, 47: 79-83.
- Lockwood, M.P. 1976. *Effects of pollutants on aquatic organisms*. Cambridge University Press, New York.
- Lombi, E., F.J. Zhao, S.J. Dunham and S.P. McGrath. 2000. Cadmium accumulation in populations of *Thlaspi caerulescens* and *Thlaspi goesingense*. *New Phytol.*, 145: 11-20.
- Madejón, P., J.M. Murilloa, T. Marañoña, F. Cabrera and M.A. Soriano. 2003. Trace element and nutrient accumulation in sunflower plants two years after the Aznalcóllar mine spill. *Sci. Total Environ.*, 307: 239-257.
- McGrath, S.P., F.J. Zhao and E. Lombi. 2002. Phytoremediation of metals, metalloids and radionuclides. *Adv. Agron.*, 75: 1-56.
- Nordberg, G. 2003. Cadmium and human health: A perspective based on recent studies in China. *J. Trace Elem. Exp. Med.*, 16: 307-319.

- Pena, L.B., L.A. Pasquini, M.L. Tomaro and S.M. Gallego. 2006. Proteolytic system in sunflower (*Helianthus annuus* L.) leaves under cadmium stress. *Plant Sci.*, 171: 531-537.
- Salt, D.E., I.J. Pickering, R.C. Prince, D. Gleba, S. Dushenkov, R.D. Smith and I. Raskin. 1997. Metal accumulation by aquacultured seedlings of Indian mustard. *Environ. Sci. Technol.*, 31: 1636-1644.
- Salt, D.E., M. Blaylock, P.B.A.N. Kumar, V. Dushenkov, B.D. Ensley, I. Chet and I. Raskin. 1996. Phytoremediation: a novel strategy for the removal of toxic metals from the environment using plants. *Biotechnology*, 13: 468-474.
- Salt, D.E., R.C. Prince, I.J. Pickering and I. Raskin. 1995. Mechanisms of cadmium mobility and accumulation in Indian mustard. *Plant Physiol.*, 109: 1427-1433.
- Salt, D.E., R.D. Smith and I. Raskin. 1998. Phytoremediation. *Annu. Rev. Plant Physiol.*, 49: 643-648.
- Sanità, di Toppi L. and R. Gabbrielli. 1999. Response to cadmium in higher plants. *Environ. Exp. Bot.*, 41: 105-130.
- Saxena, P.K., S. KrishnaRaj, T. Dan, M.R. Perras and N.N. Vettakkorumakankav. 1999. Phytoremediation of heavy metal contaminated and polluted soils. In: *Heavy metal stress in plants. From molecules to ecosystems*. pp. 305-329. (Eds): Prasad, M.N.V. and J. Hagemeyer. Berlin: Springer-Verlag.
- Soudek, P., R. Tykva, R.Vaňková, and T. Vaněk. 2006. Accumulation of radioiodine from aqueous solution by hydroponically cultivated sunflower (*Helianthus annuus* L.). *Environ. Exp. Bot.*, 57: 220-225.
- Stephan, U.W. and Z. Prochazka. 1989. Physiological disorders of the nicotianamine-auxotroph tomato mutant *chloronerva* at different levels of iron nutrition. I. Growth characteristics and physiological abnormalities as related to iron and nicotianamine supply. *Acta Bot. Neerl.*, 38: 147-153.
- Wagner, G.J. 1994. Accumulation of cadmium in crop plants and its consequences to human health. *Adv. Agron.*, 51: 173-212.

(Received for publication 14 December 2007)