

## EVALUATION OF ZN DISTRIBUTION AMONG GRAIN AND STRAW OF TWELVE INDIGENOUS WHEAT (*TRITICUM AESTIVUM* L.) GENOTYPES

\*MUHAMMAD AAMER MAQSOOD<sup>1\*</sup>, RAHMATULLAH<sup>1</sup>, SHAMSA KANWAL<sup>1</sup>, TARIQ AZIZ<sup>2</sup> AND MUHAMMAD ASHRAF<sup>1</sup>

<sup>1</sup>Institute of Soil & Environmental Sciences, University of Agriculture, Faisalabad,

<sup>2</sup>Depalpur Sub-Campus, University of Agriculture, Faisalabad.

### Abstract

Zinc (Zn) deficiency is widespread in wheat grown on alkaline calcareous soils. A large population of the world as result of this also lacks adequate Zn nutrition. A pot study was conducted on Pindorian series (*Udic Haplustalf*) to evaluate the distribution of Zn in grain and straw of 12 indigenous wheat genotypes. Zinc was applied @ 6mg Zn kg<sup>-1</sup> soil as ZnSO<sub>4</sub>.7H<sub>2</sub>O. Various fertilizer doses were imposed in triplicate according to complete randomized design. All 72 pots received uniform dose of 60 mg N kg<sup>-1</sup> soil as urea, 60 mg P kg<sup>-1</sup> as monoammonium phosphate and 30 mg K kg<sup>-1</sup> soil as K<sub>2</sub>SO<sub>4</sub>. Plant tops were harvested 30 days after sowing. There was a significant ( $P<0.05$ ) main and interactive effect of wheat genotypes and zinc application on grain and straw yield, Zn concentration and total uptake of Zn by wheat plants. Sehar-06 produced maximum grain and straw yield whereas Iqbal-2000 produced minimum grain and straw yield. The genotypes Sehar-06, Shafaq-06 and SARC-1 were able to retain more Zn in grain compared to straw. However, further verification of the results is warranted under field conditions.

### Introduction

Wheat is an important staple food crop of the entire world as well as Pakistan. But the average yield of wheat is low due to many factors. Nutrient deficiency is one of the important factors. The universal deficiency of nitrogen and phosphorus is followed by Zn deficiency. Almost 50% of the world soils used for cereal production are Zn deficient (Gibbson, 2006). As a result, approximately 2 billion people suffer from Zn deficiency all over the world. Bhutta *et al.*, (1999) have reported Zn deficiency in Pakistan in children less than five years of age and women of reproductive age. Low soil Zn is attributed to a number of soil and environmental factors including low soil organic matter, high soil pH, calcareousness, water logging and arid climate (Cakmak, 1998; Tandon, 1995; Mortvedt, 1991). Several approaches have been investigated to overcome the common problem of Zn deficiency. These include Zn supplementation, food diversification as well as food fortification. Increasing the Zn content of food crops can be a good strategy to overcome its deficiency in people of developing countries. The Zn content of crops can be increased *via* Zn fertilization but the resource poor farmers are unable to bear the relatively high cost of Zn fertilizers.

Like many other crop species, wheat genotypes possess great variation in their Zn acquisition and utilization (Oikeh *et al.*, 2003; Banziger & Long, 2000). Most studies emphasize on the effect of Zn application on vegetative growth and grain production and not on the Zn status in grain. The objective of the present study was to compare the response of 12 indigenous wheat genotypes to Zn deficiency on a calcareous soil and to estimate the Zn status in grain.

---

\*Corresponding author's e-mail: mohamgill@hotmail.com

## Materials and Methods

**Soil:** Bulk surface soil sample (0-15 cm) was collected for Pindorian series (*Udic Haplustalf*). The sample was air-dried and ground to pass through a 2 mm sieve. A portion of the prepared soil sample was analyzed for various physico-chemical properties. Soil had pH 7.36 which was measured in 1: 1 soil: water suspension by calomel glass electrode assembly by using a Beckman pH meter. Soil texture determined by hydrometer method was sandy loam (Gee & Bauder, 1982). Organic matter content in the soil sample was 1.66 % according to Walkley-Black method (Nelson & Sommer, 1982). Free lime was 1.5% which was estimated by acid dissolution (Allison & Moodie, 1965). Plant available Zn in the soil was 0.75 mg kg<sup>-1</sup> which was extracted with 0.005 M DTPA (Lindsay & Norvell, 1978) and determined by atomic absorption spectrophotometer.

**Pot study:** Five kilogram of thoroughly mixed soil was filled each in 72 polyethylene lined plastic pots. Thirty mg N kg<sup>-1</sup> soil as urea, 45 mg P kg<sup>-1</sup> soil as monoammonium phosphate (MAP) and 30 mg K kg<sup>-1</sup> soil as K<sub>2</sub>SO<sub>4</sub> was applied uniformly to all pots in solution form. Two rates of Zn (0, 6 mg Zn kg<sup>-1</sup> soil) were applied as ZnSO<sub>4</sub>·7H<sub>2</sub>O in solution form. Before planting, soil in all the pots was moistened with distilled water, dried and thoroughly mixed for equilibration. The pots were arranged according to a completely randomized design in the green house (Steel & Torrie, 1980). During the experimental period, the average temperatures in the green house were 20±5°C at different times of the day and 12± 3 °C during the night. Light intensity varied from 300 to 1400 μmol photon m<sup>-2</sup> s<sup>-1</sup> and relative humidity varied from 35% (midday) to 85 % (midnight). Eight seeds of each of the 12 wheat genotypes were sown per pot. After germination two plants per pot were allowed to grow. Distilled water was used to maintain moisture contents at field capacity in all the pots during the experimental period. Two weeks after sowing, a second dose of 30 mg N kg<sup>-1</sup> soil as urea and 15 mg P kg<sup>-1</sup> soil as MAP were applied uniformly to all pots in solution form. Plant were harvested at maturity, washed with distilled water and blotted dried with tissue paper. The grains were separated from the straw. The straw samples were air-dried and then oven-dried at 70 °C to a constant weight for dry matter yield in a forced air oven. The straw and grain samples were then finely ground with a Wiley mill fitted with a stainless steel chamber and blades. A portion of finely ground plant and grain samples was digested in a diacid (HNO<sub>3</sub>:HClO<sub>4</sub>) mixture. The Zn concentration in the digest was estimated by atomic absorption spectrophotometer.

The data obtained for straw yield, grain yield, Zn concentration and its uptake by wheat plants were statistically analyzed using Microsoft Excel and MSTAT-C computer software (Russell & Eisensmith, 1983).

## Results

The wheat genotypes produced significantly ( $P<0.01$ ) different grain yield. The sandy loam soil had marginally available DTPA extractable Zn up to 0.75 mg kg<sup>-1</sup>. The application of Zn therefore had a significant ( $P<0.05$ ) effect on grain yield (Table 1). Maximum grain yield was produced by Sehar-06 and minimum by Iqbal-2000 by Zn application and also under control treatment. Overall there was a 9.5% increase in grain production when Zn was applied. There was a significant effect ( $P<0.05$ ) of zinc application on straw production of wheat genotypes over control (Table 1). Under adequate Zn supply maximum straw was produced by Sehar-06 (15.83g) and minimum was produced by Iqbal-2000 (11.23 g).

Table 1. Grain yield, straw yield and grain to straw ratio of 12 wheat genotypes grown at adequate and deficient zinc levels.

Wheat genotypes	Grain yield (g)		Straw yield (g)		Grain/straw ratio	
	-Zn (Control)	+Zn @ 6 $\mu\text{g g}^{-1}$	-Zn (Control)	+Zn @ 6 $\mu\text{g g}^{-1}$	-Zn (Control)	+Zn @ 6 $\mu\text{g g}^{-1}$
Inqilab-91	14.06	14.42	12.53	15.01	1.12	0.96
Bhakar-2000	14.14	12.73	12.64	13.03	1.12	0.98
Pari-73	11.72	12.92	11.03	12.91	1.06	1.00
Yaqora	14.65	15.24	10.43	12.09	1.40	1.26
As-2002	10.29	10.84	13.65	13.85	0.75	0.78
Shafaq-06	11.73	14.28	13.01	13.46	0.90	1.06
Auqab-2000	13.47	13.80	12.86	13.45	1.05	1.03
Sehar-06	14.38	15.35	14.01	15.83	1.03	0.97
Dirk	10.84	11.24	11.61	13.24	0.93	0.85
Iqbal-2000	8.26	8.73	10.38	11.23	0.80	0.78
Vatan	10.31	10.83	12.19	12.64	0.85	0.86
SARC-I	10.91	12.49	13.96	15.20	0.78	0.82
<b>Mean</b>	<b>12.06</b>	<b>12.74</b>	<b>12.36</b>	<b>13.50</b>	<b>0.98</b>	<b>0.95</b>
LSD <sub>0.05</sub> (Gentype x Zn level)	2.85		1.36		0.078	

There was a significant effect ( $P < 0.05$ ) of wheat genotypes and Zn application on the grain to straw ratio. Grain to straw ratio ranged between 1.4 to 0.78. Yaqora, Sehar-06 and Auqab-2000 had a higher grain to straw ratio compared to other genotypes.

**Zinc concentration and uptake:** There was a significant ( $P < 0.05$ ) main and interactive effect of different wheat genotypes and Zn application on concentration and uptake of Zn in wheat grain and straw (Table 2). Zinc concentration in wheat straw ranged from 29.80 to 51.22  $\mu\text{g g}^{-1}$ . An 18.1% increase in straw Zn concentration was observed by Zn application. Both under control and adequate Zn supply, Zn uptake was maximum in Sehar-6 and minimum in Iqbal-2000. There was a significant ( $P < 0.05$ ) main and interactive effect of wheat genotypes and Zn application on the grain Zn concentration and uptake (Table 2). Zinc concentration in wheat grain ranged from 34.9 to 69.93  $\mu\text{g g}^{-1}$ . Maximum grain Zn uptake was observed by Shafaq-06 under adequate Zn supply and by Sehar-06 under deficient Zn supply.

## Discussion

The genotypes exhibited wide variation for straw as well as grain biomass production. Aycicek & Yildirim, 2006 also reported differences in grain yield and yield components in different wheat genotypes. Cakmak *et al.*, (1998) and Hoffland *et al.*, (2006) reported that by Zn addition to the root medium, straw production in different crops increases. They correlated genotypic variation of crop plants to their tolerance to low Zn availability in soil. The genotypes Shafaq-06, Sehar-06 and SARC-1 showed an overall increase in grain production by Zn application over control compared to increase in straw production (Fig. 1). This attribute is an important parameter in improving nutrient content of the edible parts of plants. Total Zn uptake both in grain and straw ranged from 975.23 to 351.24  $\mu\text{g plant}^{-1}$ . Wheat grains are mostly used for human consumption. A 48.3 to 62.2% of Zn was stored in grains.

Zinc contents in straw and grain is an important parameter indicating the relative acquisition efficiencies of these genotypes from a Zn deficient soil. Efficient genotypes such as Sehar-06, Shafaq-06 and Inqalab-91 accumulated more Zn even under deficient soil Zn conditions. Zinc use efficiency is the amount of dry matter produced per unit of Zn absorbed (Siddiqui & Glass, 1981). Present results also indicated significant differences in wheat genotypes for Zn utilization efficiency. Zinc use efficiency was positively correlated with dry matter production (Fig. 2). Differences in Zn use efficiency have been reported in various crops such as wheat, rye, barley, oats (Cakmak *et al.*, 1998) and cotton (Shukla & Raj, 1987).

Relative reduction in shoot dry matter (SDM) or Zn stress factor (ZnSF) is a useful parameter in assessing relative tolerance of crop genotypes to Zn deficiency (Babikar, 1986; Yaseen, *et al.*, 2000; Irshad *et al.*, 2004). Significant differences for ZnSF (Fig. 3) were exhibited among wheat genotypes. The maximum relative reduction in SDM (%) due to Zn deficiency was exhibited by Inqalab-91 (16.5 %) while As-2002 and Bhakar-2000 showed negligible reduction in SDM.

## Conclusion

Useful genetic variations existed among wheat genotypes for Zn acquisition and its utilization to produce biomass. Cultivars efficient in Zn acquisition and utilization (Sehar-06, Shafaq-06 and Inqalab-91) showed more growth than inefficient cultivars. Efficient cultivars also translocated more Zn from the straw to the grains. Nevertheless verification of the results is needed under field conditions.

Table 2. Zinc concentration and zinc uptake in straw and grain of 12 wheat genotypes grown at adequate and deficient zinc levels.

Genotype	Zn concentration in straw ( $\mu\text{g g}^{-1}$ )		Zn concentration in grain ( $\mu\text{g g}^{-1}$ )		Total Zn uptake in straw ( $\mu\text{g plant}^{-1}$ )		Total Zn uptake in grain ( $\mu\text{g plant}^{-1}$ )	
	-Zn (Control)	+Zn @ 6 $\mu\text{g g}^{-1}$	-Zn (Control)	+Zn @ 6 $\mu\text{g g}^{-1}$	-Zn (Control)	+Zn @ 6 $\mu\text{g g}^{-1}$	-Zn (Control)	+Zn @ 6 $\mu\text{g g}^{-1}$
Inqalab-91	29.80	35.58	52.41	60.58	373.66	534.72	735.52	865.99
Bhakar-2000	41.87	44.76	54.29	59.90	433.63	583.30	761.26	753.61
Pari-73	34.73	39.32	40.00	55.82	381.57	515.12	472.63	719.91
Yaqora	33.71	40.85	51.05	59.05	351.24	497.37	746.45	899.25
As-2002	33.64	42.72	38.64	51.73	456.56	591.44	390.23	562.58
Shafaq-06	36.77	43.76	34.90	68.23	481.05	592.43	409.83	975.23
Auqab-2000	32.35	34.05	52.76	54.80	414.89	459.21	710.53	750.24
Sehar-06	34.19	51.22	54.12	57.35	585.89	809.46	778.94	883.74
Dirk	36.77	39.49	55.65	69.93	433.77	519.47	591.04	787.52
Iqbal-2000	37.96	40.68	45.61	61.60	398.17	455.36	372.82	538.43
Vatan	39.15	45.27	51.39	51.90	477.61	572.87	531.97	542.89
SARC-I	32.58	40.78	48.16	56.16	454.72	620.58	519.53	703.45
<b>Mean</b>	<b>35.29</b>	<b>41.54</b>	<b>48.25</b>	<b>58.92</b>	<b>436.90</b>	<b>562.61</b>	<b>585.06</b>	<b>748.57</b>
LSD <sub>0.05</sub> (Gentye x Zn level)	5.57		2.85		75.45		112.19	




Fig. 1. Percent increase (%) over control by Zn application in straw and grain yield of 12 wheat genotype.




Fig. 2. Correlation between shoot dry matter and Zn use efficiency of wheat genotypes.




Fig. 3. Relative reduction in shoot dry matter (ZnSF%) in 12 wheat genotypes due to Zn deficiency.

## References

- Allison, L.E. and C.D. Moodie. 1965. In: *Methods of soil analysis*, (Ed.): C.A. Black Part 2: Chemical and microbiological properties. *Am Soc. Agron.*, Madison, Wisconsin. pp. 1379-1396
- Aycicek, M. and T. Yildirim. 2006. Path coefficient analysis of yield and yield components in bread wheat (*Triticum aestivum* L.) genotypes. *Pak. J. Bot.*, 38(2): 417-424.
- Babikar, J.S.H. 1986. The effect of zinc sulfate levels on rice (*Oryza sativa* L.) growth and productivity. *Alexandria J. Agric. Res.*, 31(2): 480.
- Banziger, M. and J. Long. 2000. The potential for increasing the iron and zinc density of maize through plant breeding. *Food Nutr. Bull.*, 21(4): 397-400.
- Bhutta, Z.A., S.Q. Nizami and Z. Isani. 1999. Zinc supplementation in malnourished children with persistent diarrhoea in Pakistan. *Pediatrics*, 103(4): 1-9.
- Cakmak, I., B. Torun, B. Erenoglu, L. Ozturk, H. Marschner, M. Kalayci, H. Ekiz and A. Yilmaz. 1998. Morphological and physiological differences in the response of cereals to zinc deficiency. *Euphytica*, 100(1-3): 349-357.
- Cakmak, O., L. Ozturk, S. Karanlik, H. Ozkan, Z. Kaya and I. Cakmak. 2001. Tolerance of 65 durum wheat genotypes to zinc deficiency in a calcareous soil. *J. Plant Nutr.*, 24(11): 1831-1847.
- Gee, G.W. and J.W. Bauder. 1986. Particle-size analysis. In: *Methods of soil analysis*. Part 1: Physical and mineralogical methods. (Ed.): A. Klute. Agronomy Monogr. 9. *Soil Sci. Soc. Am.*, Madison, WI. pp. 383-409.
- Gibson, R.S. 2006. Zinc: the missing link in combating micronutrient malnutrition in developing countries. *Proceedings of the Nutrition Society*, University of East Anglia, Norwich, June 28 - July 1, 2005.
- Hoffland, E., C. Wei and M. Wissuwa. 2006. Organic anion exudation by lowland rice (*Oryza sativa* L.) at zinc and phosphorus deficiency. *Plant and Soil*, 283: 155-162.
- Irshad, M., M.A. Gill, T. Aziz, Rahmatullah and I. Ahmed. 2004. Growth response of cotton cultivars to zinc deficiency stress in chelator-buffered nutrient solution. *Pak. J. Bot.*, 36(2): 373-380.
- Lindsay, W.L. and W.A. Norvell. 1978. Development of a DTPA soil test for zinc, iron manganese and copper. *Soil Sci. Soc. Am J.*, 42(3): 421-428.
- Maftoun, M. and N. Karimian. 1989. Relative efficiency of two zinc sources for maize (*Zea mays* L.) in two calcareous soils from an arid area of Iran. *Agronomi*, 9(8): 771-775.
- Mortvedt, J.J., F.R. Cox, L.M. Shuman and R.M. Welch. 1991. In: *Micronutrients in Agriculture*, 2nd Edition. (Eds.): J.J. Mortvedt, F.R. Cox, L.M. Shuman and R.M. Welch. *Soil Sci. Soc. Am.*: Madison, WI.
- Nelson, D.W. and L.E. Sommers. 1982. Total carbon, organic carbon and organic matter. In: *Methods of soil analysis*, Part 2: Chemical and microbiological properties. *Agronomia*. Mongr. 9; *Soil Sci. Soc. Am.*: Madison, WI. pp. 570-571.
- Oikeh, S.O., A. Menkir, B. Maziya-Dixon, R. Welch and R.P. Glahn. 2003. Genotypic differences in concentration and bioavailability of kernel-iron in tropical maize varieties grown under field conditions. *J. Plant Nutr.*, 26(10&11): 2307-2319.
- Russell, D.F. and S.P. Eisensmith. 1983. *MSTAT-C*. Crop Soil Sci. Dept., Michigan State Univ. USA.
- Shukla, U.C. and H. Raj. 1987. Influence of genotypical variability on zinc response in cotton. *Plant and Soil*, 104(1): 151-154.
- Siddiqi, M.Y. and A.D.M. Glass. 1981. Utilization index: a modern approach to the estimation and comparison of nutrient utilization efficiency in plants. *J. Plant Nutr.*, 4: 289-302.
- Steel, R.G.D. and J.H. Torrie. 1980. *Principles and procedures of statistics*. McGraw-Hill, New York.
- Tandon, H.L.S. 1995. *Micronutrients in Soils, Crops and Fertilizers*. Fertilizer Development and Consultation Organization: New Delhi, India.
- Yaseen, M., R.H.N. Khan, M.A. Gill, A. Aziz, M. Aslam and A.R. Khan. 2000. Genetic variability among different rice cultivars for Zn uptake and utilization. *Pak. J. Biol. Sci.*, 3(7): 1174-11.

(Received for publication 7 November 2008)