

SURVEY OF FRESHWATER ALGAE FROM KARACHI, PAKISTAN

R. ALIYA¹, A. ZARINA² AND MUSTAFA SHAMEEL¹

¹*Department of Botany, University of Karachi, Karachi-75270, Pakistan*

²*Department of Botany, Federal Urdu University of Arts, Science & Technology,
Gulshan-e-Iqbal Campus, Karachi-75300, Pakistan.*

Abstract

Altogether 214 species of algae belonging to 86 genera of 33 families, 15 orders, 10 classes and 6 phyla were collected from various freshwater habitats in three towns of Karachi City during May 2004 and September 2005. Among various phyla, Cyanophycota was represented by 82 species (38.32%), Volvophycota by 78 species (36.45%), Euglenophycota by 4 species (1.87%), Chrysophycota by 2 species (0.93%), Bacillariophycota by 38 species (17.76%) and Chlorophycota by 10 species (4.67%). Members of the phyla Cyanophycota and Volvophycota were most prevalent (74.8%) and those of Euglenophycota and Chrysophycota poorly represented (2.8%).

Introduction

Karachi, the largest city of Pakistan is spread over a vast area of 3,530 km² and includes a variety of ponds, streams, water falls, artificial and natural water reservoirs and two small ephemeral rivers with their branchlets, which inhabit several groups of freshwater algae. Only a few studies have been carried out in the past on different groups of these algae, either from a point of view of their habitats and general occurrence (Parvaiz & Ahmed, 1981; Shameel & Butt, 1984; Aisha & Hasni, 1991; Aisha & Zahid, 1991; Leghari *et al.*, 2002) or from taxonomic viewpoint (Salim, 1954; Aizaz & Farooqui, 1972; Farzana & Nizamuddin, 1979; Ahmed *et al.*, 1983). Recently, a study was made on the occurrence of algae within Karachi University Campus (Mehwish & Aliya, 2005). Thus the available information about the algal flora of this city was not sufficient. Therefore, a large survey was initiated in May 2004 to collect and identify the freshwater algae from different areas of Karachi.

Materials and Methods

Specimens were collected from various freshwater habitats in three towns (Saddar, Jamshaid and Gulshan-e-Iqbal) of Karachi City during May 2004 and September 2005. They were brought to the laboratory and preserved in 3% formaline solution. The specimens were mounted on glycerin and semi-permanent slides were prepared by sealing the cover-slip with nail polish. They were examined under binocular light microscope and taxonomically determined with the help of authentic literature (Desikachary, 1959; Krieger & Gerloff, 1962, 1965, 1969; Prescott, 1962; Starmach, 1964, 1966; Taft & Taft, 1971; Tiwari, 1975; John *et al.*, 2002; Wehr & Sheath, 2003).

Results and Discussion

The collected specimens of freshwater algae were investigated under microscope and identified up to species level. On the basis of their morphological and cytological characteristics 214 species belonging to 86 genera of 33 families, 15 orders, 10 classes and 6 phyla were observed in three towns of Karachi City (Table 1). They have been systematically arranged according to the recently proposed classification (Shameel, 2001, 2008), therefore the new taxonomic nomenclature has been used here. The phylum Cyanophycota was found to be represented by 82 species (38.32%), Volvophycota by 78 species (36.45%), Euglenophycota by 4 species (1.87%), Chrysophycota by 2 species (0.93%), Bacillariophycota by 38 species (17.76%) and Chlorophycota by 10 species (4.67%). Members of the phyla Cyanophycota and Volvophycota were most prevalent (74.8%) and those of Euglenophycota and Chrysophycota poorly represented (2.8%). No member of the phyla Charophycota and Vaucherophycota was found to occur. These results are nearly similar with the observations made by Leghari *et al.*, (1999) at the water reservoir of Chotiari (Sanghar District, near Karachi), where 157 species of freshwater algae were recorded. Among them, 65.36% species belonged to the phyla Cyanophycota and Volvophycota. Members of the phylum Bacillariophycota were 17.5%, which is almost equal to our observation.

The phylum Cyanophycota exhibited largest diversity with 82 species. Our results are also comparable with the observations made by Leghari *et al.*, (2003) at the water reservoirs of Wah Garden (district Attock), where species of the phyla Cyanophycota and Volvophycota were dominant during summer seasons because physico-chemical factors such as high temperature, alkaline pH, high light intensity and stagnant water conditions were suitable for their excessive growth.

Gulshan-e-Iqbal Town exhibited largest number of algal flora than other two towns, although these towns included the water reservoirs of Hill Park and Askari Park. The reason may be that the large area of Karachi University Campus is also included in Gulshan-e-Iqbal Town, where several water ponds, ditches and water channels are situated in educational departments showing a luxuriant growth of algae, as has been previously observed by Mehwish & Aliya (2005). The smallest number of algal species were observed in Saddar Town, where the collections were mainly obtained from Zoological Garden. The water at most of the pools and ditches in the garden is highly polluted due to waste thrown by children and visitors, which adversely affects the growth of algae.

It was observed that the growth of microscopic species was abundant, especially in the planktonic state. The differences in the percentage of species in various towns is due to different environmental conditions like temperature, pH and growth conditions as was observed in a variety of water pools at Rawal Dam (Leghari *et al.*, 2005). Species of the phylum Chrysophycota were found in smallest proportion (0.93%), as high water temperature (25-29°C) does not appear to be suitable for their growth. Similar observations were made in the previous studies at Karachi (Leghari *et al.*, 2002; Mehwish & Aliya, 2005).

Table 1. Freshwater algae of Karachi City during May 2004-September 2005.

Algal Taxa	Sadar Town	Jamshaid Town	Gulshan-e-Iqbal Town
KINGDOM MONERA:			
PHYLUM CYANOPHYCOTA			
CLASS CHROOCOPHYCEAE			
Order Chroococcales			
Family Chroococcaceae			
<i>Aphanocapsa</i> Nägeli			
<i>A. elachista</i> W. West et G. S. West	-	-	+
<i>A. endophytica</i> G. M. Smith	-	-	+
<i>A. grevillei</i> (Hassall) Rabenhorst	-	-	+
<i>Aphanothece</i> Nägeli			
<i>A. castagnei</i> (de Brébisson) Rabenhorst	-	-	+
<i>A. gelatinosa</i> (Henning) Lemmermann	-	-	+
<i>A. microscopica</i> Nägeli	-	-	+
<i>A. prasina</i> A. Braun	-	-	+
<i>A. saxicola</i> Nägeli	-	-	+
<i>Chroococcopsis</i> Geitler			
<i>C. chroococcoides</i> (Fritsch) Komárek et Anagnostidis	-	-	
<i>Chroococcus</i> Nägeli			
<i>C. aurantiacus</i> Bernard			+
<i>C. disperses</i> (Keissler) Lemmermann	-	-	+
<i>C. gelatinosus</i> (Geitler)	-	-	+
<i>C. giganteus</i> W. West	-	-	+
<i>C. gomontii</i> Nygaard	-	-	+
<i>C. limneticus</i> Lemmermann	-	-	+
<i>C. minutus</i> (Kützing) Nägeli	-	-	+
<i>C. minor</i> (Kützing) Nägeli	-	-	+++
<i>C. montanus</i> Hansgirg	-	-	++
<i>C. pallidus</i> (Nägeli) Nägeli	-	-	++
<i>C. turicensis</i> (Nägeli) Hansgirg	-	-	+
<i>C. turgidus</i> (Kützing) Nägeli	-	-	++
<i>C. tenax</i> (Kirchner) Hieron	-	-	+
<i>C. varius</i> A. Braun	-	-	+
<i>Coelosphaerium</i> Nägeli			
<i>C. dubium</i> Grunow	-	-	+
<i>C. pallidum</i> Lemmermann			++
<i>Gloeocapsa</i> Kützing			
<i>G. aeruginosa</i> Kützing	-	-	+
<i>G. alpina</i> (Nägeli) Brand	-	-	+
<i>G. dermochroa</i> Nägeli	-	-	+
<i>G. gelatinosa</i> (Meneghini) Kützing	-	-	+
<i>G. helvetica</i> (Nägeli) Starmach	-	-	+
<i>G. haematodes</i> Kützing	-	-	+
<i>G. minuta</i> (Kützing) Hollerbach	-	-	+
<i>G. rupestris</i> Kützing	-	-	+
<i>G. siderochlamys</i> (Skuja) Starmach	-	-	+
<i>G. turgida</i> (Kützing) Hollerbach	-	-	+
<i>Gloeothece</i> Nägeli			
<i>G. linearis</i> Nägeli	-	-	+
<i>Merismopedia</i> Meyen			
<i>M. punctata</i> Meyen	-	+	+
<i>Microcystis</i> Kützing ex Lemmermann			
<i>M. aeruginosa</i> (Kützing) Kützing	-	+	+
<i>M. flosaquae</i> (Wittrock) Kirchner	-		+
<i>M. incerta</i> (Lemmermann) Lemmermann	-	+	+++
<i>M. sp.</i>	-	+	-

Table 1. (Cont'd.).

Algal Taxa	Sadar Town	Jamshaid Town	Gulshan-e-Iqbal Town
CLASS NOSTOCOPHYCEAE			
Order Nostocales			
Family Microchaetaceae			
<i>Microchaete</i> Thuret ex Bornet et Flahault			
<i>M. diplosiphon</i> Gomont ex Bornet et Flahault	-	-	++
Family Nostocaceae			
<i>Anabaena</i> Bory de Saint-Vincent ex Bornet et Flahault			
<i>A. azollae</i> Strasburger	-	-	+
Family Oscillatoraceae			
<i>Borzia</i> Cohn ex Gomont			
<i>B. trilocularis</i> Cohn ex Gomont	-	-	+
<i>Lyngbya</i> C. Agardh ex Gomont			
<i>L. aerugineocoerulea</i> Gomont	-	-	+
<i>L. hieronymusii</i> Lemmermann	-	-	+
<i>L. versicolor</i> Wartmann ex Gomont	-	-	+
<i>Oscillatoria</i> Vaucher ex Gomont			
<i>O. acutissima</i> Kufferath	-	-	+
<i>O. angustissima</i> W. West et G. S. West	-	-	+
<i>O. animalis</i> C. Agardh	-	-	+
<i>O. articulate</i> Gardner	-	-	+
<i>O. brevis</i> Kützing ex Gomont	-	-	+
<i>O. chalybea</i> Mertens	-	-	+
<i>O. curviceps</i> C. Agardh	-	-	+
<i>O. claricentrosa</i> Gardner	-	-	+
<i>O. earlei</i> Gardner	-	-	+
<i>O. formosa</i> (Bory de Saint-Vincent) Gomont	-	-	+
<i>O. minima</i> Gicklhorn	-	-	+
<i>O. pseudogeminata</i> G. Schmid	-	-	+
<i>O. quadripunctulata</i> Brühl et Biswas	-	-	+
<i>O. subbrevis</i> Schmidle	-	-	+
<i>O. tenuis</i> (C. A. Agardh) Rabenhorst	-	-	+
<i>O. sp.</i>	+	-	+
<i>Planktolynghya</i> Anagnostidis et Komárek			
<i>P. circumcreta</i> (G. S. West) Anagnostidis et Komárek	-	+	-
<i>P. contorta</i> (Lemmermann) Anagnostidis et Komárek	-	+	-
<i>P. tallingii</i> J. Komárek et H. Kling	-	+	-
<i>Pseudospirulina</i>			
<i>P. amoena</i>	-	+	-
<i>Romeria</i> Koczwara in Geitler			
<i>R. chlorine</i> Bocher	-	+	-
<i>R. gracilis</i> Koszwara	-	+	-
<i>R. leopoliensis</i> (Raciborski) Koczwara	-	+	-
Family Rivularaceae			
<i>Calothrix</i> C. Agardh ex Bornet et Flahault			
<i>C. parietina</i> (Nägeli) Thuret	-	-	+
<i>Dichothrix</i> Zanardini ex Bornet et Flahault			
<i>D. orsiniana</i> (Kützing) Bornet et Flahault	-	-	+
<i>Homoeothrix</i> (Thuret ex Bornet et Flahault) Kirchner			
<i>H. fluviatilis</i> Jao	-	-	+
<i>Phormidium</i> Kützing ex Gomont			
<i>P. ambiguum</i> Gomont	-	-	+
<i>P. luridum</i> (Kützing) Gomont	-	-	+
<i>Schizothrix</i> Kützing ex Gomont			
<i>S. muelleri</i> Nägeli ex Gomont	-	-	+
<i>Spirulina</i> Turpin ex Gomont			
<i>S. laxa</i> G. M. Smith	-	-	+

Table 1. (Cont'd.).

Algal Taxa	Sadar Town	Jamshaid Town	Gulshan-e-Iqbal Town
<i>S. laxissima</i> G. S. West	-	-	+
<i>S. major</i> Kützing	-	-	+
<i>S. princeps</i> (W. West et G. S. West) G. S. West	-	-	+
<i>S. subsalsa</i> Örsted	-	-	++
<i>S. sp.</i>	+	-	-
KINGDOM PROTISTA:			
PHYLUM VOLVOPHYCOTA			
CLASS VOLVOPHYCEAE			
Order Chlorococcales			
Family Characiaceae			
<i>Characium</i> A. Braun in Kützing			
<i>C. acuminatum</i> A. Braun	-	-	+
Family Chlorococcaceae			
<i>Chlorococcum</i> Meneghini			
<i>C. humicola</i> (Nägeli) Rabenhorst	-	-	+
<i>Tetracystis</i> R. M. Brown et Bold			
<i>T. isobilateralis</i> R. M. Brown et Bold	-	-	+
Family Coleastraceae			
<i>Coelastrum</i> Nägeli			
<i>C. pseudomicroporum</i> Korshikov	-	+	-
<i>C. verrucosum</i> (Reinsch) Reinsch	-	+	-
Family Hydrodictyaceae			
<i>Pediastrum</i> Meyen			
<i>P. boryanum</i> (Turpin) Meneghini	-	+	-
<i>P. duplex</i> Meyen	+	+	-
<i>P. simplex</i> Meyen	-	-	+++
<i>P. tetras</i> (Ehrenberg) Ralfs	++	-	-
Family Micractinaceae			
<i>Golenkinia</i> R. Chodat			
<i>G. paucispina</i> W. West et G. S. West	-	+	-
<i>G. radiata</i> Chodat	-	-	-
Family Oocystaceae			
<i>Ankistrodesmus</i> Corda			
<i>A. convolutus</i> Corda	+	++	-
<i>A. densus</i> Korshikov	-	+	-
<i>Kirchneriella</i> Schmidle			
<i>K. elongata</i> G. M. Smith	-	-	+
<i>K. lunaris</i> (Kirchner) K. Möbius	-	-	-
<i>K. obesa</i> (G. S. West) Schmidle	-	+	++
<i>Lauterborniella</i> W. Schmidle			
<i>L. elegantissima</i> W. Schmidle	-	+	-
<i>Nephrocytium</i> Nägeli			
<i>N. agardhianum</i> Nägeli	+	-	-
<i>N. lunatum</i> West	+	-	-
<i>Oocystis</i> A. Braun			
<i>O. borgei</i> J. Snow	-	+	-
<i>O. marssonii</i> Lemmermann	-	+	-
<i>Schroederia</i> Lemmermann			
<i>S. setigera</i> (Schröder) Lemmermann	-	+	-
<i>Sorastrum</i> Kützing			
<i>S. americanum</i> (Bohlin) Schmidle	-	+	-
<i>S. spinulosum</i> Nägeli	-	+	-
<i>Tetraedron</i> Kützing			
<i>T. caudatum</i> (Corda) Hansgirg	-	-	+
<i>T. pentaedricum</i> W. West et G. S. West	-	-	+

Table 1. (Cont'd.).

Algal Taxa	Sadar Town	Jamshaid Town	Gulshan-e-Iqbal Town
<i>T. minimum</i> (A. Braun) Hansgirg	-	-	+
<i>T. muticum</i> (A. Braun) Hansgirg	-	++	-
<i>T. quadratum</i> (Reinsch) Hansgirg	+	-	+
Treubaria C. Bernad			
<i>T. crassispina</i> G. M. Smith	-	-	+
Westella De Wildermann			
<i>W. linearis</i> G. M. Smith	+	-	-
Family Scenedesmaceae			
Scenedesmus Meyen			
<i>S. abundans</i> (O. Kirchner) Chodat	-	+	-
<i>S. acuminatus</i> (Lagerheim) Chodat	-	+	++
<i>S. acutiformis</i> Schröder	-	+	+
<i>S. acutus</i> Meyen	-	+	-
<i>S. arcuatus</i> Lemmermann	-	-	+
<i>S. bijuga</i> (Reinsch.) Hansgirg	-	-	+
<i>S. circumfusus</i> Hortobagyi	-	+	-
<i>S. dimorphus</i> (Turpin) Kützing	+	++	+
<i>S. denticulatus</i> Lagerheim	+	-	-
<i>S. falcatus</i> Chodat	-	+	-
<i>S. hystrix</i> Lagerheim	-	-	+
<i>S. incrassatulus</i> Bohlin	+	-	-
<i>S. obliquus</i> (Turpin) Kützing	-	+	-
<i>S. perforatus</i> Lemmermann	-	-	+
<i>S. quadricauda</i> (Turpin) de Brébisson	+	-	++
<i>S. raciborskii</i> Wolozynska	-	+	-
<i>S. semipulcher</i> Hortobagyi	-	+	-
<i>S. sp.</i>	-	+	-
Tetrastrum Chodat			
<i>T. glabrum</i> (Y. V. Roll) Ahlstrom et Tiffany	-	+	-
<i>T. heteracanthum</i> (Nordstedt) Chodat	+	-	-
Order Chlorosarcinales			
Family Chlorosarcinaceae			
Chlorosarcinopsis W. Hernodon			
<i>C. minuta</i> R. D. Groover et H. C. Bold	-	-	+
Family Palmellaceae			
Chloranomala Mitra			
<i>C. cuprecola</i> Mitra	-	-	+
Gloeocystis Nägeli			
<i>G. major</i> Gerneck ex Lemmermann	-	-	+
Order Volvocales			
Family Volvocaceae			
Pandorina Bory de Saint-Vincent			
<i>P. morum</i> (O. F. Müller) Bory de Saint-Vincent	-	-	+
Order Tetrasporales			
Family Coccomyxaceae			
Diogenes W. Pennington			
<i>D. bacillaris</i> (Naum.) W. Pennington	-	-	+
Dispora H. Printz			
<i>D. crucigenioides</i> H. Printz	-	-	+
Family Gloeocystaceae			
Palmellopsis O. Korshikov			
<i>P. gelatinosa</i> O. Korshikov	-	-	+

Table 1. (Cont'd.).

Algal Taxa	Sadar Town	Jamshaid Town	Gulshan-e-Iqbal Town
CLASS DESMIDOPHYCEAE			
Order Desmiales			
Family Desmidiaceae			
<i>Arthrodesmus</i> C. G. Ehrenberg ex Ralfs			
<i>A. fuellebornei</i> Schmidle	-	-	+
<i>Closterium</i> Nitzsch ex Ralfs			
<i>C. moniliferum</i> Ehrenberg	-	-	+
<i>C. parvulum</i> Nägeli	-	-	+
<i>Cosmarium</i> Ralfs			
<i>C. angulosum</i> de Brébisson	-	+	-
<i>C. arctoum</i> Nordstedt et Krieger	-	+	-
<i>C. leave</i> Rabenhorst	-	+	-
<i>C. geometricum</i> W. West et G. S. West	-	+	-
<i>C. margaritatum</i> (Lund) Roy et Bisset	-	-	+
<i>C. phaseolus</i> de Brébisson ex Ralfs	-	+	-
<i>C. pseudoprotuberans</i> Krichner	-	+	-
<i>C. sulcatum</i> Nordstedt	-	+	-
<i>Eurastrum</i> C. G. Ehrenberg ex Ralfs			
<i>E. pectinatum</i>	-	-	+
<i>Staurastrum</i> (Meyen) Ralfs			
<i>S. anatinum</i> Cooke et Wills	-	-	+
<i>S. cyrtocerum</i> Brébisson	-	+	-
<i>S. leptocladum</i> L. N. Johanson	-	-	+
<i>S. manfeldtii</i> Deleponte	-	-	+
<i>S. muticum</i> de Brébisson ex Ralfs	-	-	+
<i>S. sebalzii</i> Nordstedt	-	+	-
<i>S. taylorii</i> R. L. Grönblad	-	+	-
<i>S. sp.</i>	-	+	-
PHYLUM EUGLENOPHYCOTA			
CLASS EUGLENOPHYCEAE			
Order Euglenales			
Family Euglenaceae			
<i>Euglena</i> Ehrenberg			
<i>E. spirogyra</i> Ehrenberg	-	-	+
<i>E. sp.</i>	-	++	-
<i>Phacus</i> Dujardin			
<i>P. curvicauda</i> Swirenko	+	-	-
<i>P. orbicularis</i> K. Hübner	+	-	-
PHYLUM CHRYSOPHYCOTA			
CLASS CHRYSOPHYCEAE			
Order Ochromonales			
Family Chrysocapsaceae			
<i>Chrysocapsa</i> Pascher			
<i>C. planctonica</i> (West et West) Pascher	-	-	+
Family Synuraceae			
<i>Mallomonas</i> Perty			
<i>M. caudata</i> Iwanoff	-	-	+
PHYLUM BACILLAROPHYCOTA			
CLASS BACILLAROPHYCEAE			
Order Bacillariales			
Family Achnantheaceae			
<i>Achnanthes</i> Bory de Saint-Vincent			
<i>A. linearis</i> W. Smith	-	-	++

Table 1. (Cont'd.).

Algal Taxa	Sadar Town	Jamshaid Town	Gulshan-e-Iqbal Town
Cocconeis Ehrenberg			
<i>C. pediculus</i> Ehrenberg	-	-	+
Family Cymbellaceae			
Amphora Ehrenberg ex Kützing			
<i>A. coffeaformis</i> Agardh	-	-	+
<i>A. ovalis</i> Kützing	-	-	+
<i>A. sp.</i>	-	-	+
Cymbella C. A. Agardh			
<i>C. cistula</i> (Hempr.) Kirchner	+	-	+
<i>C. ventricosa</i> Kützing	-	+	-
<i>C. sp.</i>	-	+	+
Family Epithemaceae			
Denticula Ehrenberg			
<i>D. elegans</i> Kützing	-	-	+
Epithemia de Brébisson			
<i>E. ocellata</i> Kützing	-	+	-
<i>E. sorex</i> Kützing	-	+	-
Rhopalodia O. F. Müller			
<i>R. gibba</i> (Ehrenberg) O. Müller	-	-	+
<i>R. sp.</i>	-	-	++
Family Fragilaraceae			
Ceratonei Ehrenberg			
<i>C. arcus</i> (Hempr.) Kirchner	-	-	+
Fragilaria Lyngbye			
<i>F. crotonensis</i> Kitt.	-	+	-
<i>F. alpestris</i> Krasske	-	+	-
<i>F. ulna</i> Nitzsch) Lange-Bertalot	+	+	-
Fragilariforma (J. Ralfs) D. M. Williams et F. E. Round			
<i>F. virescens</i> (J. Ralfs) D. M. Williams et F. E. Round	-	+	-
<i>F. sp.</i>	-	+	+
Grammatophora Ehrenberg			
<i>G. serpentine</i> Ralfs	-	-	++
Synedra Ehrenberg			
<i>S. acus</i> Ehrenberg	+	-	-
<i>S. minuscula</i> Grunow	+	+	-
Family Naviculaceae			
Anomoeoneis Pfitzer			
<i>A. sphaerophora</i> (Kützing) Pfitzer	-	-	+
Diatomella Greville			
<i>D. sp.</i>	-	-	+
Mastogloia Thwaites ex W. Smith			
<i>M. danseii</i> Thwait	-	-	+
<i>M. smithii</i> Thwait	-	-	+
Navicula Bory emend. Cleve			
<i>N. anglica</i> Ralfs	-	+	-
<i>N. cocconeiformis</i> Gregory	-	+	-
<i>N. platystoma</i> Ehrenberg	-	+	-
<i>N. viridis</i> Kützing	-	-	+
<i>N. viridula</i> (Kützing) Kützing emend. Van Heurck	-	-	+
<i>N. sp.</i>	-	+	-

Table 1. (Cont'd.).

Algal Taxa	Sadar Town	Jamshaid Town	Gulshan-e-Iqbal Town
<i>Pinnularia</i> (Ehrenberg) Ehrenberg			
<i>P. major</i> Kützing	-	+	-
<i>P. sp.</i>	-	+	-
Family Nitzschaceae			
<i>Gyrosigma</i> Hassall			
<i>G. sp.</i>	-	-	+
<i>Nitzchia</i> Hassall			
<i>N. sp.</i>	-	-	+
Order Biddulphales			
Family Coscinodiscaceae			
<i>Cyclotella</i> (Kützing) de Brébisson			
<i>C. glomerata</i> Bochmann	-	-	+
<i>C. sp.</i>	+	-	+
KINGDOM PHYCOTA:			
PHYLUM CHLOROPHYCOTA			
CLASS ULVOPHYCEAE			
Order Ulotrichales			
Family Cylirocapsaceae			
<i>Cylindrocapsa</i> Reinsch			
<i>C. conferta</i> W. West	-	-	+
<i>Gloetotanium</i> Itasgr			
<i>G. ioilesbergereanum</i> Hasgird	-	-	+
Family Ulotrichaceae			
<i>Geminella</i> Turpin emend. Lagerheim			
<i>G. mutabilis</i> (de Brébisson) Wille	-	-	+
<i>Ulothrix</i> Kützing			
<i>U. sp.</i>	+	-	-
CLASS ZYGNEMOPHYCEAE			
Order Oedogonales			
Family Oedogonaceae			
<i>Oedogonium</i> Link			
<i>O. grande</i> Kützing	-	-	+
Order Zygnemales			
Family Zygnemaceae			
<i>Spirogyra</i> Link			
<i>S. longata</i> (Vaucher) Kützing	-	-	+
<i>Zygnemopsis</i> (Skuja) Transeau			
<i>Z. desmidioides</i> (W. West et G. S. West) Transeau	-	-	+
CLASS SIPHONOCALADOPHYCEAE			
Order Cladophorales			
Family Cladophoraceae			
<i>Pithophora</i> Wittrock			
<i>P. oedogonia</i> (Montagne) Wittrock	-	+	-
<i>Rhizoclonium</i> Kützing			
<i>R. crassipellitum</i> W. West et G. S. West	-	-	++
<i>R. hookeri</i> Kützing	-	-	++

+ = Common, ++ = Frequent, +++ = Dominant, - = Absent.

References

- Ahmed, J., M. Begum and S. Begum. 1983. Survey and cultivation of freshwater algae from Karachi and its suburbs. *Biblioth. Phycol.*, 66: 389-445.
- Aisha, K. and S. Hasni. 1991. Occurrence of *Enteromorpha* Link (Chlorophyceae) in freshwater of Pakistan. *Pak. J. Bot.*, 23: 143.
- Aisha, K. and P.B. Zahid. 1991. Occurrence of algae in industrial waste water around Karachi. *Pak. J. Bot.*, 23: 155-159.
- Aizaz, I.G. and P.B. Farooqui. 1972. On the morphology and development of some Chlorococcales from Karachi. *Pak. J. Bot.*, 23: 171-181.
- Desikachary, T.V. 1959. *Cyanophyta*. I. C. A. R., New Delhi, 686 pp.
- Farzana, A. and M. Nizamuddin. 1979. Studies on some members of Cyanophyta from Karachi. *Nova Hedw.*, 31: 247-256.
- John, D.M., B.A. Whitton and A.J. Brook. 2005. *The Freshwater Algal Flora of the British Isles*. Camb. Univ. Press, Cambridge. 702 pp.
- Krieger, W. and J. Gerloff. 1962. Die Gattung *Cosmarium*. Weinheim, 1: 1-112.
- Krieger, W. and J. Gerloff. 1965. Die Gattung *Cosmarium*. Weinheim, 2: 113-140.
- Krieger, W. and J. Gerloff. 1969. Die Gattung *Cosmarium*. Weinheim & Lehre, 1-4: 241-410.
- Leghari, S.M., S.I.H. Jafri, M.A. Mahar, K.H. Lashari, S.S. Ali, M.Y. Khuhawar and T.M. Jahangir. 1999. Biodiversity of Chotiari Reservoir, distt. Sanghar, Sindh, Pakistan. *In: Aquatic Biodiversity of Pakistan*. p. 139-157.
- Leghari, S.M., T.M. Jahangir, M.Y. Khuhawar and A. Leghari. 2002. Study on the natural springs at Clifton, Karachi, Sindh, Pakistan. *Proc. Pak. Congr. Zool.*, 22: 125-131.
- Leghari, M.K, M.Y. Leghari, M. Shah and S.N. Arbani. 2003. Ecological study of algal flora of Wah Garden, district Attock, Pakistan. *Pak. J. Bot.*, 35: 705-716.
- Leghari, M.K, M. Shah and M.Y. Leghari. 2005. Ecological study of blue-green algae from Rawal Dam Water, Islamabad. *Int. J. Phycol. Phycochem.*, 1: 5-12.
- Mehwish, H. and R. Aliya. 2005. Occurrence of freshwater algae at different localities of Karachi University. *Int. J. Phycol. Phycochem.*, 1: 117-124.
- Parvaiz, K. and J. Ahmed. 1981. A preliminary survey of sewage algae from Karachi. *Pak. J. Bot.*, 13: 221-223.
- Prescott, G.W. 1962. *Algae of the Western Great Lakes Area*. 2nd ed., Wm C. Brown Co., Dubuque, Iowa, 977 pp.
- Salim, K.M. 1954. Some planktonic diatoms from Karachi. *Pak. J. Sci.*, 6: 106-110.
- Shameel, M. 2001. An approach to the classification of algae in the new millennium. *Pak. J. Mar. Biol.*, 7: 233-250.
- Shameel, M. 2008. Change of divisional nomenclature in the Shameelian classification of algae. *Int. J. Phycol. Phycochem.*, 4: 225-232.
- Shameel, M. and N.I. Butt. 1984. On the occurrence of Cyanophyta from Karachi, Pakistan. *Pak. J. Bot.*, 16: 75-79.
- Starmach, K. 1964. Chrysophyta. II. Bacillariophyta-Okrzemki. *Flora Slodkow Polski*, 6: 1-610.
- Starmach, K. 1966. Cyanophyta-Sinice, Glaucophyta-Glaukofity. *Flora Slodkow Polski*, 2: 1-808.
- Taft, C.E. and C.W. Taft. 1971. *The Algae of the Western Lake Erie*. Ohio State Univ., Columbus, 4: 1-189.
- Tiwari, G.L. 1975. A study of the blue green algae from paddy field soils of India. II. Taxonomic consideration of non heterocystous blue green algae. *Nova Hedw.*, 26: 765-798.
- Wehr, J.D. and R.G. Sheath. 2003. *Freshwater Algae of the North America: Ecology and Classification*. Acad. Press, London, 918 pp.

(Received for publication 30 December 2008)