

RELATIONSHIPS BETWEEN SODIUM ION ACCUMULATION AND PHYSIOLOGICAL CHARACTERISTICS IN RICE (*ORYZA SATIVA* L. SPP. *INDICA*) SEEDLINGS GROWN UNDER ISO-OSMOTIC SALINITY STRESS

KONGAKE SIRINGAM¹, NIRAN JUNTAWONG¹, SURIYAN CHA-UM^{2*}
AND CHALERMPOL KIRDMANEE²

¹*Department of Botany, Faculty of Sciences, Kasetsart University, Chatuchak, Bangkok, 10900, Thailand*

²*National Center for Genetic Engineering and Biotechnology, National Science and Technology Development Agency (NSTDA), Klong Luang, Pathumthani, 12120, Thailand.*

**Corresponding author: suriyanc@biotec.or.th*

Abstract

The objective of this research was to elucidate the role of sodium ion (Na^+) on photosynthetic machinery and growth characteristics in both salt-tolerant (HJ) and salt-sensitive (PT1) rice varieties grown under iso-osmotic salinity stress. The Na^+ and Na^+/K^+ in HJ and PT1 rice seedlings were increased with increasing salt concentrations in the culture media. K^+ in salt-stressed HJ seedlings was increased but that in salt-stressed PT1 seedlings was unchanged. The Na^+ accumulation in salt-stressed seedlings was negatively related to the water potential in HJ ($r^2 = 0.78$) and PT1 ($r^2 = 0.72$), leading to pigment degradation i.e. chlorophyll a (Chl_a), chlorophyll b (Chl_b) and total carotenoids (C_{x+c}). The water potential in both HJ and PT1 salt-stressed seedlings was positively related to the Chl_a concentration ($r^2 = 0.78$ and $r^2 = 0.58$). Furthermore, Chl_a and total chlorophyll (TC) concentrations in HJ and PT1 salt-stressed seedlings were positively related to water oxidation in PSII (PSII) as maximum quantum yield of PSII (F_v/F_m) ($r^2 = 0.73$ and $r^2 = 0.68$) and quantum efficiency of PSII (Φ_{PSII}) ($r^2 = 0.92$ and $r^2 = 0.91$), respectively. The Φ_{PSII} in HJ and PT1 salt-stressed seedlings was positively related to the dry matter ($r^2 = 0.86$ and $r^2 = 0.67$). The K^+ accumulation in HJ salt-stressed seedlings may play a major role in salt defense mechanisms, leading to enhance on photosynthesis capacity, water oxidation in photosystem II (PSII) and growth abilities.

Introduction

Salinity stress is the major factor to limit plant growth and productivity in many areas of the world (Allakhverdiev *et al.*, 2000; Ashraf, 2009). Excessive amounts of salts in the soil solution cause both osmotic and ionic stresses (Khelil *et al.*, 2007). These stresses are generated from the major salt contaminated soil, namely NaCl, which is readily dissolved in the water to give rise Na^+ and Cl^- (Taiz & Zeiger, 2002). The Na^+ is generally reported as a strong cation to interact with other cations, especially K^+ (Hu *et al.*, 2007; Aziz & Khan, 2001; Ashraf, 2004) in the membrane system, since Na^+/K^+ exchanges for intracellular influx to increase the Na^+/K^+ selectivity (Parida & Das, 2005). Enhanced membrane permeability in salt stressed plants has been well documented (Hasegawa *et al.*, 2000), resulting in Na^+ accumulation and increasing Na^+/K^+ , conversely reducing K^+ in many plant species i.e. rice (Dionisio-Sese & Tobita, 1998), green bean (Yasar *et al.*, 2006), winter wheat (Zheng *et al.*, 2008), umbu plant (da Silva *et al.*, 2008).

Furthermore, a deteriorated membrane affects on the reduction of the physiological and biochemical processes (Taiz & Zeiger, 2002; Noreen & Ashraf, 2008), especially photosynthesis. Photosynthetic pigments are necessary for photosynthesis mechanism

including; chlorophyll a (Chl_a), chlorophyll b (Chl_b), total chlorophyll (TC) and total carotenoids (C_{x+c}) which play a role in the light-harvesting complex of photosystem II (PSII). Chl_a, Chl_b and TC in higher plants play an important role on the water oxidation and electron transport in photosystem II. In addition, C_{x+c} play a key role for antioxidative functions to protect the plants from the reactive oxygen species (ROS), which generally generate by salinity stress. Under salinity stress, the plasma membrane and activities of antioxidant enzymes are regulated by Na⁺ accumulation to prevent the damage of chloroplasts, which contain photosynthetic pigments such as Chl_a, Chl_b, TC and C_{x+c} (Demiral & Türkan, 2005). The chloroplast is very sensitive to damage both the properties and efficiencies in the light-harvesting complexes, leading to disturb in the chloroplast ultrastructure (Zheng *et al.*, 2008). Many reports show salt induced reduction in photosynthetic pigments in many plant species i.e. rice (Cha-um *et al.*, 2007a), castor bean (Pinheiro *et al.*, 2008), raspberry (Neocleous & Vasilakakis, 2007).

The efficiency of photosynthetic pigments is indicated by the function of chlorophyll pigments or chlorophyll a fluorescence, which demonstrate the functioning of water oxidation and photosynthetic electron transport in photosystem II (PSII) under stress condition (Gray *et al.*, 1997). The photosynthetic pigments in the chloroplast are composed of light harvesting complexes namely photosynthetic apparatus. Chlorophyll a fluorescence parameters such as maximum quantum yield (F_v/F_m), photosystem II quantum efficiency (Φ_{PSII}), photochemical quenching (qP) and non-photochemical quenching (NPQ) have been investigated (Maxwell & Johnson, 2000) and applied as effective indices for abiotic stress screening. There are many reports which show the reduction of chlorophyll a fluorescence in salt-stressed plant species i.e., rice (Cha-um *et al.*, 2007a), sunflower (Santos, 2004), Rumex (Chen *et al.*, 2004) and citrus (Lopez-Climent *et al.*, 2008).

Rice (*Oryza sativa* L. spp. *indica*) is a staple food for feeding and providing the necessary daily calories for three billions of people in the world (Kush, 1997). However, the growth and yield are dramatically reduced by abiotic extreme conditions, especially salinity stress, since the rice has been reported as salinity susceptible crop (Yokoi *et al.*, 2002; Flowers & Yeo, 1995). From some previous studies, the results showed that rice growth and development are reduced when exposed to the salinity stress (Morsy *et al.*, 2007; Cha-um *et al.*, 2007a; Dionisio-Sese & Tobita, 1998; Shannon *et al.*, 1998). However, the response of rice to salt stress depends on its salt tolerance abilities. The objective of this study was to investigate the role of Na⁺/K⁺ on physiological responses of salt-tolerant and salt-sensitive rice varieties grown under iso-osmotic salinity stress.

Materials and Methods

Plant materials and treatments: Seeds of salt-tolerant (HJ) and salt-sensitive (PT1) varieties (Cha-um *et al.*, 2007b) of indica rice (*Oryza sativa* L. spp. *indica*) were obtained from Rice Research Institute, Pathumthani Rice Research Center, Pathumthani, Thailand. Rice seeds were dehusked, disinfected once in 5% (v/v) Clorox[®] [5.25% (w/v) sodium hypochlorite solution, Clorox Co. Ltd., USA] for 12 h, once in 25% (v/v) Clorox[®] for 30 min., and then rinsed thrice with sterilized distilled water. Surface sterilized seeds were germinated on 25 mL MS-gelled (Murashige & Skoog, 1962) with 3% sucrose (photomixotrophic condition) in a 250 mL glass vessel. The medium pH was adjusted to 5.7 before autoclaving. *In vitro* rice seedlings were cultured in a culture room under conditions of 25±2°C air temperature, 60±5% relative humidity (RH), 60±5 μmol m⁻² s⁻¹ photosynthetic photon flux density (PPFD) provided by fluorescent lamps (TLD 36 W/84 Cool White 3350 Im, Philips, Thailand) for 16 h d⁻¹ photoperiod. Fourteen-day-old

seedlings were aseptically transferred to MS sugar-free liquid media (photoautotrophic condition) by using vermiculite as a supporting material for 7 days. Air-exchange rate in the glass vessels was adjusted to 2.32 h^{-1} by punching a hole in the plastic cap ($\varnothing 1 \text{ cm}$) and covering the hole with a gas-permeable microporous polypropylene film ($0.22 \mu\text{m}$ pore size, Nihon Millipore Ltd., Japan). The culture medium was adjusted to 0.0, 85.5, 171.0, 256.5 or 342.0 mM NaCl with iso-osmotic level using mannitol solution for controlling the range of water potential in the medium ($-1.4 \pm 0.2 \text{ MPa}$) for 4 days.

Ion contents: One-hundred-milligrams fresh weight of leaf tissues were ground in liquid-nitrogen and extracted by acidic method. Na^+ and K^+ concentrations were measured by an atomic absorption spectrophotometer (AA; Model M6 Thermo Elemental, MA, USA) according to Dionisio-Sese & Tobita method (1998).

Photosynthetic pigment concentrations: Chlorophyll a (Chl_a), chlorophyll b (Chl_b), total chlorophyll (TC) and total carotenoids (C_{x+c}) concentrations were determined following Shabala *et al.*, (1998) and Lichtenthaler (1987) methods, respectively. One-hundred-milligram leaf tissues were placed in a glass vessel, added 10 mL of 95.5% acetone and blended with a homogenizer (T25 Basic ULTRA-TURRAX[®]; IKA, Kuala Lumpur, Malaysia). The glass vessels were sealed with parafilm to prevent evaporation and then stored at 4°C for 48 h. The Chl_a , Chl_b , TC and C_{x+c} concentrations were measured using a UV-visible spectrophotometer (DR/4000; HACH, Loveland, Colorado, USA) at 662, 644 and 470 nm. A solution 95.5% acetone was used as a blank. The Chl_a , Chl_b , TC and C_{x+c} ($\mu\text{g g}^{-1} \text{ FW}$) concentrations in the leaf tissues were calculated according to the following equations:

$$\begin{aligned} \text{Chl}_a &= 9.784D_{662} - 0.99D_{644} \\ \text{Chl}_b &= 21.42D_{644} - 4.65D_{662} \\ \text{TC} &= \text{Chl}_a + \text{Chl}_b \\ C_{x+c} &= \frac{1000D_{470} - 1.90\text{Chl}_a - 63.14\text{Chl}_b}{214} \end{aligned}$$

where D_i is an optical density at the wavelength i .

Chlorophyll a fluorescence measurement: Chlorophyll a fluorescence emission of adaxial leaf surface was monitored by a Fluorescence Monitoring System (FMS 2; Hansatech Instruments Ltd, Norfolk, UK) in the pulse amplitude modulation mode, as previously described by Loggini *et al.*, (1999). The maximum quantum yield of PSII photochemistry (F_v/F_m), quantum efficiency of PSII (Φ_{PSII}), photochemical quenching (qP) and non-photochemical quenching (NPQ) were evaluated by FMS software for Windows (Fluorescence Monitoring System Software; Hansatech Instruments Ltd), and were calculated as described by Maxwell & Johnson (2000).

Growth abilities: Fresh weight was measured after exposing salt stress for 4 days. The plant sample was then dried at 110°C in a hot-air oven (Memmert, Model 500, Germany) for 48 h and then incubated in a desiccator before measuring the dry weight. Dry matter was calculated with the following equation:

$$\text{Dry matter (\%)} = \frac{\text{Dry weight}}{\text{Fresh weight}} \times 100$$

Statistical analysis of data: The experiment was designed as 2×5 factorials in a Completely Randomized Design (CRD) with four replications and five seedlings per replicate. Significant level was determined by one-way analysis of variance (ANOVA) using the SPSS software (SPSS for Windows, SPSS Inc., USA). Differences between means were compared by the Duncan's Multiple Range Test (DMRT). The relations between Na⁺ accumulation and water potential, water potential and Chl_a, Chl_a and F_v/F_m, TC and Φ_{PSII}, Φ_{PSII} and dry matter were evaluated.

Results and Discussions

Sodium ion (Na⁺) accumulation in salt-tolerant (HJ) and salt-sensitive (PT1) rice seedlings was directly enhanced, relating to NaCl salt concentrations (85.5, 171.0, 256.5 and 342.0 mM NaCl) in the culture medium (Table 1). The Na⁺ in HJ seedlings cultured on 85.5, 171.0, 256.5 and 342.0 mM NaCl enhanced for 23.1, 68.0, 88.5 and 146.9 folds when compared to those on without NaCl. In a parallel way, the Na⁺ in PT1 salt-stressed seedlings was accumulated 6.5, 21.3, 24.0 and 28.4 folds when compared to those cultured on that without NaCl (Table 1). Moreover, the Na⁺ accumulation in HJ and PT1 salt-stressed seedlings was negatively related to water potential ($r^2 = 0.78$ and $r^2 = 0.72$) (Fig. 1A and 1B). In contrast, potassium ion (K⁺) in HJ salt-stressed seedlings was significantly accumulated, depending on NaCl salt concentrations in the media. The K⁺ in HJ seedlings cultured on 85.5, 171.0, 256.5 and 342.0 mM was 2.8, 2.7, 3.3 and 4.4 folds higher than those cultured on without NaCl. On the other hand, K⁺ accumulation in PT1 seedlings was not significantly changed when exposed to NaCl stress (Table 1). In addition, Na⁺/K⁺ ratio in HJ and PT1 salt-stressed was continuously increased with increasing the NaCl concentrations (Table 1). The Na⁺/K⁺ ratio in HJ seedlings cultured on 85.5, 171.0, 256.5 and 342.0 mM NaCl was higher than those cultured on without NaCl for 8.4, 25.3, 27.3 and 35.2 folds, respectively. As well as the Na⁺/K⁺ ratio in PT1 salt-stressed seedlings was higher than those cultured on without NaCl for 7.2, 18.9, 26.7 and 25.8 folds (Table 1).

Na⁺ accumulation and Na⁺/K⁺ ratio in salt-stressed plants depend on salt stress treatments, while K⁺ accumulation is generally decreased (Parida & Das, 2005; Parida *et al.*, 2004; Levigneron *et al.*, 1995; Sabir & Ashraf, 2007). The Na⁺ and K⁺ enrichment and Na⁺/K⁺ ratio have been utilized to identify the salt tolerant ability in many plant species such as rice (Zeng, 2005; Nakamura *et al.*, 2002; Dionisio-Sese & Tobita, 1998), winter wheat (Zheng *et al.*, 2008) umbu plant (da Silva *et al.*, 2008), *Arbutus unedo* (Navarro *et al.*, 2007) and maize (Çiçek & Cakirlar, 2002). In this study, the Na⁺ and K⁺ accumulation and Na⁺/K⁺ ratio in salt-tolerant HJ seedlings was increased with increasing iso-osmotic salinity stress, while the K⁺ in PT1 salt-stressed seedlings remained unchanged. The K⁺ accumulation in HJ rice seedlings grown under iso-osmotic salinity stress may play a role as salt defense mechanism when compared to that in PT1 salt sensitive seedlings. There are many reports, which show an important role of K⁺ in different physiological processes such as osmotic adjustment, ionic balance and ion homeostasis (Rejili *et al.*, 2007; Nguyen *et al.*, 2005; Serrano & Navarro, 2001). In addition, an increase in K⁺ is well established (Yasar *et al.*, 2006; Chow *et al.*, 1990), which leads to reduced Na⁺ uptake in the cells (Parida & Das, 2005; Saqib *et al.*, 2005) thereby maintaining low [Na⁺/K⁺] ratio in plant tissues exposed to salt stress (Rejili *et al.*, 2007; Nguyen *et al.*, 2005).

Table 1. Sodium ion (Na⁺), potassium ion (K⁺) and Na⁺:K⁺ ratio in leaves of salt-tolerant (HJ) and salt-sensitive (PT1) rice varieties cultured on MS medium and exposed to 0.0, 85.5, 171.0, 256.5 and 342.0 mM NaCl with iso-osmotic level using mannitol application for 4 days.

Rice varieties	NaCl (mM)	Na ⁺ (mg g ⁻¹ FW)	K ⁺ (mg g ⁻¹ FW)	Na ⁺ :K ⁺
HJ	0.0	0.30 f	22.82 e	0.013 f
	85.5	6.92 de	64.15 bc	0.109 ef
	171.0	20.36 c	62.00 bc	0.328 cde
	256.5	26.51 c	74.72 b	0.355 cd
	342.0	44.01 a	99.27 a	0.456 bc
PT1	0.0	1.17 ef	43.04 d	0.027 f
	85.5	7.58 d	39.26 de	0.193 def
	171.0	24.71 c	48.44 cd	0.511 abc
	256.5	22.85 c	37.37 de	0.630 ab
	342.0	33.04 b	48.01 cd	0.696 a
Significant level				
Rice varieties		ns	**	**
NaCl		**	**	**
Rice varieties × NaCl		*	**	ns

Means with the different letters are significantly different at $p \leq 0.01$ (**), $p \leq 0.05$ (*) and non-significant (ns) by Duncan's New Multiple Range Test.




Fig. 1. Relationships between Na⁺ accumulation (mg g⁻¹ FW) and water potential (MPa) of salt-tolerant (A) and salt-sensitive (B) rice seedlings cultured *In vitro* and exposed to salt stress with iso-osmotic level using mannitol application for 4 days.

Table 2. Chlorophyll a (Chl_a), chlorophyll b (Chl_b), total chlorophyll (TC) and total carotenoid (C_{x+c}) concentrations of salt-tolerant (HJ) and salt-sensitive (PT1) rice varieties cultured on MS medium and exposed to 0.0, 85.5, 171.0, 256.5 or 342.0 mM NaCl with iso-osmotic level using mannitol application for 4 days.

Rice varieties	NaCl (mM)	Chl _a (µg g ⁻¹ FW)	Chl _b (µg g ⁻¹ FW)	TC (µg g ⁻¹ FW)	C _{x+c} (µg g ⁻¹ FW)
HJ	0.0	320 b	661 b	905 b	40.1 a
	85.5	260 d	529 c	802 c	36.4 b
	171.0	239 e	451 e	702 d	29.7 d
	256.5	232 e	440 ef	700 d	29.2 d
	342.0	213 f	413 f	632 e	26.5 e
PT1	0.0	375 a	698 a	967 a	37.2 b
	85.5	289 c	493 d	779 c	31.4 c
	171.0	192 g	467 de	629 e	23.2 f
	256.5	178 g	412 f	558 f	21.6 g
	342.0	154 h	170 g	425 g	21.4 g
Significant level					
Rice varieties		**	**	**	**
NaCl		**	**	**	**
Rice varieties × NaCl		**	**	**	**

Means with the different letters are significantly different at $p \leq 0.01$ (**) by Duncan's New Multiple Range Test

Photosynthetic pigment concentrations in both salt tolerant and salt sensitive varieties were sharply dropped when exposed to high salinity stress (Table 2). Chl_a, Chl_b, TC and C_{x+c} concentrations in salt-stressed seedlings were significantly decreased, depending on rice varieties, NaCl concentration and their interaction (Table 2). The Chl_a in HJ salt-stressed seedlings cultured on 85.5, 171.0, 256.5 and 342.0 mM NaCl was lower than those cultured on without (Table 2). The Chl_a degradation in salt-stressed seedlings was positively related to water potential in both HJ salt tolerant ($r^2 = 0.78$) (Fig. 2A) and PT1 salt sensitive plants ($r^2 = 0.58$) (Fig. 2B). The Chl_a degradation percentage of salt stressed HJ was gradually increased from 18.75-33.43% whereas that of salt stressed PT1 was sharply exhibited (22.9-58.9%), relating to salt concentration in the culture media. In parallel way, the Chl_b, TC and C_{x+c} in HJ rice seedlings were maintained better than those in PT1 seedlings when exposed to salt stress, especially in 342 mM NaCl. The Chl_a contents in salt stressed seedlings of rice were positively correlated with maximum quantum yield of PSII (F_v/F_m) in both HJ ($r^2 = 0.73$) (Fig. 3A) and PT1 ($r^2 = 0.68$) (Fig. 3B). Chlorophyll a fluorescence parameters, F_v/F_m , Φ_{PSII} , qP and NPQ in salt stressed seedlings were decreased significantly, depending on rice varieties, salt concentration in the media and their interactions (Table 3). In addition to, a positive relationship was found between Φ_{PSII} and TC in HJ ($r^2 = 0.92$) (Fig. 4A) and PT1 ($r^2 = 0.91$) (Fig. 4B). Diminishing of Φ_{PSII} in salt stressed rice seedlings was positive related to plant dry matter of HJ ($r^2 = 0.92$) (Fig. 4A) and PT1 ($r^2 = 0.88$) (Fig. 4B). Growth characteristics, fresh weight, dry weight and dry matter percentage, in rice seedlings was decreased by the factors of rice varieties, salt concentration and their interaction, except in the dry matter parameter was unchanged by rice varieties (Table 4). As well as, the growth reduction percentage of HJ salt tolerance was extensively maintained better than those in PT1 salt susceptible.




Fig. 2. Relationships between water potential (MPa) and chlorophyll *a* concentration ($\mu\text{g g}^{-1}$ FW) of salt-tolerant (A) and salt-sensitive (B) rice seedlings cultured *In vitro* and exposed to salt stress with iso-osmotic level using mannitol application for 4 days.

The photosynthetic pigments in both rice seedlings exposed to iso-osmotic salinity stress decreased significantly (Table 2). There are many reports which show the reduction of photosynthetic pigments in the salt-stressed leaves of different plants such as bean (Parida & Das, 2005; Hernandez *et al.*, 1995), sunflower (Santos, 2004), castor bean (Pinheiro *et al.*, 2008), winter wheat (Sairam & Srivastava, 2002; Zheng *et al.*, 2008), rice (Cha-um *et al.*, 2007a), tomato (Khavarinejad & Mostofi, 1998) and cotton (Meloni *et al.*, 2003). In this study, the photosynthetic pigments including Chl_a, Chl_b, TC and C_{x+c} in salt-stressed leaves of HJ were stabilized better than those in PT1. Under salinity stress, the photosynthetic apparatus, especially chloroplast may be damaged by toxic ion, especially sodium ion (Na^+). Since, the Na^+ can generate the lipid peroxidation in the

Table 3. Maximum quantum yield (F_v/F_m), photosystem II quantum efficiency (Φ_{PSII}), photochemical quenching (qP) and non-photochemical quenching (NPQ) in the leaf tissues of salt-tolerant (HJ) and salt-sensitive (PT1) rice varieties cultured on MS medium and exposed to 0.0, 85.5, 171.0, 256.5 or 342.0 mM NaCl with iso-osmotic level using mannitol application for 4 days

Rice varieties	NaCl (mM)	F_v/F_m	Φ_{PSII}	qP	NPQ
HJ	0.0	0.824 a	0.618 a	0.817 a	0.072 a
	85.5	0.771 b	0.526 b	0.678 b	0.051 b
	171.0	0.729 cd	0.466 d	0.622 d	0.037 d
	256.5	0.620 e	0.414 f	0.581 f	0.026 e
	342.0	0.514 f	0.297 g	0.527 g	0.017 f
PT1	0.0	0.737 c	0.507 c	0.659 c	0.044 c
	85.5	0.728 cd	0.497 c	0.594 e	0.037 d
	171.0	0.709 d	0.446 e	0.528 g	0.011 g
	256.5	0.627 e	0.411 f	0.516 h	0.010 g
	342.0	0.620 e	0.403 f	0.508 h	0.010 g
Significant level					
Rice varieties		**	**	**	**
NaCl		**	**	**	**
Rice varieties \times NaCl		**	**	**	**

Means with the different letters are significantly different at $p \leq 0.01$ (**) by Duncan's New Multiple Range Test.

chloroplast to damage the ultrastructure and chloroplast functions (Hernandez *et al.*, 1995) as well as produce reactive oxygen species (ROs) (Vaidyanathan *et al.*, 2003) to disturb the function of photosynthetic pigments (Demiral & Türkan, 2005). The Na ions generated by salt stress have been reported as the major toxicity to be damaged by the photosynthetic pigments. The function of photosynthetic pigments has been well established as light harvesting and water oxidation in photosystem II (PSII) which is identified by chlorophyll a fluorescence. It is a simple, rapid and sensitive procedure as physiological changes in plant responses to abiotic stresses (Gray *et al.*, 1997; Sudhir *et al.*, 2005; Cha-um *et al.*, 2009). In present study, the chlorophyll a fluorescence including; F_v/F_m , Φ_{PSII} , qP and NPQ in salt-stressed seedlings were significantly reduced (Table 3). The reduction of chlorophyll a fluorescence in higher plant species such as citrus (Lopez-Climent *et al.*, 2008), red raspberry (Neocleous & Vasilakakis, 2007), sunflower (Santos, 2004), rice (Cha-um *et al.*, 2007a) and Rumex (Chen *et al.*, 2004) has been reported. Decrease in chlorophyll a fluorescence is indicated as the loss of photosynthetic pigment function to gain the energy from light in the light reaction of photosynthesis, leading to growth inhibition (Neocleous & Vasilakakis, 2007). There are many documents to report on growth reduction of plant species when exposed to salt stress such as rice (Morsy *et al.*, 2007; Cha-um *et al.*, 2007a; Nguyen *et al.*, 2005; Dionisio-Sese & Tobita, 1998), guava (Ali-Dinar *et al.*, 1999), pea (Hernandez *et al.*, 1995), citrus (Lopez-Climent *et al.*, 2008), safflower (Siddiqi *et al.*, 2007), cotton (Chachar *et al.*, 2008), red raspberry (Neocleous & Vasilakakis, 2007), sunflower (Santos, 2004) and Rumex (Chen *et al.*, 2004).




Fig. 3. Relationships between chlorophyll *a* concentration (µg g⁻¹ FW) and F_v/F_m of salt-tolerant (A) and salt-sensitive (B) seedlings cultured *In vitro* and exposed to salt stress with iso-osmotic level using mannitol application for 4 days.

Conclusions

Na^+ ions in the culture media enriched into root tissues of salt tolerant HJ and salt sensitive PT1 genotypes and subsequently accumulated in the leaf tissues whereas K^+ markedly reduced, especially in HJ rice. The K^+ homeostasis in salt tolerant HJ rice may play a key role as salt defense mechanism. The Na^+ enrichment in the salt-stressed leaf tissues caused to low water potential, pigment degradation and chlorophyll *a* fluorescence diminishing, leading to growth reduction, especially in PT 1 salt sensitive genotype.

Table 4. Fresh weight, dry weight and dry matter of salt-tolerant (HJ) and salt-sensitive (PT1) rice varieties cultured on MS medium and subjected to 0.0, 85.5, 171.0, 256.5 or 342.0 mM NaCl with iso-osmotic level using mannitol application for 4 days.

Rice varieties	NaCl (mM)	Fresh weight (mg)	Dry weight (mg)	Dry matter (%)
HJ	0.0	145.2 a	47.0 a	35.8 a
	85.5	132.1 b	33.7 b	26.5 b
	171.0	116.4 c	25.4 cd	20.3 c
	256.5	112.4 cd	23.9 cde	19.0 cd
	342.0	110.4 cd	20.4 def	16.6 d
PT1	0.0	108.8 d	28.4 c	35.7 a
	85.5	77.1 e	19.6 ef	22.2 c
	171.0	72.7 ef	17.4 f	21.9 c
	256.5	69.3 f	16.3 f	20.6 c
	342.0	68.6 f	16.0 f	19.3 cd

Significant level			
Rice varieties	**	**	ns
NaCl	**	**	**
Rice varieties × NaCl	**	**	*

Means with the different letters are significantly different at $p \leq 0.01$ (**), $p \leq 0.05$ (*) and non-significant (ns) by Duncan's New Multiple Range Test




Fig. 4. Relationships between total chlorophyll concentration ($\mu\text{g g}^{-1}$ FW) and Φ_{PSII} of salt-tolerant (A) and salt-sensitive (B) rice seedlings cultured *In vitro* and exposed to salt stress with iso-osmotic level using mannitol application for 4 days.




Fig. 5. Relationships between Φ_{PSII} and dry matter (%) of salt-tolerant (A) and salt-sensitive (B) rice seedlings cultured *In vitro* and exposed to salt stress with iso-osmotic level using mannitol application for 4 days.

Acknowledgements

This research was supported by The National Center for Genetic Engineering and Biotechnology (BIOTEC; Grant number BT-B-02-RG-BC-4905) and partially supported by the Thailand Graduate Institute of Science and Technology (TGIST; Grant number TGIST 01-48-041), National Science and Technology Development Agency (NSTDA).

References

- Ahmad, M.S.A., F. Javed and M. Ashraf. 2007. Iso-osmotic effect of NaCl and PEG on growth, cations and free proline accumulation in callus tissue of two indica rice (*Oryza sativa* L.) genotypes. *Plant Growth Regul.*, 53: 53-63.

- Ali-Dinar, H.M., G. Ebert and P. Ludders. 1999. Growth, chlorophyll content, photosynthesis and water relations in guava (*Psidium guajava* L.) under salinity and different nitrogen supply. *Gartenbauwissenschaft*, 64: 54-59.
- Allakhverdiev, S.I., A. Sakamoto, Y. Nishiyama, M. Inaba and N. Murata. 2000. Ionic and osmotic stress effect of NaCl-induced inactivation of photosystem I and II in *Synechococcus* sp. *Plant Physiol.*, 123: 1047-1056.
- Ashraf, M. 2004. Some important physiological selection criteria for salt tolerance in plants. *Flora*, 199(5): 361-376.
- Ashraf, M. 2009. Biotechnological approach of improving plant salt tolerance using antioxidants as markers. *Biotechnol. Adv.*, 27: 84-93.
- Aziz, I. and M.A. Khan. 2001. Experimental assessment of salinity tolerance of *Ceriops tagal* seedlings and saplings from the Indus delta, Pakistan. *Aquat. Bot.*, 70: 259-268.
- Chachar, Q.I., A.G. Solangi and A. Verhoef. 2008. Influence of Sodium chloride on seed germination and seedling root growth of cotton (*Gossypium hirsutum* L.). *Pak. J. Bot.*, 40: 183-197.
- Cha-um, S. and C. Kirdmanee. 2009. Effect of salt stress on proline accumulation, photosynthetic ability and growth characters in two maize cultivars. *Pak. J. Bot.*, 41: 87-98.
- Cha-um, S., K. Supaibulwatana and C. Kirdmanee. 2007a. Glycinebetaine accumulation, physiological characterizations and growth efficiency in salt-tolerant and salt-sensitive lines of indica rice (*Oryza sativa* L. ssp. *indica*) in response to salt stress. *J. Agron. Crop Sci.*, 193: 157-166.
- Cha-um, S., P. Vejchasarn and C. Kirdmanee. 2007b. An effective defensive response in Thai aromatic rice varieties (*Oryza sativa* L. spp. *indica*) to salinity. *J. Crop Sci. Biotech.*, 10: 257-264.
- Chen, H.X., W.J. Li, S.Z. An and H.Y. Gao. 2004. Characterization of PSII photochemistry and thermostability in salt treated *Rumex* leaves. *J. Plant Physiol.*, 161: 257-264.
- Chow, W.S., M.C. Ball and J.M. Anderson. 1990. Growth and photosynthetic response of spinach to salinity: implication of K⁺ nutrition for salt tolerance. *Aust. J. Plant Physiol.*, 17: 563-578.
- Çiçek, N. and H. Cakirlar. 2002. The effect of salinity on some physiological parameters in two maize cultivars. *Bulg. J. Plant Physiol.*, 28: 66-74.
- da Silva, E.C., R.J.M.C. Nogueira, F.P. de Araujo, N.F. de Melo and A.D. de Azevedo Neto. 2008. Physiological responses to salt stress in young umbu plants. *Environ. Exp. Bot.*, 63: 147-157.
- Demiral, T. and I. Türkan. 2005. Comparative lipid peroxidation, antioxidant defense systems and proline content in roots of two rice cultivars differing in salt tolerance. *Environ. Exp. Bot.*, 53: 247-257.
- Dionisio-Sese, M.L. and S. Tobita. 1998. Antioxidant responses of rice seedlings to salinity stress. *Plant Sci.*, 135: 1-9.
- Flower, T.J. and A.R. Yeo. 1995. Breeding for salinity resistance in crop plants: where next? *Aust. J. Plant Physiol.*, 22: 875-884.
- Gray, R., L.P. Chauvin, F. Sarhan and N.P.A. Huner. 1997. Cold acclimation and freezing tolerance (A complex interaction of light and temperature). *Plant Physiol.*, 114: 467-474.
- Hasegawa, P.M., R.A. Bressan, J. Zhu and H.J. Bohnert. 2000. Plant cellular and molecular responses to high salinity. *Annu. Rev. Plant Physiol. Plant Mol. Biol.*, 51: 463-499.
- Hernandez, J.A., E. Olmos, F.J. Corpas, F. Sevilla and L.A. Del Rio. 1995. Salt-induced oxidative stress in chloroplasts of pea plants. *Plant Sci.*, 105: 151-167.
- Hu, Y., Z. Burucs, S. von Tucher and U. Schmidhalter. 2007. Short-term effects of drought and salinity on mineral nutrient distribution along growing leaves of maize seedlings. *Environ. Exp. Bot.*, 60: 268-275.
- Khavarinejad, R.A. and Y. Mostofi. 1998. Effects of NaCl on photosynthetic pigments, saccharides, and chloroplast ultrastructure in leaves of tomato cultivars. *Photosynthetic.*, 35: 151-154.
- Khelil, A., T. Menu and B. Ricard. 2007. Adaptive response to salt involving carbohydrate metabolism in leaves of a salt-sensitive tomato cultivar. *Plant Physiol. Biochem.*, 45: 551-559.
- Kush, G.S. 1997. Origin, dispersal, cultivation and variation of rice. *Plant Mol. Biol.*, 35: 25-34.

- Levigneron, A., F. Lopez, G. Vasuyt, P. Berthomieu, P. Fourcroy and F. Casse Delbart. 1995. Plants toward salt stress. *Cahiers d'Agricultures*, 4: 263-273.
- Lichtenthaler, H.K. 1987. Chlorophylls and carotenoids: pigments of photosynthetic biomembranes. *Method Enzymol.*, 148: 350-380.
- Loggini, B., A. Scartazza, E. Brugnoli and F. Navari-Izzo. 1999. Antioxidant defense system, pigment composition and photosynthetic efficiency in two wheat cultivars subjected to drought. *Plant Physiol.*, 119: 1091-1099.
- Lopez-Climent, M.F., V. Arbona, R.M. Perez-Clemente and A. Gomez-Cadenas. 2008. Relationship between salt tolerance and photosynthetic machinery performance in citrus. *Environ. Exp. Bot.*, 62: 176-184.
- Luo, Q., B. Yu and Y. Liu. 2005. Differential sensitivity to chloride and sodium ions in seedlings of *Glycine max* and *G. soja* under NaCl stress. *J. Plant Physiol.*, 162: 1003-1012.
- Maxwell, K. and G.N. Johnson. 2000. Chlorophyll fluorescence -a practical guide. *J. Exp. Bot.*, 51: 659-668.
- Meloni, D.A., M.A. Oliva, C.A. Martinez and J. Cambraia. 2003. Photosynthesis and activity of superoxide dismutase, peroxidase and glutathione reductase in cotton under salt stress. *Environ. Exp. Bot.*, 49: 69-76.
- Morsy, M.R., L. Jouve, J.F. Hausman, L. Hoffmann and J. McD. Stewart. 2007. Alteration of oxidative and carbohydrate metabolism under abiotic stress in two rice (*Oryza sativa* L.) genotypes contrasting in chilling tolerance. *J. Plant Physiol.*, 164: 157-167.
- Murashige, T. and F. Skoog. 1962. A revised medium for rapid growth and bioassays with tobacco tissue cultures. *Physiol. Plant.*, 15: 473-479.
- Nakamura, I., S. Murayama, S. Tobita, B. Ba Bong, S. Yanagihara, Y. Ishimine and Y. Kawamitsu. 2002. Effect of NaCl on the photosynthesis, water relations and free proline accumulation in the wild *Oryza* species. *Plant Prod. Sci.*, 5: 305-310.
- Navarro, A., S. Banon, E. Olmos and M.J. Sanchez-Blanco. 2007. Effect of sodium chloride on water potential components, hydraulic conductivity, gas exchange and leaf ultrastructure of *Arbutus unedo* plants. *Plant Sci.*, 172: 473-480.
- Neocleous, D. and M. Vasilakakis. 2007. Effects of NaCl stress on red raspberry (*Rubus idaeus* L. 'Autumn Bliss'). *Sci. Hort.*, 112: 82-289.
- Nguyen, H.T.T., I.S. Shim, K. Kobayashi and K. Usui. 2005. Effects of salt stress on ion accumulation and antioxidative enzyme activities of *Oryza sativa* L., and *Echinochloa oryzicola* Vasing. *Weed Biol. Manage.*, 5: 1-7.
- Noreen, S. and M. Ashraf. 2008. Alleviation of adverse effects of salt stress on sunflower (*Helianthus annuus* L.) by exogenous application of salicylic acid: Growth and photosynthesis. *Pak. J. Bot.*, 40(4): 1657-1663.
- Parida, A.K. and A.B. Das. 2005. Salt tolerance and salinity effects on plant: a review. *Ecotoxic. Environ. Safe*, 60: 324-349.
- Parida, A.K., A.B. Das, Y. Sanada and P. Mohanty. 2004. Effects of salinity on biochemical components of the mangrove, *Aegiceras corniculatum*. *Aquat. Bot.*, 80: 77-87.
- Pinheiro, H.A., J.V. Silva, L. Endres, V.M. Ferreira, C.A. Camara, F.F. Cabral, J.F. Oliveira, L.W.T. de Carvalho, J.M. dos Santos and B.G. dos Santos Filho. 2008. Leaf gas exchange, chloroplast pigments and dry matter accumulation in castor bean (*Rinicus communis* L) seedlings subjected to salt stress conditions. *Ind. Crop Prod.*, 27: 385-392.
- Rejili, M., A.M. Vadel, A. Guetet and M. Neffatti. 2007. Effect of NaCl on the growth and the ionic balance K^+/Na^+ of two populations of *Lotus creticus* (L.) (Papilionaceae). *South Afri. J. Bot.*, 73: 623-631.
- Sabir, P. and M. Ashraf. 2007. Screening of local accessions of *Panicum maliaceum* L., for salt tolerance at seedling stage using biomass production and ion accumulation as selection criteria. *Pak. J. Bot.*, 39(5): 1655-1661.
- Sairam, R.K. and G.C. Srivastava. 2002. Changes in antioxidant activity in sub-cellular fractions of tolerant and susceptible wheat genotypes in response to long term salt stress. *Plant Sci.*, 162: 897-904.

- Santos, C.V. 2004. Regulation of chlorophyll biosynthesis and degradation by salt stress in sunflower leaves. *Sci. Hort.*, 103: 93-99.
- Saqib, M., C. Zorb, Z. Rengel and S. Schubert. 2005. The expression of the endogenous vacuolar Na⁺/K⁺ antiporters in roots and shoots correlates positively with the salt resistance of wheat (*Triticum aestivum* L.). *Plant Sci.*, 169: 959-965.
- Serrano, R. and A.R. Navarro. 2001. Ion homeostasis during salt stress in plants. *Curr. Opin. Cell Biol.*, 13: 399-404.
- Shabala, S.N., S.I. Shabala, A.I. Martynenko, O. Babourina and I.A. Newman. 1998. Salinity effect on bioelectric activity, growth, Na⁺ accumulation and chlorophyll fluorescence of maize leaves: a comparative survey and prospects for screening. *Aust. J. Plant Physiol.*, 25: 609-616.
- Shannon, M.C., J.D. Rhoades, J.H. Draper, S.C. Scardaci and M.D. Spyres. 1998. Assessment of salt tolerance in rice cultivars in response to salinity problems in California. *Crop Sci.*, 38: 394-398.
- Siddiqi, E.H., M. Ashraf and N.A. Akram. 2007. Variation in seed germination and seedling growth in some diverse lines of safflower (*Carthamus tinctorius* L.) under salt stress. *Pak. J. Bot.*, 39: 1937-1944.
- Sudhir, P.R., D. Pogoryelov, L. Kovacs, G. Garab and S.D.S. Murthy. 2005. The effects of salt stress on photosynthetic electron transport and thylakoid membrane proteins in the *Cyanobacterium, Spirulina platensis*. *J. Biochem. Mol. Biol.*, 38: 481-485.
- Taiz, L. and E. Zieger. 2002. *Plant Physiology* 3rd edition. Sinauer Associate, Inc. Publisher, Sunderland Massachusetts. 690 p.
- Tonon, G., C. Kevers, O. Faivre-Rampant, M. Graziani and T. Gaspar. 2004. Effect of NaCl and mannitol iso-osmotic stresses on proline and free polyamine levels in embryogenic *Fraxinus angustifolia* callus. *J. Plant Physiol.*, 161: 701-708.
- Vaidyanathan, H., P. Sivakumar, R. Chakrabarty and G. Thomas. 2003. Scavenging of reactive oxygen species in NaCl-stressed rice (*Oryza sativa* L.) differential response in salt-tolerant and sensitive varieties. *Plant Sci.*, 165: 1411-1418.
- Yasar, F., O. Uzal, S. Tufenkci and K. Yildiz. 2006. Ion accumulation in different organs of green bean genotypes grown under salt stress. *Plant Soil Environ.*, 52: 476-480.
- Yokoi, S., R.A. Bressan and P.M. Hasegawa. 2002. Salt stress tolerance of plants. JIRCAS Working Report, pp. 25-33.
- Zeng, L. 2005. Exploration of relationships between physiological parameters and growth performance of rice (*Oryza sativa* L.) seedlings under salinity stress using multivariate analysis. *Plant Soil*, 268: 51-59
- Zheng, Y., Z. Wang, X. Sun, A. Jia, G. Jiang and Z. Li. 2008. Higher salinity tolerance cultivars of winter wheat relieved senescence at reproductive stage. *Environ. Exp. Bot.*, 62: 129-138.

(Received for publication 15 April 2009)