

ISOLATION OF FUNGI ASSOCIATED WITH SHISHAM TREES AND THEIR EFFECT ON SEED GERMINATION AND SEEDLING MORTALITY

NASIR AHMED RAJPUT*¹, M.A. PATHAN^{1,2}, A. QAYOOM RAJPUT³,
M.M. JISKANI¹, A. MUBEEN LODHI¹, SABIT AHSAN RAJPUT⁴
AND M.I. KHASKHALI⁵

¹Department of Plant Pathology, Sindh Agriculture University, Tandojam, Pakistan.

²Department of Plant Pathology, Lasbela University of Agriculture,
Water and Marine Sciences

³EFUPVS & NPP, Ministry of Food & Agriculture, Pakistan

⁴FSC & RD, Ministry of Food & Agriculture, Pakistan.

Abstract

A total of 10 fungi viz., *Fusarium solani*, *F. moniliforme*, *F. equiseti*, *F. oxysporum*, *F. semitectum*, *Rhizoctonia solani*, *Alternaria alternata*, *Curvularia lunata*, *Aspergillus niger* and *Penicillium* sp. were isolated from infected roots, bark, seed and stem of shisham. *F. solani* was most predominant followed by *R. solani* and *C. lunata*. Maximum infection frequency of 75.00% was exhibited by *F. solani* colonizing stem tissues followed by 47.39% from bark tissues, 29.83% from seeds and 15.62% from roots. The colonization percent of *F. solani* was highest in stem tissues collected from Ghotki (85.50%) as compared to Pano Akil (70.00%), Mirpur Mathelo (65.50%), Daharki (56.50%) and Hala (34.60%) followed by *F. moniliforme* ranging from 3.00-18.50% as compared to other isolated fungi. Seed germination percentage was also reduced (50.00%) in soil infested with *F. solani* and seedling mortality was 93.33% followed by soil infested with *R. solani* (60.00%) with mortality rate (66.66%) and *C. lunata* (70.00%), seedling mortality rate of 42.85% as compared to *F. moniliforme* and *F. oxysporum*, respectively.

Introduction

Shisham (*Dalbergia sissoo* Roxb.), native to Haryana and other parts of India, Pakistan and Nepal is of great importance because of its multiple uses, such as furniture wood, building timber, plywood and fuel. It is medium to large sized deciduous tree with a light crown and is perpetuated by seeds and suckers (Hassan, 2005). It is cultivated in forest plantation, along water channel and canal banks, road sides and railway lines. Unfortunately shisham is susceptible to dieback, wilt and several other soilborne pathogens (Sah *et al.*, 2003). Previously Bakshi (1974) isolated *Phellinus gilvus* from roots of trees affected by dieback. Richardson (1990) reported several species of *Aspergillus*, *Penicillium*, *Rhizopus*, *Alternaria*, *Fusarium*, *Chaetomium*, *Drechslera* and *Curvularia* from forest tree seeds. Parajuli *et al.*, (1999) reported *Fusarium oxysporum* from *Dalbergia sissoo* on water-logged soils in Nepal. Manadhar & Shrestha (2000) isolated *Botryodiplodia* sp., and *Fusarium solani* from five diseased samples of *D. sissoo*. Khan *et al.*, (2001b) detected *Aspergillus niger*, *A. flavus*, *A. terreus*, *Aspergillus* sp., *Alternaria alternata*, *Chaetomium* sp., *Drechslera australiensis*, *Fusarium pallidoroseum*, *F. solani*, *Fusarium* sp., *Penicillium* sp., *Rhizopus* sp., and *Geotrichum* sp. from seeds of shisham trees. Rajput *et al.*, (2008) isolated *F. solani*, *Rhizoctonia solani* and *Curvularia lunata* as predominant fungi from shisham die-back trees.

*Corresponding author: E-mail: nasir_ahmedrao@hotmail.com

The insufficient production of shisham is due to dieback and decline and high mortality of seeds and seedlings (Khan & Khan, 2000). The pathogens penetrate into the seed coat and embryo during storage and are responsible for poor germination as well as seedling mortality (Bhansli & Jindal, 1997). Mortality as high as 20-28% has been obtained in Nepal (Joshi & Baral, 2000). Bakhisa & Bask (2000) reported that shisham decline resulted in sudden death of 10% plantation within 6 months after crowns exhibited symptoms of the disease. Bajwa *et al.*, (2003) found 20-28% shisham trees affected by decline during a survey of different areas of Punjab. The present studies were taken up for the first time in Sindh to isolate and identify fungi associated with shisham dieback and their impact on seed germination and seedling mortality.

Materials and Methods

Isolation from root and stem: A survey of shisham growing areas of Sindh *viz.*, Ghotki, Pano Akil, Mirpur Mathelo, Dahrki, Hyderabad, Hala, Tandojam and Dadu was carried out. From these localities the samples were collected from shisham trees showing infected branches. Diseased samples including root bark from collar portion, and stem, collected in sterilized polythene bags and brought to the laboratory for the isolations of the associated pathogens. At least 50 samples were taken from five different sites of each locality.

The infected roots, bark, and stem were used for isolation as described by Pathak (1987). Isolations were made on three layers of moistened blotter papers and potato dextrose agar medium (Saleem & Nasir, 1991). Samples were surface sterilized with 0.01% HgCl₂ for two minutes, rinsed with sterile distilled water for 2-3 minutes and then placed on blotter papers and PDA under aseptic conditions. Plates were incubated at 25°C for 7 days.

The fungi growing from infected tissues were identified on the basis of colony characteristics and conidial morphology using keys of Barnett & Hunter (1972), Booth (1977), Neergaard (1979) and Hawksworth *et al.*, (1995).

Isolation from seed: Samples of shisham seed were collected from various areas of Sindh including Ghotki, Pano Akil, Mirpur Mathelo, Daharki, Dadu, Hyderabad, Tandojam and Hala. Seeds were assayed for the presence of seed-borne fungi by standard blotter method (Anon., 1996) on PDA, 100 seeds from each seed lot were placed on moistened blotter papers and freshly prepared PDA medium with five seeds per plate. Plates were incubated at 25°C for 7 days and observed for fungal growth. The frequency of fungi was estimated using the following formula:

$$\text{Colonization (\%)} = \frac{\text{Number of root pieces/ seeds colonized by a pathogen}}{\text{Total number of pieces/seeds studied}} \times 100$$

Seed germination and seedling mortality: Thirty shisham seeds were grown in earthen pots containing 2kg steam sterilized soil previously infested with either *Fusarium solani*, *F. moniliforme*, *F. oxysporum*, *Rhizoctonia solani* or *Curvularia lunata* separately. Seeds were allowed to germinate under natural day light conditions and watered when ever needed.

Germination percent was recorded as under:

$$\text{Germination (\%)} = \frac{\text{Number of seeds germinated}}{\text{Total number of seeds used}} \times 100$$

The seedling mortality was also estimated using the following formula:

$$\text{Mortality (\%)} = \frac{\text{Number of dead seedlings}}{\text{Total number of seeds germinated}} \times 100$$

Results and Discussion

Isolation of fungi: *Fusarium solani* (Mart.) Sacc. was isolated most abundantly from all plant parts. Mycelial growth was sparse to dense and grayish to white. Micro-conidia formed abundantly on the aerial mycelium, may be single septate, 9-16 x 2-4 u. Macro-conidia 4-5 septate were well developed and were 40-100 x 5-7.5 u.

Fusarium semitectum var. *majus* Wollenw; showed cottony white with abundant conidiophores in aerial mycelium.

Fusarium oxysporum Schlecht; showed white but usually with light purple colour mycelium. Micro and macro-conidia formed frequently.

Fusarium equiseti (Corda) Sacc., showed first white but later deep olive in colour mycelium.

Fusarium moniliforme Sheld., produces extensive cottony mycelial growth. Conidia are multi-septate and larger formed on typical conidiophores.

Alternaria alternata Auct; develops typical mycelial growth with small conidia but in long chains.

Curvularia lunata (Walker) Boedjin; produces septate mycelium, at first hyaline which later becomes brown.

Rhizoctonia solani Kuhn; produces a dense pale to dark brown mycelium which becomes darker with age.

F. solani was isolated predominantly from stem tissues collected from all locations (34.6-85.50%) as compared to root pieces followed by *F. moniliforme* (3.00-18.50%), *Rhizoctonia solani* (7.50-15.50%) and *Curvularia lunata* (5.50-14.50%), respectively, than all other fungi which were isolated with low frequency (Table 1).

Maximum infection frequency (75.00%) was exhibited by *F. solani* stem tissues followed by 47.39% bark tissues, 29.83% seeds and 15.62% root tissues (Table 2) as compared to *F. moniliforme* 20.00% from bark tissues, 15.00% stem tissues, 11.66% seeds and 9.37% root tissues, *Rhizoctonia solani* (25.00%) from bark, 19.37% stem tissues, 15.00% root tissues and *Curvularia lunata* 25.50% from seeds, 23.75% bark tissues, 17.50% stem tissues and 14.75% root tissues (Table 2). The other fungi were isolated with very low frequency from all parts of shisham tree (Table 2). Ahmad & Bhutta (1993) isolated *F. solani*, and *F. pallidoroseum* from seeds of *Dalbergia sissoo* and *Leucaena leucocephala*. Shakir *et al.*, (1999) reported the association of *F. solani* with roots of affected shisham trees alongwith species of *Aspergillus*, *Cladosporium*, *Fusarium*, *Verticillium* and *Tylenchorhynchus*, *Helicotylenchus*, *Hoplolaimus*, *Dorylaimus* and *Xiphinema* nematodes. Manadhar & Shrestha (2000) found *Botryodiplodia* sp., and *Fusarium solani* associated with 5 diseased samples of *D. sissoo*. The species of *Alternaria*, *Aspergillus* and *Fusarium* were also detected from seeds of *D. sissoo* (Manadhar *et al.*, 2000). Results of our studies also agreed with those reported by Khalid *et al.*, (2002). Mustafa *et al.*, (2004) isolated *Rhizoctonia solani*, *Fusarium solani*, *F. oxysporum*, *F. moniliforme*, *Aspergillus niger*, *Alternaria alternata* and *Helminthosporium* sp., as seed-borne fungi from seed samples of shisham.

Table 1. Fungi isolated from root and stem of shisham dieback samples collected from different localities of Sindh.

Fungi isolated	Colonization (%) at different localities																	
	Ghotki		Pano Akil		Mirpur Mathelo		Daharki		Dadu		Hyderabad		Tandojam		Hala			
	R*	S**	R	S	R	S	R	S	R	S	R	S	R	S	R	S		
<i>Fusarium solani</i>	12.5	85.5	15.5	70.0	20.2	65.5	15.6	56.5	15.5	55.5	25.5	35.5	23.5	40.2	10.5	34.6		
<i>Fusarium moniliforme</i>	15.5	18.5	12.5	15.5	10.5	12.5	15.4	16.5	12.5	20.5	9.5	7.5	3.5	5.5	6.5	3.0		
<i>Fusarium equiseti</i>	11.5	12.5	15.5	10.4	9.5	11.5	20.5	15.4	0	0	0	0	6.5	8.4	0	0		
<i>Fusarium oxysporum</i>	9.5	2.5	5.4	3.5	0	0	2.5	1.5	0	0	0	0	0	0	0	0		
<i>Fusarium semitectum</i>	7.5	1.5	5.2	1.5	0	0	1.5	1.0	0	0	0	0	0	0	0	0		
<i>Rhizoctonia solani</i>	18.5	15.5	20.5	21.5	20.5	12.5	22.5	16.5	17.5	16.4	9.5	7.5	18.4	15.2	15.4	7.5		
<i>Alternaria alternata</i>	10.5	12.5	8.5	13.4	6.4	12.5	11.4	13.5	5.4	3.5	7.5	10.4	2.5	12.2	2.5	8.4		
<i>Curvularia lunata</i>	15.6	14.5	18.4	20.3	15.4	10.4	12.4	9.5	10.2	8.5	6.5	3.5	1.5	7.5	0	5.5		
<i>Aspergillus niger</i>	7.5	3.4	6.5	2.5	5.4	1.5	6.4	0	5.4	0	7.2	1.5	5.4	1.2	0	1.3		
<i>Penicillium</i> sp.	5.5	2.2	5.3	2.1	0	0	4.3	2.2	3.5	0	2.4	1.0	3.2	0	2.5	0		

R* = Root, S** = Stem

Table 2. Overall infection of fungi isolated from different parts of shisham.

Fungi isolated	Bark infection (%)		Stem infection (%)		Root infection (%)		Seed infection (%)	
<i>Fusarium solani</i>	47.39	75.00	15.62	29.83	15.62	29.83	15.62	29.83
<i>Fusarium moniliforme</i>	20.00	15.00	9.37	11.66	9.37	11.66	9.37	11.66
<i>Fusarium equiseti</i>	15.00	10.00	14.37	0	14.37	0	14.37	0
<i>Fusarium oxysporum</i>	0	14.37	4.37	0	4.37	0	4.37	0
<i>Fusarium semitectum</i>	0	0	0	17.91	0	17.91	0	17.91
<i>Rhizoctonia solani</i>	25.00	19.37	15.00	0	15.00	0	15.00	0
<i>Alternaria alternata</i>	12.00	8.75	8.75	10.41	8.75	10.41	8.75	10.41
<i>Curvularia lunata</i>	23.75	17.50	14.75	25.50	14.75	25.50	14.75	25.50
<i>Aspergillus niger</i>	08.00	2.50	5.62	8.75	5.62	8.75	5.62	8.75
<i>Penicillium</i> sp.	0	1.25	3.75	6.66	3.75	6.66	3.75	6.66

Table 3. Effect of predominant fungi on seed germination and seedling mortality.

Fungi	Inoculated germination (%)	No. of dead seedlings	Mortality (%)
<i>Fusarium solani</i>	50	14	93.33
<i>Rhizoctonia solani</i>	60	12	66.66
<i>Curvularia lunata</i>	70	9	42.85
<i>Fusarium moniliforme</i>	80	6	25.00
<i>Fusarium oxysporum</i>	90	3	11.11

Seed germination percentage was also affected in soil infested with *Fusarium solani* (50.00%) followed by *Rhizoctonia solani* (60.00%), *Curvularia lunata* (70.00%), *F. moniliforme* (80.00%) and *F. oxysporum* (90.00%) respectively, as compared to uninfested soil (Table 3). Seedling mortality rate was maximum (93.33%) for *F. solani* followed by *R. solani* (66.66%), *Curvularia lunata* (42.85%), and *F. moniliforme* (25.00%). The least mortality rate of 11.11% was obtained in pots infested with *F. oxysporum* (Table 3). Vigayan & Rehill (1990) and Pathan *et al.*, (2007) found reduction in germination of shisham seeds infected with *Aspergillus flavus*, *A. niger*, *F. oxysporum* and *F. solani*. Saleem (1999) observed that disease progressed in mango downward and bark was discolored some distance from the tip. Khan *et al.*, (2004) observed mortality ranging from 25.00-30.00% and a disease incidence of 20.50-40.00% in shisham trees in various districts of Punjab including Kasur, Toba Tek Singh, Hafizabad and Gujranwala. Our results also agreed with Shailendra *et al.*, (2004), who recorded highest mortality in shisham plantation in India due to *Fusarium solani*.

References

- Ahmad, I. and A.R. Bhutta. 1993. Fungi associated with land scape tree seed in Islamabad, Pakistan. *Pak. J. Phytopathol.*, 5: 126-129.
- Anonymous. 1996. *International Rules for seed testing*. Proc. of Intl. Seed Testing Association, Zurich.
- Bajwa, R., J. Arshad and A. Saleh. 2003. Extent of shisham (*Dalbergia sissoo* Roxb.) decline in Sialkot, Gujranwala, Lahore and Sargodha districts. *Mycopath.*, 1: 1-5.
- Baksha, M.W. and A.C. Bask. 2000. Mortality of sissoo in Bangladesh. *Proc. of 3rd Natl. Conf. of Pl. Pathol. 2001, NARC, Islamabad*. 33-37.
- Bakshi, B.K. 1974. Control of root disease in plantation in reforested stands. *Indian Forester*, 100: 77-78.
- Barnett, H.L. and B.B. Hunter. 1972. *Illustrated Genera of Imperfect Fungi*. Sant Paul Minnesota, USA. 241 pp.
- Bhansli, R.R. and S.K. Jindal. 1997. *Diseases of trees and their management. Agroforestry sustained productivity in Arid Region*. Scientific Publisher, India.
- Booth, C. 1977. *The genus Fusarium*. CMI, Kew, Surrey, England.
- Hassan, B. 2005. Disease destroying shisham trees. Dawn May 9th 2005. Lahore, Pakistan.
- Hawksworth, D.L., P.M. Kirk, B.C. Sutton and D.N. Pegler. 1995. *Ainsworth and Bisby's Dictionary of the Fungi. 8th Ed.* CAB Intl. Wallingford, Oxan Ox 108 DE, UK.
- Joshi, R.B. and S.R. Baral. 2000. A report on dieback of *Dalbergia sissoo* Roxb., in Nepal. In: *Proc. of the sub-regional seminar on dieback of sissoo (Dalbergia sissoo Roxb.), Katmandu, Nepal, 25-28 April 2000*, pp. 17-22.
- Khalid, N., A.S. Anwar, M.I. Haque and A. Riaz. 2002. Study on occurrence of seed-borne fungi and their impact on seed germination of five forest trees. *Pak. J. Phytopathol.*, 14: 47-50.

- Khan, M.M. and M.H. Khan. 2000. Dieback of *Dalbergia sissoo* Roxb., in Pakistan. In: *Proc. of the sub-regional seminar on dieback of sissoo (Dalbergia sissoo Roxb.), Katmandu, Nepal, 25-28 April 2000*, pp. 51-56.
- Khan, S.H., M. Idress, F. Muhammad, A. Mahmood and S.H. Zaidi. 2004. Incidence of shisham (*Dalbergia sissoo* Roxb.) decline and *in vitro* response of isolated fungus species to various fungicides. *Int. J. of Agril. Biol.*, 6: 611-614.
- Khan, S.M., A. Rehman and M.K. Samiya. 2001a. Problem and progress of shisham decline in the Punjab. *Proc. Of 3rd Natl. Conf. of Pl. Pathol., Oct. 1-3, 2001, NARC, Islamabad*, pp. 49-51.
- Khan, S.M., A.S. Shakir, M.A. Tabssum and A. Rehman. 2001b. Isolation and identification of different fungi from diseased shisham tree. *Proc. of 3rd Natl. Conf. of Pl. Pathol. Oct. 1-3, 2001. NARC, Islamabad*, pp. 44-46.
- Manadhar, G. and S.K. Shrestha. 2000. Fungi associated with dieback of *D. sissoo*. In: *Proc. the sub-regional seminar on dieback of sissoo (Dalbergia sissoo Roxb.), Katmandu, Nepal, 25-28 April 2000*, pp. 27-30.
- Manadhar, G., S.K. Shrestha, S. Appanah, G. Allard and S.M. Amatya. 2000. Fungi associated with dieback of sissoo. *Proc., of Intl. Seminar, Nepal*, (18): 27-29.
- Mustafa, A., S.M. Khan and A. Rehman. 2004. Fungi associated with shisham (*Dalbergia sissoo* Roxb.) seed and their control. *Pak. J. Phytopathol.*, 16: 73-75.
- Neergaard, P. 1979. *Seed Pathology*. Vol. 1. The McMillan Press Ltd. London. 240 pp.
- Parajuli, A.V., B. Bhatta, M.K. Adhikary, J. Tuladhar and B. Thapa. 1999. Causal agents responsible for the dieback of *Dalbergia sissoo* Roxb., in the eastern Nepal Terai. *Ban Ko Jankari*, 9: 7-14.
- Pathak, V.N. 1987. *Laboratory Manual of Plant Pathology*. 2nd ed. pp. 23-50. Oxford IBH Pub. Co. New Delhi.
- Pathan, M.A., N.A. Rajput, M.M. Jiskani and K.H. Wagan. 2007. Studies on intensity of shisham dieback in Sindh and impact of seed-borne fungi on seed germination. *Pak. J. Agric., Agril. Engg., Vet. Sci.*, 23: 12-17.
- Rajput, N.A., M.A. Pathan, M.M. Jiskani, A.Q. Rajput and R.R. Arain. 2008. Pathogenicity and host range of *Fusarium solani* (Mart.) Sacc., causing dieback of Sisham (*Dalbergia sissoo* Roxb.). *Pak. J. Bot.*, 40(6): 2631-2639.
- Richardson, M.J. 1990. *An Annotated List of Seed-borne Diseases*. 4th ed. ISTA, Zurich.
- Sah, S.P., C.K. Sharma and F. Sehested. 2003. Possible role of the soil in the sissoo forest (*Dalbergia sissoo* Roxb.) decline in the Nepal Terai. *Plant Soil Environ.*, 49: 378-385.
- Saleem, A. 1999. Recent development in the management of mango disease. In: *Proc. of 2nd Natl. Conf of Plant Pathol. Sept. 27-29, 1999. Univ. of Agri. Faisalabad*, pp. 6-11.
- Saleem, A. and M.A. Nasir. 1991. *Culture Media*. Directorate of Agric. Information, Agriculture Department, Government of the Punjab, Lahore.
- Shailendra, K., K. Surinder, K.S. Kapoor, S. Ranjeet and S. Chakrabarti. 2004. Mortality of *Dalbergia sissoo* Roxb. (Shisham) in Subathu Forest Range of Solan, Himachal Pardesh: a case study. *Indian Forester*, 130: 349-350.
- Shakir, A.S., S.M. Khan and R. Ahmad. 1999. First report on shisham (*Dalbergia sissoo* Roxb.) decline in Pakistan. *Pak. J. Phytopathol.*, 11: 106.
- Vigayan, A.K. and P.S. Rehill. 1990. Effect of culture filtrates of some seed-borne fungi of *Dalbergia sissoo* Roxb., on seed germination and seedling growth. *Indian Forester*, 116: 559-563.

(Received for publication 21 March 2009)