

EFFECT OF NITROGEN APPLICATION ON YIELD AND YIELD COMPONENTS OF BARLEY (*HORDEUM VULGARE* L.)

MOHAMMAD SHAFI^{1*}, JEHAN BAKHT², FAZAL JALAL¹, MOHAMMAD AMAN KHAN³, AND SABIR GUL KHATTAK⁴

¹Department of Agronomy, KPK. Agricultural University Peshawar Pakistan.

²Institute of Biotechnology and Genetic Engineering, KPK. Agricultural University Peshawar, Pakistan

³Department of Biotechnology and Bioinformatics QAU Islamabad Pakistan

⁴Agricultural Research Institute Tarnab Peshawar, KPK

*Corresponding author E-mail: mshafi.aup@gmail.com

Abstract

The present research was aimed to study the effect of different levels of inorganic fertilizer N on the yield and yield components of barley varieties at KPK Agricultural University, Peshawar, Pakistan. The experiment was laid out in randomized complete block design with split plot arrangements. Barley varieties (local and sterling) were sown in the main plots whereas different nitrogen levels (i.e. 0, 20, 40, 60, 80 and 100 kg ha⁻¹) were allotted to sub-plots. The results showed that different varieties and nitrogen levels had significantly ($p < 0.05$) affected plant height, 1000 grains weight, grains spike⁻¹, straw yield, grain yield, biological yield, grain and plant N. Maximum grain spike⁻¹, thousand grain weight and grain yield was recorded in sterling. Taller plants, highest biological yield, straw yield, grain and plant N was observed in local variety. Nitrogen applied at the rate of 60 kg ha⁻¹ resulted in maximum grain spike⁻¹, thousand grain weight, biological yield, grain yield, grain and plant N.

Introduction

Barley (*Hordeum vulgare* L.) is the major cereal in many dry areas of the world and is vital for the livelihoods of many farmers. Barley is an annual cereal crop and grown in environments ranging from the desert of the Middle East to the high elevation of the Himalayas (Hayes *et al.*, 2003). It is the major food source in many North African countries. In Pakistan, it is mainly grown for grain and straw for small ruminants during winter, with green fodder sometimes used for winter grazing. Barley can replace wheat as the dominant crop due to its tolerance to drought and salinity. Barley assumes fourth position in total cereal production in the world after wheat, rice, and maize. Barley is more productive under adverse environments than other cereals. Barley serves as a major animal fodder, base malt for beer and certain other distilled beverages.

Nitrogen is commonly the most limiting nutrient for crop production in the major world's agricultural areas and therefore adoption of good N management strategies often result in large economic benefits to farmers. Among the plant nutrients, nitrogen plays a very important role in crop productivity (Zapata & Cleenput, 1986; Ahmad, 1999; Miao *et al.*, 2006; Oikeh *et al.*, 2007; Worku *et al.*, 2007). Efficient use of N is also important for minimizing environmental contamination (Scharf & Alley, 1988). Nitrogen is the key element in achieving consistently high yields in cereals. Nitrogen is a constituent of many fundamental cell components such as nucleic acids, amino acids, enzymes, and photosynthetic pigments. The rate of uptake and partition of N is largely determined by supply and demand during various stages of plant growth. Soil N supply, for example must be high at tillering, stem elongation, booting, heading and grain filling requiring a greater amount of the development and growth of its reproductive organs and for an enhanced and high accumulation of proteins in the kernel. Nitrogen is considered one of the most important factors affecting crop morphology (Amanullah *et al.*, 2008a), crop growth rate and grain yield (Amanullah *et al.*, 2008b) in Northwest Pakistan. The amount of nitrogen that a barley crop needs to maximize yield and quality will depend on the seasonal conditions, soil type, and rotational history of the soil as well as the potential yield of the crop. Nitrogen is needed for early tiller development of barley to set up the crop for a high yield potential. Ayoub *et al.*, (1994) reported that split N application had little effect on yield, but decreased lodging and spike population with increased grain weight. Singh & Uttam (1992)

recorded increased grain yield with increase in nitrogen level. However, increasing N fertility beyond a certain limit induced lodging and ultimately decreased grain yield and its components. Nowadays, both to avoid pollution by nitrates and to maintain a sufficient profit margin, farmers have to reduce the use of nitrogen fertilizer. These objectives can be met through efficient farming techniques, but also by using plant varieties that have better nitrogen use efficiency (Gallais & Hirel, 2004). For NUE, genetic variability and genotype x nitrogen fertilization level interactions reflecting differences in responsiveness have been observed in maize (Bertin & Gallais, 2000). The present study was carried out to investigate the response of barley varieties to different nitrogen levels.

Materials and Methods

The present study was conducted at KPK Agricultural University, Peshawar, Pakistan using randomized complete block design with split plot arrangements replicated four times. Six levels of nitrogen (0, 20, 40, 60, 80 and 100 kg N ha⁻¹) and two barley varieties (sterling and local variety) were studied during the present experiment. A plot size of 2.5 m x 2 m having 6 rows, 30 cm apart was used. Phosphorus at the rate of 30 kg ha⁻¹ was applied as basal dose. All other input and agronomic practices were carried out uniformly.

Procedures for data recording: Data on days to emergence was recorded by counting the days taken by each treatment from the date of sowing till the completion of emergence. Emergence m⁻² was noted by counting number of plants emerged in one meter row length at three randomly selected rows in each plot. Days to anthesis was recorded by counting days from date of sowing till about 50% of plants in subplot gave a clear indication of anthesis formation. Height of five plants in each treatment at random was measured at physiological maturity from soil surface to the tip of spike. Data on total and productive tillers m⁻² was recorded by counting tillers in three points of each sub plot and converted into m⁻². Spike length data was recorded by measuring spike length of six plants in each sub plot and averaged. Number of grain spike⁻¹ data was observed by counting the number of grain for five spikes randomly taken in each sub-plot and then averaged. After threshing, 1000 grain weight was recorded with the help of electronic balance at random from the produce of each subplot. Grain yield was recorded in four central rows in each subplot and then converted into kg ha⁻¹. The data on biological yield was taken on four central rows of each subplot

at maturity and then converted into kg ha^{-1} . Harvest index was calculated as percentage ratio of grain yield and biological yield. Nitrogen percentage in straw and grain was determined by kjeldhal method of Bremner and Mulvaney (1982).

Statistical analysis: All data are presented as mean values of three replicates. Data were analyzed statistically for analysis of variance (ANOVA) following the method described by Gomez & Gomez (1984). MSTATC computer software was used to carry out statistical analysis (Bricker, 1991). The significance of differences among means was compared by using Least Significant Difference (LSD) test (Steel & Torrie, 1997).

Results and Discussion

Plant growth and development: Varieties, nitrogen levels and their interactions did not significantly ($p > 0.05$) affected days to emergence and emergence m^{-2} (Figs. 1 and 2). Our results are in line with Le Gouis *et al.*, (1999) and Moselhy & Zahran (2002) who reported that nitrogen application had little or no effects on days to emergence. Days to anthesis, plant height and tillers m^{-2} was significantly ($p < 0.05$) affected by varieties and nitrogen levels. Interaction between varieties and nitrogen levels was non significant (Figs. 3, 4 and 5). Sterling variety took minimum days to anthesis as compared to local variety. Maximum days to anthesis (106 days) were recorded from plants which received 60 kg N ha^{-1} and beyond. Similar results were reported by Kernich and Halloran. (1996), who observed that nitrogen fertilizer considerably influenced duration of the pre-anthesis period and spike in barley. Our results indicated that taller plants (107.58 cm) were produced by local variety when compared with sterling (95.71 cm). Plant height was maximum (107.38 cm) when nitrogen was applied at the rate of 100 kg ha^{-1} , followed by 80 kg N ha^{-1} with plant height (103.88 cm). The probable reason might be that optimum nitrogen supply played an essential role in plant growth and development. Maqsood *et al.*, (2001), Kenbaev & Sade, (2002), Salwa *et al.*, (2005), Soyly *et al.*, (2005), Arif *et al.*, (2006) and Pervez *et al.*, (2009) reported significant increase in plant height of wheat with application of nitrogen.

More tillers m^{-2} was recorded in sterling as compared to local variety. In case of N levels, maximum tillers m^{-2} was produced with the application of 60 kg N ha^{-1} when compared with other treatments (Fig. 5). The probable reason might be that optimum nitrogen availability plays an essential role in plant growth whereas low or very high dose of nitrogen caused reduction in above ground vegetative growth of plant. Increase in the number of tillers m^{-2} of wheat due to N application was also reported by Rajput *et al.*, (1993) and Ahmad (1999). The result showed that lodging score was significantly ($p < 0.05$) affected by varieties, N levels and their interactions (Fig. 6). Maximum lodging was recorded with the application of 100 kg N ha^{-1} , followed by 80 kg N ha^{-1} . Mean value of the data also showed more lodging in local compared with sterling.

Yield and yield components: Significant ($p < 0.05$) differences were observed in productive tillers m^{-2} , spike length, grains spike $^{-1}$, 1000 grain weight, grain yield, straw yield, biological yield, harvest index, grain and straw N due to various levels of N and varieties (Figs. 8-16). More productive tillers m^{-2} (305.25), lengthy spikes (18.25 cm) highest grain spike $^{-1}$ (27.13), 1000 grain weight ((36.99 g), grain yield (2187 kg ha^{-1}) and biological yield (7481 kg ha^{-1}) was produced by the application of 60 kg N ha^{-1} . Similarly, grain and straw N was more (2.31% and 2.16% respectively) when N was applied at the rate of 60 kg N ha^{-1} . In case of varieties, sterling performed better than local for all the yield parameters except biological yield. Walter *et al.*, (1995) who observed that nitrogen

application significantly affected productive tillers m^{-2} . Cantero *et al.*, (1995), Le Gouis *et al.*, (1999) and Oweis *et al.*, (1999) observed similar results for grain spike $^{-1}$ in barley. Weight and number of grains spike $^{-1}$ was significantly increased with increasing N fertilization as reported by Moselhy & Zahran, (2002). They further revealed that application of nitrogen fertilizer significantly increased spike length, number of grains spike $^{-1}$, 1000 grain weight, grain yield and N uptake by the crop (Chaudhary & Mehmood, 1998; Bakhsh *et al.*, 1999; Tilahun *et al.*, 2000; Ennin & Clegg, 2001; Ahmad & Rashid, (2004); Imran *et al.*, 2005; Caglar & Ozturk, 2007; Zebarth *et al.*, 2009; Pervez *et al.*, 2009).




Fig. 1. Days to emergence of barley varieties as affected by different N levels. Bars shows LSD value at $p < 0.05$.




Fig. 2. Emergence m^{-2} of barley varieties as affected by different N levels. Bars shows LSD value at $p < 0.05$.




Fig. 3. Days to anthesis of barley varieties as affected by different N levels. Bars shows LSD value at $p < 0.05$.




Fig. 4. Plant height of maize varieties as affected by different N levels. Bars shows LSD value at p<0.05.




Fig. 8. Spike length (cm) of barley varieties as affected by different N levels. Bars shows LSD value at p<0.05.




Fig. 5. Tillers m² of barley varieties as affected by different N levels. Bars shows LSD value at p<0.05.




Fig. 9. Grains spike⁻¹ of barley varieties as affected by different N levels. Bars shows LSD value at p<0.05.




Fig. 6. Lodging (%) of barley varieties as affected by different N levels. Bars shows LSD value at p<0.05.




Fig. 10. Thousand grains weight of barley varieties as affected by different N levels. Bars shows LSD value at p<0.05.




Fig. 7. Productive tillers m² of barley varieties as affected by different N levels. Bars shows LSD value at p<0.05.




Fig. 11. Grain yield (kg ha⁻¹) of barley varieties as affected by different N levels. Bars shows LSD value at p<0.05.




Fig. 12. Straw yield (kg ha⁻¹) of barley varieties as affected by different N levels. Bars shows LSD value at p<0.05.




Fig. 16. Grain N (%) of barley varieties as affected by different N levels. Bars shows LSD value at p<0.05.




Fig. 13. Biological yield (kg ha⁻¹) of barley varieties as affected by different N levels. Bars shows LSD value at p<0.05.




Fig. 14. Harvest index (%) of barley varieties as affected by different N levels. Bars shows LSD value at p<0.05.




Fig. 15. Straw N (%) of barley varieties as affected by different N levels. Bars shows LSD value at p<0.05.

References

Ahmad, M.I., F.H. Taj and M.T. Jan. 1986. Effect of fertilizer levels and cutting frequency on the grain and green fodder yield of barley. *Sarhad. J. Agric.*, 2: 369-373.

Ahmad, N. 1999. Fertilizer scenario in Pakistan: policies and development. In: Proc. Conference on Agriculture Fertilizer Use by 2010. NFDC, Publication 2/99. Islamabad.

Ahmad, N. and M. Rashid. 2004. Fertilizer use in Pakistan, p: 74. NFDC, Planning and Development Division, Islamabad.

Amanullah, H. Rahman, Z. Shah and P. Shah. 2008b. Effects of plant density and N on growth dynamics and light interception of cereal. *Arch. Agron. Soil Sci.*, 54: 401-411.

Amanullah, R.A. Khattak and S.K. Khalil. 2008a. Effects of plant density and N on phenology and yield of cereal. *Plant. Nut. J.*, 32: 246-260.

Arif, M., M.A. Chohan, S. Ali, R. Gul and S. Khan. 2006. Response of wheat to foliar application of nutrients. *J. Agric. Bio. Sci.*, 1: 30-34.

Ayoub, M., S. Guertin, S. Lussier and D.L. Smith. 1994. Timing and levels of nitrogen fertility effects on spring wheat. *Crop Sci.*, 34: 748-750.

Bakhsh, A., A.H. Gurmani, A.U. Bhatti and H. Rehman. 1999. Effect of balanced application of N, P and K on the grain yield of wheat. *Sarhad. J. Agric.*, 15: 453-457.

Bertin, P. and A. Gallais. 2000. Physiological and genetic basis of nitrogen use efficiency in Maize. I. Agro-physiological results. *Maydica*, 45:53-66.

Bremner, J.M. and C.S. Mulvaney. 1982. Nitrogen-Total. In: *Methods of Soil Analysis*, (Eds.): A.L. Page, R.H. Miller, D.R. Keeney. Part II. Chemical and Microbiological Properties. Second Ed. *Am. Soc. Agron*, Madison, WI, USA, P. 595-624.

Bricker, B. 1991. MSTATC: A micro computer programe for the design, management and analysis of agronomic research experimentation. Crop and Sci. Deptt. MSU East Lansing Mi 48824 USA.

Caglar, P. and M.L. Ozturk. 2007. Response of wheat genotype to different levels of nitrogen fertilizer. *Eur. J. Agron.*, 11: 245-246.

Cantero, M.C., J.M. Villar, I. Romagosa and E. Fereses. 1995. Nitrogen fertilization of barley under semi arid rainfed conditions. *Eur. J. Agron.*, 4: 309-316.

Chaudhary, A.U. and R. Mehmood. 1998. Determination of optimum level of fertilizer nitrogen Ennin, S.A. and M.D. Clegg. 2001. Effect of soybean plant population in soybean and maize rotation. *Agron. J.*, 93: 396-403.

Gallais, A. and B. Hirel. 2004. An approach to genetics of nitrogen use efficiency in maize. *J. Exp. Bot.*, 55: 295-306.

Gomez, K.A. and A.A. Gomez. 1984. Statistical Procedures for Agricultural Research. 2nd Ed. John Willey & Sons, Inc. New York. 641.

Hayes, P.M., A. Castro, L.M. Cedillo, A. Corey, C. Henson, B.L. Jones, J. Kling, D. Matus, I. Rossi and K. Sato. 2003. Genetic

- diversity for quantitatively inherited agronomic and malting quality traits. Elsevier science publishers, Amsterdam.
- Imran, M., Z.A. Somroo, M. Qamar and Z.A. Gurmani. 2005. Effect of nitrogen and phosphorus on yield and yield contributing characteristics of wheat. *Sarhad. J. Agric.*, 21: 651-653.
- Kenbaev, B. and B. Sade. 2002. Response of field-grown barley cultivars grown on zinc-deficient soil to zinc application. *Comm. Soil Sci. Plant Anal.*, 33: 533-544.
- Kernich, G.C. and G.M. Halloran. 1996. Nitrogen fertilizer effects on the duration of the pre-anthesis period and spikelet number per spike in barley. *J. Agron.*, 177: 289-293.
- Le Gouis, J., O. Delebarre, D. Beghin, E. Heumez and P. Pluchard. 1999. Nitrogen uptake and utilization efficiency of two-row and six-row winter barley cultivars grown at two N levels. *Eur. J. Agron.*, 10: 73-79.
- Maqsood, M., A.A. Abid, A. Iqbal and M.I. Hussain. 2001. Effect of variable rates of nitrogen and phosphorus on growth and yield of maize. *J. Bio. Sci.*, 1: 19-20.
- Miao, Y., D.J. Mulla, C.P. Robert and J.A. Hernandez. 2006. Within-field variation in corn yield and grain quality responses to nitrogen fertilization and hybrid selection. *Agron. J.*, 98: 129-140.
- Moselhy, E.I. and M.A. Zahran. 2002. Effect of bio and mineral nitrogen fertilization on barley crop grown on a sandy soil. *Egypt. J. Agric. Res.*, 3: 921-936.
- Novoa, R. and R.S. Loomis. 1981. Nitrogen and plant production. *Plant and Soil*, 58: 177-204.
- Oikeh, S.O., V.O. Chude, G.J. Kling and W.J. Horst. 2007. Comparative productivity of nitrogen-use efficient and nitrogen-inefficient maize cultivars and traditional grain sorghum in the moist Savanna of West Africa. *African J. Agri. Res.*, 2: 112-118.
- Ortiz-Monasterio, J.L., K.D. Sayre, S. Rajaram and M. McMahan. 1997. Genetic progress in wheat yield and nitrogen use efficiency under four nitrogen rates. *Crop Sci.*, 37: 898-904.
- Oweis, T., M. Pala and J. Ryan. 1999. Management alternatives for improved durum wheat production under supplemental irrigation in Syria. *Eur. J. Agron.*, 11: 255-266.
- Pervez, K., M. Yousuf, M. Imitas and M. Islam. 2009. Response of wheat to foliar and soil application of urea at different growth stages. *Pak. J. Agron.*, 41: 1197-1204.
- Rajput, H.S., O.P. Lathwal, S. Tej and T. Singh. 1993. Effect of irrigation and nitrogen levels on yield attributions and yield of wheat. *Har. Agron. J.*, 8: 69-73.
- Salwa, S., B. Sade, A. Topal, N. Akgun and S. Gezgin. 2005. Responses of irrigated durum and bread wheat cultivars to boron application in low boron calcareous soil. *Turk. J. Agric.*, 29: 275-286.
- Scharf, P.C. and M.M. Alley. 1988. Nitrogen loss pathway and nitrogen loss inhibitors: A review. *J. Fert.*, 5: 109-125.
- Singh, V.P. and S.K. Uttam. 2000. Effect of farmyard manure and nitrogen levels on crop yield and economics of rice-wheat cropping system. *Crop Res.*, 5: 82-86.
- Steel, R.G.D. and J.H. Torrie. 1997. Principles and procedures of statistics: A biometrical approach. 2nd Edition McGraw Hill Book Co. Inc. New York.
- Tilahun, G., N. Kedir and T. Feyissa. 2000. Crop rotation effects on grain yield and yield components of bread wheat in the Bale highlands of southeastern Ethiopia. pp. 316-324. In: The Eleventh Regional Wheat Workshop for Eastern, Central and Southern Africa. CIMMYT, Addis Ababa, Ethiopia.
- Tripathi, S.C., K.D. Sayre, J.N. Kual and R.S. Narang. 2003. Lodging behavior and yield potential of spring wheat, effect of ethphon and genotypes. *Field Crop Res.*, 87: 207-220.
- Walter, E.B., C.B. Christianson and A.G. Lamothe. 1995. The effect of nitrogen fertilizer on growth, grain yield and yield component of malting barley. *Field Crop Res.*, 43: 87-99.
- Worku, M., B.E. Friesen, O.A. Diallob and W.J. Horst. 2007. Nitrogen uptake and utilization in contrasting nitrogen efficient tropical maize hybrids. *Crop Sci.*, 47: 519-528.
- Zapata, F. and O.V. Cleenput. 1986. Recovery of N¹⁵ labeled fertilizer by sugar beet spring wheat and winter sugar beet cropping sequences. *Fert. Res.*, 8: 269-178.
- Zebarth, B.J., P. Scout and M. Sharifi. 2009. Effect of straw and fertilizer nitrogen management for spring barley on soil nitrogen supply to a subsequent potato crop. *Amer. J. Res.*, 86: 209-217.

(Received for publication 15 October 2009)