

ACTION OF PLANT GROWTH REGULATORS IN ALLEVIATING SALINITY AND TEMPERATURE EFFECTS ON THE GERMINATION OF *PHRAGMITES KARKA*

AFSHEEN ZEHRA¹, BILQUEES GUL¹, RAZIUDDIN ANSARI¹, ABDEL REHMAN A. ALATAR²,
A.K. HEGAZY² AND M. AJMAL KHAN^{1,2,3*}

¹Institute of Sustainable Halophyte Utilization (ISHU), University of Karachi, Karachi-75270, Pakistan

²Department of Botany & Microbiology, College of Science, King Saud University, Riyadh, Saudi Arabia.

³Current address: Qatar Shell Professorial Chair for Sustainable Development & Professor, Department of International Affairs, College of Arts and Sciences, Qatar University, PO Box 2713, Doha, Qatar

Corresponding author's e-mail: ajmal.khan@qu.edu.qa

Abstract

High salinity, high temperature and absence of light may disturb the balance in endogenous growth regulators. This may put constraint on seed germination of halophytic grasses forcing them to adopt necessary measure like going into dormancy. Germination regulating chemicals may release dormancy imposed by such factors hence effect of Thiourea (10 mM), Nitrate (20 mM), Proline (0.1 mM), Betaine (0.1 mM), GA₃ (3 mM), Kinetin (0.05 mM) and Fusicoccin (5 μM) was studied in alleviating the inhibitory effect of a range of NaCl and temperature on seed germination of the halophytic grass *Phragmites karka*. Six NaCl concentrations (0, 100, 200, 300, 400 & 500 mM) were used in 12h light: 12h dark photoperiod and in complete darkness, at different temperature regimes (10/20°C, 15/25°C, 20/30°C and 25/35°C, the lower temperature corresponding to dark and the higher to light photoperiod). Highest number of seeds germinated in non-saline control and the seed germination decreased with increase in salinity at all temperature regimes. All growth regulators significantly promoted seed germination in saline and also in non-saline conditions at all thermoperiods, except at 25-35°C. Only GA₃ and fusicoccin successfully alleviated salinity enforced dormancy of seeds at this thermoperiod. Growth regulators promoted germination in darkness at all temperature regimes. Rate of germination was also significantly affected by the application of these chemicals. Salt induced dormancy of *P. karka* seeds was broken by the application of different growth regulators. These chemicals also alleviated the temperature (GA₃ and fusicoccin) and light (Nitrate) enforced dormancy from seeds of *P. karka*. It is concluded that germination regulating chemicals have differential effect on the seed germination of *P. karka*. Some of them may alleviate the dormancy while others have no effect.

Introduction

Seed germination is the first and probably the most sensitive stage in the life cycle of a plant. This attains even more significance in case of halophytes which are plants of the wild occupying saline habitats. Under such conditions, increase in NaCl concentrations may inhibit the germination of seed either by creating low osmotic potential, external to the seed preventing water uptake, or through the toxic effects of Na⁺ and Cl⁻ ions (specific ion effect) on the metabolic processes (Khajeh-Hosseini *et al.*, 2003; Kaya *et al.*, 2006). In addition to salinity, germination under such conditions may face hazards like shortage (drought) or excess of water (water logging causing anaerobic condition), temperature extremes etc. (Gulzar & Khan, 2002) resulting in reduced germination. To survive and perpetuate under such harsh conditions, halophytic seeds adopt the strategy of acquiring dormancy may it be innate, enforced or induced and wait for favorable conditions. Dormancy relieving and germination regulating chemicals are useful in improving seed germination under stressful conditions and may break the dormancy as well.

The germination promoting effect of growth hormones in breaking seed dormancy has been reported in many species (Macchia *et al.*, 2001; Duan *et al.*, 2004; El-Keblawy *et al.*, 2010; Bahrani & Pourreza, 2012). Germination regulating chemicals such as GA₃, kinetin, fusicoccin, proline, and betaine are reported to release dormancy (Ungar, 1977; Plyler & Proseus, 1996; Gulzar & Khan, 2001; El-Keblawy, 2012; Yarnia & Tabrizi, 2012). Compatible solutes, such as proline and betaine are known to play a role in the process of osmotic adjustment and they accumulate in plant under conditions of

environmental stresses. They also act as antioxidants by scavenging OH⁻ radical in oxidative stress (Heur, 2003; Hsu *et al.*, 2003). Proline is also involved in several physiological roles, such as stabilization of proteins, preservation from heat denaturation of enzymes, while betaine stabilizes the structure of a key protein like Rubisco (Thakur & Sharma, 2005).

Phragmites karka (Retz.) Trin. ex steud., is a perennial reed distributed in Pakistan, Tropical Africa, Polynesia, northern Australia and tropical Asia. It has creeping rhizomes, tall and erect culms up to 10 m high and occurs in swamps and beside streams of both coastal and inland region. *P. karka* is used as decoration, weaving material, and for making musical instruments. Freshly collected seeds of *P. karka* lack innate dormancy and germinate promptly under optimum conditions (0 mM NaCl; 20/30°C thermoperiod and 12-hour photoperiod) (Zehra & Khan, 2007). Salinity reduced germination but about 60% of the un-germinated seeds from high salt treatment (50 dS m⁻¹ NaCl) could recover after salinity was removed, indicating they had salt-enforced dormancy (Zehra & Khan, 2007). Keeping in view our recent studies indicating that *P. karka* may be a potential source of biofuel (Abideen *et al.*, 2001, 2012), the present study is designed to determine the effect of some dormancy relieving compounds in alleviating salinity, temperature and photoperiod effects in seeds of *P. karka*.

Materials and Methods

Inflorences of *P. karka* were collected during autumn 2004 from Karachi university campus. Seeds were cleaned and surface sterilized with 0.85% Clorox (sodium hypochlorite) for 1 minute before germination.

Various concentrations of NaCl (0, 100, 200, 300, 400 & 500 mM) and Thiourea (10 mM), Nitrate (20 mM), Proline (0.1 mM), Betaine (0.1 mM), Gibberillin (3 mM), Kinetin (0.05 mM), Fusicoccin (5 μ M) were used based on optimal level promoting seed germination reported for other species (Gulzar & Khan, 2002). Germination was carried out in 50 mm diameter tight fitting plastic Petri plates with 5 ml of test solution. Four replicates of 25 seeds each were used for each treatment. Seeds were germinated in programmed incubator (Percival, Boone, USA) with a 12h light: 12h dark photoperiod (Sylvania cool white fluorescent lamps, irradiance of 25 μ mol m⁻² s⁻¹, 400-700 nm) at different temperature regimes (10/20°C, 15/25°C, 20/30°C and 25/35°C). Percent germination was recorded on alternate days for 20 days. For observing response in complete darkness, Petri plates containing seeds were placed in black plastic bags and the germination was recorded at the 20th day. Rate of germination was calculated with the help of a modified Timson's germination velocity index = $\Sigma G/t$, where G is the percentage of seed germinated at 2-day intervals and t is the total germination period (Khan & Ungar, 2001a). The maximum possible values using our data were 50 (1000 seeds 20 days⁻¹). Germination data were arcsine transformed before statistical analysis using SPSS for windows release 9.0 (Anon., 1999). The differences among means were examined using three-way ANOVA and Bonferroni post-hoc test was used for significant differences between individual treatments.

Results

Four-way ANOVA of percentage germination indicated a significant ($p < 0.0001$) effect of salinity, temperature, light, chemicals and their interactions (table




Fig. 1. Effect of plant hormones (GA₃, 3 mM; Kinetin, 0.05 mM and Fusicoccin, 5 μ M) on final seed germination ($n = 4$; mean \pm s.e.) of *P. karka* under various salinities (0 to 500 mM NaCl) and photoperiods (Light: 12/12 h light/dark; Dark: 24 h dark).

not shown). A three way ANOVA showed significant effect of salinity, temperature, chemicals and their interactions on rate of germination (table not shown). Most seeds germinated in distilled water in 12 h photoperiod, however at cooler temperature regime all seed germinated while at warmer thermoperiod only 60 % seed germinated (Fig. 1). In dark, seed germination in distilled water was also inhibited in comparison to light germinated seeds and optimal seed germination was obtained at 20/30°C (Fig. 1). Increase in salinity inhibited germination but this inhibition was greater in complete darkness. Germination was severely inhibited at 25/35°C both in light and dark while only few seeds germinated at 500 mM NaCl at moderate temperatures (20/30°C) in light (Fig. 1).

Application of GA₃, kinetin and fusicoccin almost completely alleviated inhibitory effect of salinity on the seeds of *P. karka* at 10/20°C (Fig. 1). Seed germination enhanced from 0 to 70% at 500 mM NaCl in kinetin and up to 60% in GA₃ and fusicoccin and this effect decreased with increase in temperature in both light and dark treatment (Fig. 1). Fusicoccin and GA₃ also alleviated temperature, light and salinity effect at 25/35°C.

Osmotica like betaine and proline almost completely alleviated salinity effect in light at 10/20°C whereas with the increase in temperature this effect of osmotica decreased and it inhibited germination at highest temperature regime used (Fig. 2). For seed germinated in dark better results were obtained at moderate temperature regimes for both osmotica and partial alleviation was noticed at coolest regime and inhibitory effect at highest temperature regime (Fig. 2).




Fig. 2. Effect of plant osmotica (Betaine, 0.1 mM and Proline, 0.1 mM) on final seed germination ($n = 4$; mean \pm s.e.) of *P. karka* under various salinities (0 to 500 mM NaCl) and photoperiods (Light: 12/12 h light/dark; Dark: 24 h dark).

Nitrogenous compounds like thiourea and nitrate also partially to completely alleviated the salinity effects on the seed germination of *Phragmites karka* particularly at lower temperature regimes. With the increase in temperature there was no effect at 20/30°C with substantial inhibition at 25/35°C (Fig. 3). Similar trends were noted in dark germinated seeds however nitrate partially alleviated salinity effect at 20/30°C.




Fig. 3. Effect of nitrogenous compounds (Thiourea, 10 mM and Nitrate, 20 mM) on final seed germination ($n = 4$; mean \pm s.e.) of *P. karka* under various salinities (0 to 500 mM NaCl) and photoperiods (Light: 12/12 h light/dark; Dark: 24 h dark).

The effect of dormancy-regulating chemicals mentioned above on the rate of seed germination more or less followed the similar pattern as that of percent seed germination (Tables 1-3). Germination regulating chemicals did not improve the rate of germination under non-saline conditions except for GA_3 , kinetin, betaine and proline at 10/20°C. However, rate of germination under saline conditions was improved in all treatments at cooler temperature. At the higher temperature regimes of 15/25°C and 20/30°C the rate of germination was substantially improved at higher salinity concentrations except for fusicoccin (Tables 1-3). At warmer thermoperiod, magnitude of the improvement was much lower as compared to other temperature regimes.

Discussion

Seeds of *Phragmites karka* showed above 90% germination in non-saline control and were inhibited with an increase in salinity and no seed germinated at 500mM NaCl. Fusicoccin, GA_3 , kinetin, betaine, nitrate and thiourea partially to completely alleviated the inhibitory effects of salinity on the germination. These chemicals also compensated for the light need during seed germination. These effects of dormancy-regulating chemicals are greater at cooler thermoperiod while at 25/35°C seed germination was inhibited by dormancy-regulating chemicals treatment.

These chemical also improved seed germination prevented by cooler temperature.

Fusicoccin alleviated salinity induced dormancy in *P. karka*. Similar effect was also reported for the seeds of other halophytes (Gul & Khan, 2003; Khan *et al.*, 2004; El-Keblawy *et al.*, 2010; El-Keblawy, 2012). Fusicoccin may act as gibberellic acid to counter the effects of abscisic acid produced due to salinity effects. This alleviation may be due to the stimulation of ATPase production to facilitate proton extrusion and K^+ uptake (Antipova *et al.*, 2003). Fusicoccin is known to promote seed germination by the stimulation of cell elongation and cell division in germinating seeds (Obroucheva & Antipova, 1992). Exogenous fusicoccin promoted radical emergence by inducing initiation of cell elongation in hypocotyls and in radical of *Vicia faba* seeds (Antipova *et al.*, 2003).

Proline and betaine are reported to partially alleviate innate dormancy in *Zygophyllum simplex*, *Atriplex stocksii*, *A. prostrata*, *Halogeton glomeratus*, and *Halopyrum mucronatum* (Khan & Ungar, 1997; Khan *et al.*, 1998; Khan & Ungar, 2000, 2001c; Khan *et al.*, 2003). However, proline and betaine did not relieve salinity-induced dormancy in *Arthrocnemum indicum* (Khan *et al.*, 1998) *Kosteletzky virginica* (Poljakoff-Mayber *et al.*, 1994), *Halopyrum mucronatum* (Khan & Ungar, 2001c), *Salicornia rubra* (Khan *et al.*, 2002) and *Salicornia utahensis* (Gul & Khan, 2003). Both proline and betaine alleviated the innate dormancy of *Zygophyllum simplex* seeds but neither was effective at high salinities (Khan & Ungar, 1997) while Gul & Weber (1998) reported that proline and betaine both alleviated high salinity effects in *Allenrolfea occidentalis*. Our results with *P. karka* showed that both betaine and proline were very effective in alleviating salinity induced dormancy.

Thiourea and nitrate both stimulated the germination of *P. karka* seeds under saline conditions. Gul & Weber (1998) reported that nitrate and thiourea almost completely alleviated the seed germination of *Allenrolfea occidentalis* at all salinities used. A substantial promotion was also reported in *Triglochin maritima*, *Salicornia utahensis*, *Halogeton glomeratus*, *Sporobolus arabicus* and *Aeluropus lagopoides* (Khan & Ungar, 1999, 2001a,b; Gul & Khan, 2003; Gulzar & Khan, 2002). Some nitrogenous compounds such as nitric oxide, nitrate, nitrite and thiourea are known to stimulate the germination of seeds (Bethke *et al.*, 2004; Li *et al.*, 2005). Thiourea is also known to break dormancy and overcome the negative effect of temperature on seed germination (Esashi *et al.*, 1979; Aldosaro *et al.*, 1981).

Seed germination of *P. karka* was partially alleviated by the application of GA_3 and kinetin. The GA_3 is known to alleviate salinity effect in some halophytic seeds (Khan & Ungar, 1998; Khan *et al.*, 1998; Li *et al.*, 2005) while it was ineffective in other halophytes like *Suaeda fruticosa* and *Haloxylon recurvum* (Khan & Gul, 2006), *Sarcobatus vermiculatus* (Gul *et al.*, 2001), *Ceratoides lanata* (Khan *et al.*, 2004). Gibberellic acid was also found to be ineffective in alleviating salinity effects on the germination of number sub-tropical halophytes (Khan & Gul, 2006) however, it alleviated salinity effects in case of *Zygophyllum simplex* (Khan & Ungar, 1999) and *Halopyrum mucronatum* (Khan & Ungar, 2001c). Kinetin is also a more potent growth regulator known to alleviate salinity effects in a number of halophytes (Gul & Khan, 2003; Khan *et al.*, 2002, 2004; Li *et al.*, 2005; Khan & Gul, 2006).

Table 1. Effect of plant hormones (GA₃, 3 mM; Kinetin, 0.05 mM and Fusicoccin, 5 µM) on rate of seed germination (n = 4; mean ± s.e.) of *P. karka* under various salinities (0 to 500 mM NaCl) and 12/12 h light/dark photoperiod (Similar letters within each NaCl concentration indicate non-significant differences between treatments by Bonferroni test at $p < 0.05$).

	Salinity (mM)					
	0	100	200	300	400	500
10/20 °C						
NaCl	32 ± 2.69 ^a	23 ± 2.23 ^b	11 ± 1.25 ^c	5 ± 1.73 ^c	2 ± 0.80 ^b	0 ± 0.00 ^c
GA ₃	38 ± 0.60 ^a	37 ± 0.26 ^a	34 ± 3.00 ^a	30 ± 0.64 ^a	27 ± 0.95 ^a	20 ± 1.38 ^a
Kinetin	40 ± 0.64 ^a	36 ± 0.76 ^a	34 ± 0.41 ^a	33 ± 0.42 ^a	28 ± 0.92 ^a	27 ± 1.52 ^a
Fusicoccin	30 ± 4.84 ^a	29 ± 4.53 ^b	27 ± 2.97 ^b	23 ± 1.95 ^b	20 ± 1.15 ^a	13 ± 2.26 ^b
15/25 °C						
NaCl	41 ± 0.08 ^a	40 ± 0.51 ^a	27 ± 2.30 ^b	16 ± 1.15 ^c	13 ± 1.87 ^b	3 ± 1.18 ^c
GA ₃	36 ± 2.49 ^{ab}	35 ± 3.30 ^{ab}	34 ± 0.86 ^a	30 ± 0.54 ^a	23 ± 1.72 ^a	23 ± 1.45 ^a
Kinetin	35 ± 1.49 ^b	32 ± 1.64 ^b	32 ± 0.94 ^a	31 ± 1.98 ^a	27 ± 1.46 ^a	23 ± 0.76 ^a
Fusicoccin	29 ± 4.39 ^c	28 ± 0.78 ^c	27 ± 1.56 ^b	26 ± 3.17 ^b	25 ± 0.70 ^a	19 ± 2.30 ^b
20/30 °C						
NaCl	46 ± 1.41 ^a	43 ± 0.83 ^a	35 ± 1.05 ^{ab}	26 ± 0.51 ^b	19 ± 1.33 ^c	7 ± 1.90 ^b
GA ₃	34 ± 4.22 ^b	34 ± 2.14 ^b	28 ± 2.63 ^c	27 ± 0.75 ^b	25 ± 1.02 ^b	20 ± 2.14 ^a
Kinetin	35 ± 0.65 ^b	33 ± 2.47 ^b	32 ± 0.94 ^{bc}	30 ± 2.71 ^a	26 ± 0.51 ^b	22 ± 0.71 ^a
Fusicoccin	42 ± 0.70 ^a	41 ± 0.42 ^a	38 ± 0.57 ^a	35 ± 1.70 ^a	33 ± 0.17 ^a	24 ± 3.56 ^a
25/35 °C						
NaCl	46 ± 1.41 ^a	43 ± 0.83 ^a	35 ± 1.05 ^a	26 ± 0.51 ^a	19 ± 1.33 ^b	7 ± 1.90 ^b
GA ₃	34 ± 4.22 ^b	34 ± 2.14 ^b	28 ± 2.63 ^b	27 ± 0.75 ^a	25 ± 1.02 ^a	20 ± 2.14 ^a
Kinetin	35 ± 0.65 ^b	33 ± 2.47 ^b	32 ± 0.94 ^a	30 ± 2.71 ^a	26 ± 0.51 ^a	22 ± 0.71 ^a
Fusicoccin	22 ± 2.87 ^c	21 ± 2.48 ^c	18 ± 1.09 ^c	16 ± 4.53 ^b	13 ± 2.00 ^c	4 ± 1.12 ^b

Table 2. Effect of plant osmotica (Betaine, 0.1 mM and Proline, 0.1 mM) on rate of seed germination (n = 4; mean ± s.e.) of *P. karka* under various salinities (0 to 500 mM NaCl) and 12/12 h light/dark photoperiod (Similar letters within each NaCl concentration indicate non-significant differences between treatments by Bonferroni test at $p < 0.05$).

	Salinity (mM)					
	0	100	200	300	400	500
10/20 °C						
NaCl	32 ± 2.69 ^b	23 ± 2.23 ^b	11 ± 1.25 ^b	5 ± 1.73 ^b	2 ± 0.80 ^b	0 ± 0.00 ^c
Betaine	40 ± 0.83 ^a	38 ± 1.14 ^a	36 ± 1.28 ^a	33 ± 1.44 ^a	30 ± 1.42 ^a	29 ± 1.19 ^a
Proline	41 ± 0.22 ^a	40 ± 0.57 ^a	38 ± 0.33 ^a	34 ± 0.89 ^a	31 ± 1.28 ^a	23 ± 0.97 ^b
15/25 °C						
NaCl	41 ± 0.08 ^a	40 ± 0.51 ^a	27 ± 2.30 ^b	16 ± 1.15 ^b	13 ± 1.87 ^c	3 ± 1.18 ^b
Betaine	39 ± 0.33 ^a	34 ± 1.70 ^b	32 ± 1.93 ^{ab}	29 ± 3.38 ^a	26 ± 0.87 ^b	27 ± 0.81 ^a
Proline	42 ± 1.14 ^a	41 ± 1.24 ^a	38 ± 1.21 ^a	35 ± 0.41 ^a	32 ± 0.81 ^a	24 ± 2.50 ^a
20/30 °C						
NaCl	46 ± 1.41 ^a	43 ± 0.83 ^a	35 ± 1.05 ^a	26 ± 0.51 ^b	19 ± 1.33 ^b	7 ± 1.90 ^c
Betaine	38 ± 2.64 ^b	34 ± 0.49 ^b	33 ± 0.46 ^a	33 ± 0.44 ^a	31 ± 1.26 ^a	26 ± 1.07 ^a
Proline	33 ± 0.61 ^b	31 ± 0.73 ^b	29 ± 0.63 ^b	27 ± 1.02 ^b	27 ± 1.19 ^a	18 ± 1.64 ^b
25/35 °C						
NaCl	46 ± 1.41 ^a	43 ± 0.83 ^a	35 ± 1.05 ^a	26 ± 0.51 ^b	19 ± 1.33 ^c	7 ± 1.90 ^c
Betaine	38 ± 2.64 ^b	34 ± 0.49 ^b	33 ± 0.46 ^a	33 ± 0.44 ^a	31 ± 1.26 ^a	26 ± 1.07 ^a
Proline	33 ± 0.61 ^b	31 ± 0.73 ^b	29 ± 0.63 ^b	27 ± 1.02 ^b	27 ± 1.19 ^b	18 ± 1.64 ^b

Table 3. Effect of nitrogenous compounds (Thiourea, 10 mM and Nitrate, 20 mM) on rate of seed germination (n = 4; mean \pm s.e.) of *P. karka* under various salinities (0 to 500 mM NaCl) and 12/12 h light/dark photoperiod (Similar letters within each NaCl concentration indicate non-significant differences between treatments by Bonferroni test at $p < 0.05$).

	Salinity (mM)					
	0	100	200	300	400	500
10/20 °C						
NaCl	32 \pm 2.69 ^a	23 \pm 2.23 ^a	11 \pm 1.25 ^b	5 \pm 1.73 ^c	2 \pm 0.80 ^b	0 \pm 0.00 ^b
Thiourea	25 \pm 0.06 ^b	24 \pm 0.81 ^a	23 \pm 0.49 ^a	19 \pm 0.47 ^b	18 \pm 1.36 ^a	17 \pm 0.61 ^a
Nitrate	32 \pm 1.52 ^a	28 \pm 1.02 ^a	25 \pm 0.39 ^a	23 \pm 0.45 ^a	21 \pm 0.41 ^a	15 \pm 0.36 ^a
15/25 °C						
NaCl	41 \pm 0.08 ^a	40 \pm 0.51 ^a	27 \pm 2.30 ^b	16 \pm 1.15 ^c	13 \pm 1.87 ^b	3 \pm 1.18 ^c
Thiourea	41 \pm 0.59 ^a	40 \pm 0.19 ^a	40 \pm 0.10 ^a	38 \pm 0.54 ^b	30 \pm 1.30 ^a	29 \pm 0.46 ^b
Nitrate	44 \pm 0.48 ^a	44 \pm 0.34 ^a	44 \pm 0.43 ^a	42 \pm 0.43 ^a	34 \pm 0.74 ^a	34 \pm 0.62 ^a
20/30 °C						
NaCl	46 \pm 1.41 ^a	43 \pm 0.83 ^a	35 \pm 1.05 ^a	26 \pm 0.51 ^b	19 \pm 1.33 ^b	7 \pm 1.90 ^b
Thiourea	34 \pm 2.18 ^b	34 \pm 1.39 ^b	30 \pm 1.14 ^a	28 \pm 1.37 ^{ab}	28 \pm 0.53 ^a	22 \pm 0.44 ^a
Nitrate	37 \pm 2.55 ^b	34 \pm 0.54 ^b	34 \pm 0.59 ^a	31 \pm 1.37 ^a	30 \pm 0.97 ^a	27 \pm 0.31 ^a
25/35 °C						
NaCl	46 \pm 1.41 ^a	43 \pm 0.83 ^a	35 \pm 1.05 ^a	26 \pm 0.51 ^b	19 \pm 1.33 ^b	7 \pm 1.90 ^b
Thiourea	34 \pm 2.18 ^b	34 \pm 1.39 ^b	30 \pm 1.14 ^a	28 \pm 1.37 ^{ab}	28 \pm 0.53 ^a	22 \pm 0.44 ^a
Nitrate	37 \pm 2.55 ^b	34 \pm 0.54 ^b	34 \pm 0.59 ^a	31 \pm 1.37 ^a	30 \pm 0.97 ^a	27 \pm 0.31 ^a

Change in temperature also influenced the germination of *P. karka* showing inhibition at low and high temperature regimes. Temperature stress may inhibit seed germination through ABA biosynthesis and catabolism (Goani *et al.*, 2004). Low temperature (10/20°C) induced dormancy was broken by applying different growth regulators, but at high temperature regime (25/35°C) only GA₃ and fusicoccin alleviated the dormancy of seeds. There is no reported information about the effect of growth regulators in breaking temperature stress induced dormancy from seeds of halophytes.

Germination of many halophytes occurs when there is an optimal combination of day length, thermoperiod and salinity (Young *et al.*, 1980). Seeds of *P. karka* showed a light requirement for germination, they were able to germinate under light at high percentages even in salinity, while its germination was inhibited in complete darkness both in saline and non-saline conditions. Nitrate promoted germination or broke dark enforced dormancy of *P. karka* seeds from lower to moderate thermoperiods (10/30°C). Nitrate has been commonly used to break dormancy in seeds requiring light to germinate (Baskin & Baskin, 1998). For instance, seed germination of *Hypericum* spp that is affected by the absence of light was enhanced by KNO₃ application (Cirak *et al.*, 2007).

Phragmites karka which occupies a relatively wetter part of the coastal and inland areas around Karachi is highly tolerant to salinity at germination stage. In the present study its seeds exhibited dormancy in stress conditions and had a requirement of growth regulators to break this dormancy to allow seeds to germinate. Application of proline, betaine, thiourea, nitrate, fusicoccin, GA₃, and kinetin substantially alleviated the salinity effects on germination. The germination regulating chemicals were most effective at cooler temperature regime. Dark induced germination inhibition was also partially alleviated by the application of these chemicals. Higher temperature regimes decreased germination substantially where only GA₃ and fusicoccin had some alleviating effects. *Phragmites karka* could produce plenty of biomass in a waterlogged saline

soil which may be used to make bio-fuel, fodder, thatching material, baskets etc. This species could successfully be planted after monsoon rains and seedling emergence could be enhanced by priming the seed with various germination regulating chemicals.

Acknowledgement

The authors extend their appreciation to the deanship of scientific research at King Saud University for funding the work through the research group project No. RGP-VPP-175.

References

- Abideen, Z., R. Ansari and M.A. Khan. 2011. Halophytes: potential source of lignocellulosic biomass for ethanol production. *Biomass and Bioenergy*, 35: 1818-1822.
- Abideen, Z., R. Ansari, B. Gul and M.A. Khan. 2012. The place of halophytes in Pakistan's biofuel industry. *Biofuels*, 3: 211-220.
- Aldosaro, J., A. Mantilla and G. Nicholas. 1981. Effect of ABA, fusicoccin and thiourea on germination and K⁺ and glucose uptake in chickpea seeds of different temperatures. *Physiologia Plantarum*, 52: 353-362.
- Anonymous. 1999. *SPSS 9.0 for windows update*. SPSS Inc. USA.
- Antipova, O.V., L.M. Bartova, T.S. Kalashnikova, N.V. Obroucheva, V.D. Voblikova and G.S. Muromtsev. 2003. Fusicoccin-induced cell elongation and endogenous fusicoccin-like ligands in germination seeds. *Plant Physiol. & Biochem.*, 41: 157-164.
- Bahrani, A. and J. Pourreza. 2012. Gibberlic acid and salicylic acid effects on seed germination and seedlings growth of wheat (*Triticum aestivum* L.) under salt stress condition. *World App. Sci. J.*, 18: 633-641.
- Baskin, J.M. and C.C. Baskin. 1998. *Seeds: Ecology, Biogeography, and Evolution of Dormancy and Germination*. Academic Press, San Diego, California, USA.
- Bethke, P.C., M.R. Badger and R.L. Jones. 2004. Apoplastic synthesis of nitric oxide by plant tissues. *The Plant Cell*, 16: 332-341.

- Cirak, C., K. Kevseroglu and A.K. Ayan. 2007. Breaking of seed dormancy in a Turkish endemic *Hypericum* species: *Hypericum aviculariifolium* subsp. *depilatum* var. *depilatum* by light and some pre-soaking treatments. *J. Arid Environ.*, 68: 159-164.
- Duan, D., X. Liu, M.A. Khan and B. Gul. 2004. Effects of salt and water stress on the germination of *Chenopodium glaucum*. *Pak. J. Bot.*, 36: 793-800.
- El-Keblawy, A. 2012. Impacts of dormancy-regulating chemicals on innate and salinity-induced dormancy of four forage grasses native to Arabian deserts. *Grass and Forage Science*. doi: 10.1111/j.1365-2494.2012.00901.x
- El-Keblawy, A., F. Al-Ansari and N. Al-Shamsi. 2010. Impact of dormancy regulating chemicals on salinity induced dormancy in *Lasiurus scindicus* and *Panicum turgidum*: two desert glycophytic grasses. *Plant Growth Regulations*, 62: 163-170.
- Esashi, Y., Y. Ohara, M. Okazaki and K. Hishinuma. 1979. Control of cocklebur seed germination by nitrogenous compounds: Nitrite, nitrate, hydroxylamine, thiourea, azide, and cyanide. *Plant and Cell Physiology*, 20: 349-361.
- Goani, T., S. Kawahara, M. Tougou, S. Satoh, T. Hashiba, N. Hirai, H. Kawaide, Y. Kamiya and T. Yoshioka. 2004. Abscisic acid in the thermo-inhibition of lettuce seed germination and enhancement of its catabolism by gibberellin. *J. Experi. Bot.*, 55: 111-118.
- Gul, B. and D.J. Weber. 1998. Effect of dormancy compounds on the seed germination of non-dormant *Allenrolfea occidentalis* under salinity stress. *Annals of Bot.*, 82: 555-560.
- Gul, B. and M.A. Khan. 2003. Effect of growth regulators and osmotica in alleviating salinity effects on the germination of *Salicornia utahensis*. *Pak. J. Bot.*, 36: 877-886.
- Gul, B., M.A. Khan and D.J. Weber. 2001. Seed germination in *Sarcobatus vermiculatus*: A halophytic shrub from Great Basin desert. *Pak. J. Bot.*, 33: 473-482.
- Gulzar, S. and M.A. Khan. 2001. Seed germination of a halophytic grass *Aeluropus lagopoides*. *Ann. of Bot.*, 87: 319-324.
- Gulzar, S. and M.A. Khan. 2002. Alleviation of salinity-induced dormancy in perennial grasses. *Biologia Plantarum*, 45: 617-619.
- Heur, B. 2003. Influence of exogenous application of proline and glycinebetaine on growth of salt-stressed tomato plants. *Plant Sci.*, 165: 693-699.
- Hsu, S.Y., Y.T. Hsu and C.H. Kao. 2003. The effect of polyethylene glycol on proline accumulation in rice leaves. *Biologia Plantarum*, 46: 73-78.
- Ismail, A.M.A. 1990. Germination ecophysiology in population of *Zygophyllum qatarenses* Hadidi from contrasting habitats. *J. Arid Environ.*, 18: 185-194.
- Kaya, M.D., G. Okcu, M. Atak, Y. Cikili and O. Kolsarici. 2006. Seed treatments to overcome salt and drought stress during germination in sunflower (*Helianthus annuus* L.). *European J. Agron.*, 24: 291-295.
- Khajeh-Hosseini, M., A.A. Powell and I.J. Bingham. 2003. The interaction between salinity stress and seed vigor during germination of soybean seeds. *Seed Science and Technology*, 31: 715-725.
- Khan, M.A. and Gul, B. 2006. Halophyte seed germination. In: *Eco-physiology of High Salinity Tolerant Plants*. (Eds.): Khan, M.A. and D.J. Weber. Springer Publication, Netherlands. pp. 11-30.
- Khan, M.A. and I.A. Ungar. 1997. Alleviation of seed dormancy in the desert forb *Zygophyllum simplex* L., from Pakistan. *Annals of Bot.*, 80: 395-400.
- Khan, M.A. and I.A. Ungar. 1998. Seed germination and dormancy of *Polygonum aviculare* L. as influenced by salinity, temperature, and gibberellic acid. *Seed Science and Technology*, 26: 107-117.
- Khan, M.A. and I.A. Ungar. 1999. Seed germination and recovery of *Triglochin maritima* from salt stress under different thermoperiods. *Great Basin Naturalist*, 59: 144-150.
- Khan, M.A. and I.A. Ungar. 2000. Alleviation of salinity-enforced dormancy in *Atriplex griffithii* Moq. var. *stocksii* Boiss. *Seed Science and Technology*, 28: 29-37.
- Khan, M.A. and I.A. Ungar. 2001a. Role of dormancy regulating chemicals in release of innate and salinity-induced dormancy in *Sporobolus arabicus*. *Seed Science and Technology*, 29: 299-306.
- Khan, M.A. and I.A. Ungar. 2001b. Effect of dormancy regulating chemicals on the germination of *Triglochin maritima*. *Biologia Plantarum*, 44: 301-303.
- Khan, M.A. and I.A. Ungar. 2001c. Alleviation of salinity stress and the response to temperature in two seed morphs of *Halopyrum mucronatum* (Poaceae). *Aus. J. Bot.*, 49: 777-783.
- Khan, M.A., B. Gul and D.J. Weber. 2002. Improving seed germination of *Salicornia rubra* (Chenopodiaceae) under saline conditions using germination regulating chemicals. *Western North American Naturalist*, 62: 101-105.
- Khan, M.A., B. Gul and D.J. Weber. 2004. Action of plant growth regulators and salinity on the seed germination of *Ceratoides lanata*. *Canadian J. Bot.*, 82: 37-42.
- Khan, M.A., I.A. Ungar and B. Gul. 1998. Action of compatible osmotica and growth regulators in alleviating the effect of salinity on the germination of dimorphic seeds of *Arthrocnemum indicum* L. *Inter. J. Plant Sci.*, 159: 313-317.
- Khan, M.A., I.A. Ungar and B. Gul. 2003. Alleviation of salinity-enforced seed dormancy in *Atriplex prostrata*. *Pak. J. Bot.*, 36: 907-912.
- Lado, P., F. Rasi-Caldogno and R. Colombo. 1975. Promoting effect of fusicoccin on seed germination. *Physiologia Plantarum*, 34: 359-364.
- Li, W., X. Liu, M.A. Khan and S. Yamaguchi. 2005. The effect of plant growth regulators, nitric oxide, nitrate, nitrite and light on the germination of dimorphic seed of *Suaeda salsa* under saline conditions. *J. Plant Res.*, 118: 207-214.
- Macchia, M., L.G. Angelini and L. Ceccarini. 2001. Methods to overcome seed dormancy in *Echinacea angustifolia* DC. *Scientia Horticulturae*, 89: 317-324.
- Marre, E. 1979. Fusicoccin: a tool in plant physiology. *Annual Review of Plant Physiology*, 30: 273-278.
- Obroucheva, N.V. and O.V. Antipova. 1992. Fusicoccin as a probable endogenous factor of seed germination, *Doklady Akademii Nauk*, 325: 412-416.
- Plyler, D.B. and T.E. Proseus. 1996. A comparison of seed dormancy characteristics of *Saprtina alterniflora* (Poaceae). *Amer. J. Bot.*, 83: 11-14.
- Poljakoff-Mayber, A., G.F. Somers, E. Werker and J.L. Gallagher. 1994. Seeds of *Kosteletzkya virginica* (Malvaceae): Their structure, germination, and salt tolerance. II. Germination and salt tolerance. *Amer. J. Bot.*, 81: 54-59.
- Stout, R.G. 1998. Fusicoccin activity and building in *Arabidopsis thaliana*. *Plant Physiology*, 88: 999-1001.
- Ungar, I.A. 1977. Salinity, temperature, and growth regulator effects on seed germination of *Salicornia europaea* L. *Aquatic Bot.*, 3: 329-335.
- Yarnia, M. and E.F.M. Tabrizi. 2012. Effect of seed priming with different concentration of GA3, IAA and kinetin on Azarshahr Onion germination and seedling growth. *Journal of Basic and Applied Scientific Research*, 2: 2657-2661.
- Young, J.A., B.L. Kay, H. George and R.A. Evans. 1980. Germination of three species of *Atriplex*. *Agronomy Journal*, 72: 705-709.
- Zehra, A. and M.A. Khan. 2007. Comparative effect of NaCl and sea salt on germination of halophytic grass *Phragmites karka* at different temperature regimes. *Pak. J. Bot.*, 39: 1681-16, 94.