

## ECTOMYCORRHIZAE OF HIMALAYAN ALDER (*ALNUS NITIDA*) FROM THE MOIST TEMPERATE FORESTS OF PAKISTAN

MUHAMMAD HANIF<sup>1\*</sup> AND ABDUL NASIR KHALID<sup>2</sup>

<sup>1</sup>Department of Botany, GC University, Lahore, Pakistan

<sup>2</sup>Institute of Botany, University of the Punjab, Lahore, Pakistan

\*Corresponding author's email: [dr.mhanif@gcu.edu.pk](mailto:dr.mhanif@gcu.edu.pk)

### Abstract

The roots of the Himalayan alder were sampled from the rhizosphere and were investigated for ectomycorrhizal association. We described 11 ectomycorrhizal species viz; *Amphinema* sp. MHBYS-01, *Helvella* sp. MHELA-02, *Hydnobolites* sp. MHCER-02, *Inocybe* sp. MHNIT-01, *Inocybe* sp. MHNIT-02, *Inocybe* sp. MHOBL-01, *Peziza michelii*, *Sebacina* sp. MHAFF-01, *Tomentella* sp. MHSTU-01, *Tomentella* sp. MHSUB-01 and *Tylospora* sp. MHTYL-01 based on molecular phylogeny by using the Maximum Likelihood criterion. Some of the ectomycorrhizal morphotypes are also being described here on a morpho-anatomical basis.

**Key words:** Clade, Dichotomous, Khanspur, Pseudoparenchymatous, rDNA-ITS

### Introduction

*Alnus* Mill. comprises approximately 28-44 species and are widely distributed in the boreal and temperate zone of the Northern hemisphere (Rochet *et al.*, 2011). From Pakistan, only the *Alnus nitida* is reported from the Himalayan forests of Pakistan. It is native to the Himalayas and west Nepal.

*A. nitida* has many uses; decoction of its bark is used for the treatment of swellings and body pains (Gamble, 1972), tannin from the bark is used in dyeing. Its wood is used for construction, timber, firewood, as street tree and furniture (Gupta, 1945; Gamble, 1972; Khan *et al.*, 2020). *A. nitida* has been used as fodder (Rana *et al.*, 2019), medicinal, construction, furniture, fencing, roofing, utensils (Ahmad *et al.*, 2009), *A. nepalensis* used to treat Urination with bleeding in livestock and its extract is given orally (Nand and Naithani, 2018) and *A. nitida* has also been used as a pain reliever and has pain reducing the potency of catkin and cone extracts (Nagina & Ibrar, 2018).

Ectomycorrhizal status of the *Alnus* has been previously proved by a number of studies (Trappe 1964; Molina 1979; Miller *et al.*, 1991, 1992; Pritsch 1996, Pritsch *et al.*, 2000; Becerra *et al.*, 2005; Tedersoo *et al.*, 2009; Ostonen *et al.*, 2009; Kennedy and Hill 2010; McBurney *et al.*, 2017; Thiem *et al.*, 2018; Kilpeläinen *et al.*, 2019).

Alder (*Alnus*) belongs to the birch family Betulaceae. This genus is distributed throughout the north temperate zone. From Pakistan, only *Alnus nitida* (Himalayan alder) is found along the western Himalayas. This host plant remained unexplored for their belowground ectomycorrhizal partners from this region. There was only one report about the ectomycorrhizal evidence of *Alnus nitida* from Pakistan (Ashraf *et al.*, 2012). They described and illustrated ectomycorrhizae of *Peziza michelii*. In the present work additional 11 ectomycorrhizal morphotypes have been identified, described and illustrated with this photobiont.

### Material and Methods

**Sampling of ectomycorrhizae:** The Himalayan alder (*Alnus nitida*) roots were sampled from Ayubia National Park, KPK. Ectomycorrhizal roots were collected, vouchered and wrapped in polythene bags. The

morphotypes of *A. nitida* were manually sorted after removing the soil particles from the surface of the morphotypes and kept in 2% CTAB buffer for DNA extraction and in distilled autoclaved H<sub>2</sub>O for morpho-anatomic characterization. The selected morphotypes were characterized morpho-anatomically following Agerer (2002). Morphological and anatomical characterization of the ectomycorrhizal system was carried out under stereo and compound microscopes, photographed and illustrations were made with the help of Camera Lucida.

**Molecular characterization:** For molecular characterization, DNA was extracted from the selected morphotypes using the modified CTAB method given by Gardes & Bruns (1993). Amplification of the extracted DNA was performed using fungal-specific and universal primers (ITS1F, ITS1 and ITS4). Hot-start enzyme JumpStart (Sigma, St Louis, MO, USA) was used to catalyze the PCR with 2 min at 94°C, followed by 30 cycles of 30 s at 94°C, 30 s at 53°C, 40 s + 5 s per cycle at 72°C, and finishing with 5 min at 72°C. The PCR products were purified with QIAquick (Qiagen Inc., Valencia, CA, USA), sequenced bi-directionally using the reverse and forward primers and BigDye 3.1 on an ABI 3730 DNA sequencer (Applied Biosystems, Foster City, CA, USA) and edited in sequencher 4.5 (Gene Codes, Ann Arbor, MI, USA) in Jodrell Laboratory, Royal Botanical Gardens, Kew, UK. The DNA sequences were submitted to BLAST and used to query the nucleotide collection using default settings. The divergence in rDNA-ITS was measured by comparing sequence pairs reconstructed by using MegAlign (DNASTAR). DNA sequences obtained from *A. nitida* morphotypes were submitted in GenBank. These sequences were manually edited using MacClade 4.08 and Bioedit (version 7.0.9).

### Results

#### *Amphinema* sp. MHBYS-01 Plate 01 (Figs 1. A-E)

**Morphological characters:** ECTOMYCORRHIZAL SYSTEM simple to irregular, axis 30×50-1.0×1.0mm, UNRAMIFIED ENDS dichotomously branched, club-shaped, 3.0×5.0-1.0×1.0mm, young tips skin colored

while oxblood red at maturity. Texture smooth with a matte luster, no visibility of host tissue beneath the mantle sheath. EMANATING HYPHAE frequent, off-white in color, RHIZOMORPHS not observed.

**Anatomical characters of mantle:** MANTLE transitionally pseudoparenchymatous, OUTER MANTLE plectenchymatous, with irregularly arranged and densely packed hyphae (mantle type B; Agerer, 2006), up to 11.0-11.7µm in diameter, contents clear, pale yellow in color, septa and clamps absent, hyphal junctions absent, matrix material not visible. INNER MANTLE transitionally parenchymatous, hyphae intermingled with epidermoid cells, 8.0-14.0µm in length while 8.0-9.0µm in width, light brown in color, contents clear, ornamentation absent, matrix material not visible.

**Anatomical characters of emanating elements:** EMANATING HYPHAE frequent, transparent, straight, up to 2.1-2.4µm in diameter, contents clear, ornamentation absent, frequently septate, clamped, branching not observed.

***Helvella* sp. MHELA-02 Plate 01 (Figs. 2A-F)**

**Morphological characters:** ECTOMYCORRHIZAL SYSTEM simple to irregularly dichotomous, main axis 7.0×10.0-0.5×0.5mm, UNRAMIFIED ENDS bulbous to club-shaped, 2.5×3.0-1.0×1.0mm, tips light brown to beige in color. Texture of system smooth with a matte luster, host tissue not visible beneath the mantle sheath. Main axis of the Ectomycorrhizal system covered in a dense matte of short pointed echine like structures. EMANATING HYPHAE frequent, concentrated around the tips, white in color. RHIZOMORPHS not observed.

**Anatomical characters of mantle:** MANTLE pseudoparenchymatous, OUTER MANTLE pseudoparenchymatous with densely packed irregular to rectangular cells (mantle type K; Agerer, 2006), cells 4.0×4.7µm in width while 6.1×6.8µm in length. Cells pale yellow, contents clear, no matrix material, ornamentation absent. INNER MANTLE pseudoparenchymatous (type K; Agerer, 2006), cells irregular to angular, 3.8×4.6µm in width while 5.9×7.1µm in length, contents clear, no matrix material.

**Anatomical characters of emanating hyphae and Echine:** EMANATING HYPHAE frequent, straight, hyaline, up to 1.8-2.1µm in diameter, thin walled, moderately septate, clampless, contents clear, ornamentation absent. ECHINE LIKE STRUCTURES on the main axis, 47.2- 94.4µm in length, apex 2.5-2.6µm, mid 3.3-3.5µm and base 5.4-5.6µm in width, contents clear, encrusted with gelatinous material.

***Inocybe* sp. MHNIT-01 Plate 01 (Figs. 3A-F)**

**Morphological characters:** ECTOMYCORRHIZAL SYSTEM simple to irregular, axis 10.0×15.0-0.5×1.0mm, UNRAMIFIED ENDS simple to irregular, 5.0×7.0-0.5×1.0mm, color of system is rusty-brown; texture of system is coarsely grainy with a matte luster. No visibility of host tissue beneath the mantle sheath. EMANATING HYPHAE frequent, light brown, RHIZOMORPHS not observed.

**Anatomical characters of mantle:** MANTLE pseudoparenchymatous, OUETR MANTLE pseudoparenchymatous, with tightly packed angular cells (mantle type: K; Agerer, 2006), upto 7.9-8.9µm in length and 8.0-11.4µm in width, pale yellow in color, contents clear, matrix material absent. INNER MANTLE pseudoparenchymatous, densely packed angular cells, pale yellow to hyaline in color, contents clear, matrix material not observed.

**Anatomical characters of emanating elements:** EMANATING HYPHAE frequent, of two distinct types.

Type 1 is straight, dark brown in color, up to 2.1-3.2µm in width, thin-walled, rarely septate, clamps present, ramification present; contents clear except a few oil globules present, ornamentation absent.

Type 2 is straight to wavy, hyaline, up to 10.0-11.1µm in width, thin-walled, rarely septate, clamps absent, contents clear, thickly ornamented with echines, ramification present.

***Sebacina* SP. MHAFE-01 Plate 01 (Figs. 4A-F)**

**Morphological characters:** ECTOMYCORRHIZAL SYSTEM simple to irregular, axis 25.0×35.0-0.5×1.0mm, UNRAMIFIED ENDS simple to bulbous, 3.0×7.0-1.0×1.0mm, young tips light skin to beige in color while dark at maturity. Texture of system smooth with a matte luster. Host tissue not visible beneath the mantle sheath. EMANATING HYPHAE present, RHIZOMORPHS not observed.

**Anatomical characters of mantle:** MANTLE plectenchymatous, OUETR MANTLE plectenchymatous with irregularly arranged hyphae (mantle type B; Agerer, 2006), up to 3.1-5.1µm in width, hyaline, some hyphae terminally branched, contents clear, septa present, clamps absent, hyphal junctions absent, matrix material not visible. INNER MANTLE plectenchymatous, hyphae up to 3.2-4.7µm in width, contents clear, hyaline, septate, clamps absent, hyphal junctions absent, matrix material not visible.

**Anatomical characters of emanating elements:** EMANATING HYPHAE frequent, densely surrounding the ectomycorrhizal system. Two distinct types of emanating hyphae present.

Type 1 light brown, up to 5.7- 6.1µm in width, straight to wavy, thick-walled, rarely septate, unbranched, encrusted with gelatinous material, contents clear.

Type 2 hyaline, up to 1.8-2.1µm in width, straight, thin-walled, moderately septate, clamps present, ornamentation absent, contents clear.

***Tomentella* sp. MHSTU-01 Plate 01 (Figs. 5A-F)**

**Morphological characters:** ECTOMYCORRHIZAL SYSTEM simple to irregular, axis 7.0×11.0-0.5×1.0mm, UNRAMIFIED ENDS simple, 2.0×4.0-0.5×1.0mm, tips black in color, texture of system coarse with a matte luster. Host tissue not visible beneath the mantle sheath. EMANATING HYPHAE present, concentrated on tips. RHIZOMORPHS present.




Fig. 1. A-E. Morpho-anatomical features of *Amphinema* sp. MHBYS-01 (A-B) ECM morphotypes of *Amphinema* sp. MHBYS-01 (C) Transitionally pseudoparenchymatous outer mantle (D) Plectenchymatous Inner Mantle (E) Emanating Hyphae.




Fig. 2. A-F. Morpho-anatomical features of *Helvella elastica* (A-B) ECM morphotypes of *Helvella elastica* (C) Pseudoparenchymatous outer mantle (D) Pseudoparenchymatous Inner mantle (E) Emanating Hyphae (F) Scale-like Hairs.




Fig. 3. A-F: Morpho-anatomical features of *Inocybe nitidiscula* (A-B) ECM morphotypes of *Inocybe nitidiscula* (C) Pseudoparenchymatous outer mantle (D) Pseudoparenchymatous inner mantle (E-F) Emanating Hyphae.




Fig. 4. A-F: Morpho-anatomical features of *Sebacina* sp. MHAFF-01 (A-B) ECM morphotypes of *Sebacina* sp. MHAFF-01 (C) Plectenchymatous Outer mantle (D) Plectenchymatous Inner mantle (E-F) Emanating Hyphae.




Fig. 5. A-F: Morpho-anatomical features of *Tomentella* sp. MHSTU-01 (A-B) ECM morphotypes of *Tomentella* sp. MHSTU-01 (C) Plectenchymatous Outer mantle (D) Pseudoparenchymatous Inner mantle (E) Emanating Hyphae (F) Rhizomorph.

**Anatomical characters of mantle:** Mantle transitionally plectenchymatous OUTER MANTLE plectenchymatous, some irregularly shaped rounded cells also intermingled with hyphae (mantle type H; Agerer, 2006), cells up to 5.2–9.4 µm in length and 4.0–4.5 µm in width. Hyphae up to 2.6–3.1 µm in width, light brown in color, contents clear, hyphal junctions absent, matrix material not visible. INNER MANTLE pseudoparenchymatous, with hyphae intermingled with angular-shaped cells, up to 5.4–9.7 µm in length and 4.2–4.5 µm in width, pale brown in color, contents clear, hyphal junction absent, septa absent, matrix material not visible.

**Anatomical features of emanating elements:** EMANATING HYPHAE frequent, wavy to straight, up to 4.0–4.2 µm in width, dark brown, contents clear, frequently septate, thick-walled, clamps absent, ornamentation absent, ramification absent, anastomoses absent.

RHIZOMORPHS present, light brown, aseptate, clampless hyphae, neatly aggregated together, contents clear, gelatinous material abundant on hyphae, up to 118–141.6 µm in diameter with a restricted point of attachment.

#### ***Tomentella* sp. MHSUB-01 Plate 01 (Figs. 6A-E)**

**Morphological characters:** ECTOMYCORRHIZAL SYSTEM simple to irregular, axis 15.0×22.0–0.5×1.0 mm, UNRAMIFIED ENDS straight to club-shaped, 5.0×7.0–0.5×1.0 mm, color of tips blackish brown while black at maturity, texture of system is coarse with a matte luster. No visibility of host tissue beneath the mantle sheath. EMANATING HYPHAE infrequent, RHIZOMORPHS present.

**Anatomical characters of mantle:** MANTLE pseudoparenchymatous, OUTER MANTLE pseudoparenchymatous, with tightly arranged rectangular cells (mantle type L; Agerer, 2006), up to 7.2–8.0 µm in length while 7.7–7.8 µm in width, light brown in color, contents clear, ornamentation absent, matrix material not visible. INNER MANTLE pseudoparenchymatous, cells up to 7.0–7.7 µm in length and 7.5–7.8 µm in width, contents clear, ornamentation absent, matrix material not observed.

**Anatomical characters of emanating elements:** EMANATING HYPHAE infrequent, straight, hyaline, up to 10.0–13.0 µm in width, thin walled, moderately septate, clamped, ramification absent, contents clear, ornamentation absent. RHIZOMORPHS abundant, off-white, numerous hyphae septate hyphae aggregated together, up to 180.5–182.3 µm in width with the restricted point of attachment.

**Molecular and Phylogenetic Identification of fungal taxa:** From the living roots system of *A. nitida*, 12 fungal taxa were isolated, sequenced and BLAST searched for the closest match in GenBank (Table 1).

The distribution of these 12 phylotypes was based on the cladistic distribution of fungal taxa associated with *A. nitida*. The cladogram included ectomycorrhizal phylotypes distributed in 7 clades (Fig. 7).

In the Agaricales clade, we included 3 ectomycorrhizal fungal species from the genus *Inocybe* associated with this host tree. *Inocybe* sp. MHNIT-01 isolates (ENA16.42, ENA68.44, ENA76.41, 3ENA74.43 and 3ENA73.43) shared 97–100% analyzed genetic characters/nucleotides with each other and shared 93–95% analyzed genetic characters/nucleotides with similarity with *Inocybe nitidiuscula* (JF908249.1) while *Inocybe* sp. MHOB-01.2ENA52.58 shared 93% of analyzed genetic characters/nucleotides with *Inocybe oblectabilis* (AM882831.2). These phylotypes/OTUs from Pakistan grouped in clade Agaricales with strong (92%) bootstrap support (Fig. 1).

Atheliales clade was represented by the genus *Amphinema* and *Tylospora*. From Pakistan, *Amphinema* was first time being reported from belowground community associated with *A. nitida*. All isolates of *Amphinema* sp. MHBYS-01 (3ENA89.48, ENA60.45, ENA52.48, 3ENA90.48 and ENA15.49) shared 100% of analyzed genetic characters with each other and 83% analyzed genetic characters/nucleotides with *Amphinema byssoides* (AY219839.1) while *Tylospora* sp. MHTYL shared 99% of analyzed genetic characters with *T. fibrillosa* (HM1900017). The clade formed by Atheliales was not highly resolved by bootstrapping (54%) for the present analysis. Species of order Thelephorales grouped in two paraphyletic clades. Clade Thelephorales-I represented by *Pseudotomentella* species/isolates and Clade Thelephorales-II presented by *Tomentella* species. Both OUTs of *Pseudotomentella* sp. MHTRI (3ENA75.43 and 3ENA85.44) shared 98% of their analyzed genetic characters with each other and shared 89% analyzed genetic characters with *Pseudotomentella tristis* (AJ889968.1). Other clade of the Thelephorales (Clade-II) included only species of *Tomentella* and two OUTs from Pakistan clustered with strong (82%) bootstrap support. *Tomentella* sp. MHSTU.ENA77.41 shared 78% analyzed genetic characters with *T. stiposa* (EU819523.1) while *Tomentella* sp. MHSUB.2ENA09.40 shared 96% analyzed genetic characters with *T. subtestacea* (JQ711878.1).

Clade Sebaciales represented by only one phylotype (*Sebacia* sp. MHEPI-2ENA51.67) with strong (98%) bootstrap support from Pakistan. This morpho-phylotype shared 94.5% analyzed genetic characters with *Sebacia epigaea* (JQ665492.1).

Lastly, species of order Pezizales were also grouped in two clusters. *Helvella* sp. MHELA (2ENA08.40 and 2ENA07.40) shared 100% genetic characters with each other and 87% with *H. elastica* (AF335455.1). All phylotypes grouped together with strong bootstrapping (99%). Two morpho-phylotypes were represented in the second cluster (Clade Pezizales-II). *Hydnobolites* sp. MHCER01 (ENA17.42) shared 82% genetic characters with *H. cerebriformis* (EU784272.1) and *Peziza* sp. MHMIC (3ENA88.45) shared 99% genetic characters with *P. michelii* (JF908562.1) and all phylotypes grouped together with strong bootstrap (99%) support (Fig. 01).






Fig. 6. A-F. Morpho-anatomical features of *Tomentella* sp. MHSUB-01 (A-B) ECM morphotypes of *Tomentella* sp. MHSUB-01 (C) Pseudoparenchymatous Outer mantle (D) Pseudoparenchymatous Inner Mantle (E) Emanating Hyphae (F) Rhizomorph.

Table 1. rDNA-ITS sequences from the below-ground mycoflora of <i>Alnus nitida</i> with their closest matches in the Genbank and their country of origin.								
S. No.	Fungal species	Voucher No.	Accession No.	Closest match in GenBank	GenBank Accession No.	Origin country	Max. Identity (%)	Query coverage (%)
1.	<i>Amphinema</i> sp. MHBYS-01.	3ENA89.48	JN133916	<i>Amphinema byssoides</i>	AY219839.1	Canada	84%	99%
2.	<i>Helvella</i> sp. MHELA.	2ENA08.40	Awaiting	<i>Helvella elastica</i>	AF335455.1	Canada	92%	89%
3.	<i>Hydnobolites</i> sp. MHCER.	ENA17.42	Awaiting	<i>Hydnobolites cerebriformis</i>	EU784272.1	England	88%	90%
4.	<i>Inocybe</i> sp. MHNIT-01	ENA16.42	Awaiting	<i>Inocybe nitidiuscula</i>	JF908249.1	USA	95%	99%
5.	<i>Inocybe</i> sp. MHNIT-02	.ENA76.41	Awaiting	<i>Inocybe nitidiuscula</i>	JF908249.1	USA	94%	93%
6.	<i>Inocybe</i> sp. MHOBL-01	.2ENA52.58	Awaiting	<i>Inocybe oblectabilis</i>	AM882831.2	Sweden	95%	99%
7.	<i>Peziza michelii</i>	3ENA88.45	JN836749	<i>Peziza michelii</i>	JF908562.1	USA	99%	100%
8.	<i>Pseudotomentella</i> sp. MHTRI-01	.3ENA75.43	Awaiting	<i>Pseudotomentella tristis</i>	AJ889968.1	Denmark	93%	87%
9.	<i>Pseudotomentella</i> sp. MHTRI-02	.3ENA85.44	Awaiting	<i>Pseudotomentella tristis</i>	AJ889968.1	Denmark	100%	88%
10.	<i>Sebacina</i> sp. MHEPI	.2ENA51.67	Awaiting	<i>Sebacina epigaea</i>	JQ665492.1	Germany	95%	98%
11.	<i>Tomentella</i> sp. MHSTU.	ENA77.41	Awaiting	<i>Tomentella stupea</i>	EU819523.1	USA	79%	93%
12.	<i>Tomentella</i> sp. MHSUB.	2ENA09.40	Awaiting	<i>Tomentella subtestacea</i>	IQ711878.1	Canada	96%	99%

## Discussion

The Himalayan Alder's ectomycorrhizae were first documented by Ashraf *et al.*, (2012) from Pakistan. They reported *Peziza michelii* as ectomycorrhizal with this photobiont. Recently, Tapwal *et al.*, (2021) reported *Ramaria stricta* as ectomycorrhizal with this host. McBurney *et al.*, (2017) reported 22 ectomycorrhizal fungi from *Alnus* sp. (*A. rubra*). Kilpeläinen *et al.*, (2019) reported *Amanita muscaria*, *Hebeloma*, *Laccaria laccata*, *Lactarius aspideus*, *Leccinum aurantiacum* and *Paxillus involutus* were reported as ectomycorrhizal with another *Alnus* sp. (*A. incana*). Present investigation documents 11 ectomycorrhizae of Himalayan alder. These were characterized using the conventional and the modern molecular methods and were identified.

*Amphinema* sp. MHBYS-01 Ectomycorrhizae (closely matched with *Amphinema byssoides*) are being described and illustrated first time in association with *Alnus nitida*. Previously ectomycorrhizae of *A. byssoides* were reported with *Picea abies* (Agerer, 1987-1995; Berg, 1989; Danielson & Pruden, 1989; Haug & Pritsch, 1992; Ingleby *et al.*, 1990; Weiss, 1988, 1989, 1990; Weiss & Agerer, 1988), *Pinus* (Fassian de Vecchi, 1962; Haug *et al.*, 1994; Raidl, 1997), and *Quercus ilex* (Montecchio *et al.*, 2002). With *Picea abies*, it formed monopodial-pinnate ramification, with *Pinus* dichotomous, with *Quercus* irregularly pinnate to dichotomous-like. While with *Alnus nitida*, *Amphinema* sp. MHBYS-01, forms simple to irregular system with dichotomously branched unramified ends. Besides ramification, the mantle anatomy also differed when compared with morphotypes from other photobionts. *Picea abies*, *Pinus* and *Quercus ilex* morphotypes have plectenchymatous mantle organization while *Alnus nitida* has transitionally pseudoparenchymatous mantle organization. Despite this, *Amphinema* sp. MHBYS-01 has hyphae that lacked septa and clamps and also lacked rhizomorphs, young tips skin-colored, oxblood red at maturity, frequent emanating hyphae. The fruitbody of this fungus was previously found growing on the stump of *Pinus wallichianain* forests of Murree (Ahmad, 1972). Phylogenetically, *Amphinema* sp. MHBYS-01 grouped with *A. byssoides* with strong (96%) bootstrap support. All these morpho-anatomic and molecular phylogenetic differences are significant for delimiting *Amphinema* sp. MHBYS-01 from *A. byssoides*. *Amphinema* seems to generalist mycorrhizal forming

fungus as it is part of many ecologically different belowground communities (Kjøller *et al.*, 2008; Taylor & Finlay, 2003; Qian *et al.*, 1998; Taylor & Brand, 1992). *Picea abies* and *Pinus cembra* were also found to form ectomycorrhizal associations with *Amphinema* (Margit *et al.*, 2010). It was also found with *Pinus sylvestris* (Aučina *et al.*, 2007).

Ectomycorrhiza of the *Helvella* sp. MHELA-02 (closely matched with *Helvella elastica*) had light brown to beige color. Ectomycorrhizal system is characterized by simple to irregularly dichotomous with bulbous to club-shaped unramified ends.

Ectomycorrhiza of the *Helvella* sp. MHELA-02 (closely matched with *Helvella elastica*) can be compared with *Helvella* sp. Ho-TS601 which seems a close relative of *Helvella* sp. MHELA-02. Ectomycorrhizae of *Helvella* sp. Ho-TS601 was described by Tedersoo *et al.*, (2006) and was characterized by orange-brown to red-brown morphotypes and whitish tips when young. The mantle organization was cellular with large spherical to subepidermoid cells, without emanating hyphae. The species in genera *Helvella* are treated with unresolved trophic status (Hansen and Pfister 2006; Læssøe & Hansen, 2007). There was no previous molecular evidence about its trophic status. Tedersoo *et al.*, (2007a) designated this genus as ectomycorrhizal on the basis of stable isotopic analysis. O'Donnell *et al.*, (1997), Hansen & Pfister (2006) and Tedersoo *et al.*, (2006a) provided molecular (rDNA-ITS based) evidence about the mycorrhizal status of this group. On a similar rDNA-ITS barcoding basis, *Helvella* sp. MHELA-02 from Pakistan was assigned as mycorrhizal and its mycorrhizae are first time described and illustrated from Asia. This phylotype was recently reported as part of different belowground ectomycorrhizal communities of the Himalayan Moist temperate forests of Pakistan (Hanif, 2012).

We did not find any published description of ectomycorrhizae of *Inocybe nitidiuscula* (closely match of *Inocybe* sp. MHNIT-01) ectomycorrhizae to compare its morpho-anatomic features with *Inocybe* sp. MHNIT-01. Phylogenetically, this phylotype clustered with *I. nitidiuscula* sequences and thus was identified as a sister species to *I. nitidiuscula*. Previously, it was reported as ectomycorrhizal in the association of seedlings of *Pyrola chlorantha* (Hynson *et al.*, 2013).

*Sebacina* sp. MHAFF-01 (closely matched with *Sebacina incrustans*) was found growing as mycorrhizal with roots of *A. nitida*. Their mycorrhizal morphotypes differed significantly from *Sebacina incrustans*. Urban *et al.*, (2003) described mycorrhizae of *Sebacina incrustans* in the association of *Picea*. These ectomycorrhizae have monopodial-pyramidal ramification and lack rhizomorphs. The color of their mycorrhizal roots ranged from yellow pale cream or faintly yellowish to yellowish or white; mantle organization plectenchymatous, hyphae arranged net-like, repeatedly and squarrosely branched. On the other hand, *Sebacina* sp. MHAFF-01 ectomycorrhizae have simple to the irregular ectomycorrhizal system with simple to bulbous unramified ends. Members of

*Sebacinales* are part of many ectomycorrhizal communities (Takahide *et al.*, 2007; Selosse *et al.*, 2007; Martin *et al.*, 2009; Setaro *et al.*, 2006; Morris *et al.*, 2008; Smith *et al.*, 2007). Hanif (2012) also reported *Sebacinales* from Himalayan Moist Temperate Forests of Pakistan.

The mycorrhizal interactions in *Sebacinales* may have arisen from an ancestral endophytic habit by specialization (Weiß *et al.*, 2011). Considering their proven beneficial influence on plant growth and their ubiquity, endophytic *Sebacinales* may be a previously unrecognized universal hidden force in plant ecosystems.

In the present work, two more ectomycorrhizal morphotypes of the genus *Tomentella* are described and illustrated morpho-anatomically and phylogenetically. Ectomycorrhizal morphotypes of *Tomentella* sp. MHSTU-01 (closely matched with *Tomentella stuposa*) are simple to irregular with simple unramified ends with black tips, coarse texture of system with a matte luster. Phylogenetically, *Tomentella* sp. MHSTU-01 clustered with *Tomentella stuposa* (Fig. 01). Ectomycorrhizal morphotypes of *T. stuposa* were described previously (Agerer *et al.*, 2002; Jakucs *et al.*, 2005) with *Pinus* sp. These were brown in color and had monopodial-pinnate or monopodial-pyramidal ramification, rhizomorphs presence present, mantle pseudoparenchymatous with angular cells, bearing mounds of roundish cells (type K), emanating hyphae presence. *T. stuposa* ectomycorrhizae were also observed in association with *Salix reticulata* and *Dryas octopetala* communities (Ryberg *et al.*, 2009) and many other ectomycorrhizal communities (Cline *et al.*, 2005; Tedersoo *et al.*, 2003; Smith *et al.*, 2007; McCormick *et al.*, 2004). Hanif *et al.*, (2012) described and illustrated isolates of *Tomentella* spp. with *Cedrus deodara* from Pakistan.

Phylogenetically, *Tomentella subtestacea* (closely matched) appeared as sister species of *Tomentella* sp. MHSUB-01 (Fig. 01). Both these phylotypes differed significantly on the basis of shared analyzed genetic characters and rDNA-ITS genetic divergence. Ectomycorrhizae of the *Tomentella* sp. MHSUB-01 differed from *T. subtestacea* in having a simple to the irregular ectomycorrhizal system with straight to club-shaped unramified ends. The tips of the system are blackish brown while black at maturity, texture of the system is coarse with a matte luster. While, *T. subtestacea* has a monopodial-pinnate ectomycorrhizal system with straight unramified ends (Jakucs & Agerer, 2001). Previously *T. subtestacea* was reported it ectomycorrhizal from terrestrial Orchids below ground community (McCormick *et al.*, 2004).

The existence of the diverse types of ectomycorrhizal fungal species with *A. nitida* growing in the Himalayan moist temperate forests of Pakistan indicates its preference for more mycobiont partners. To complete the below-ground picture of the ectomycorrhizal communities associated with *A. nitida*, more comprehensive sampling would be required.




Fig. 7. Phylogenetic tree for Ectomycorrhizal MOTUs associated with *Alnus nitida* and rDNA-ITS sequences retrieved from GenBank.

## References

- Agerer, R. 1987-1995. *Colour Atlas of Ectomycorrhizae*. 1st-9th delivery. Einhorn, Schwäbisch Gmünd.
- Agerer, R. 2006. Fungal relationships and structural identity of their ectomycorrhizae. *Mycol. Prog.*, 5(2): 67-107.
- Agerer, R., R. Grote and S. Raidl. 2002. The new method 'micromapping', a means to study species-specific associations and exclusions of ectomycorrhizae. *Mycol. Prog.*, 1(2): 155-166.
- Ahmad, H., A.M. Khan, S. Ghafoor and N. Ali. 2009. Ethnobotanical study of upper Siran. *J. Herbs Spices Med. Plants.*, 15: 86-97.
- Ahmad, S. 1972. *Basidiomycetes* of Pakistan. Biological Society of Pakistan. *Monograph*, 6. pp. 141.
- Ashraf, T., M. Hanif and A.N. Khalid. 2012. *Peziza michelii* and its ectomycorrhizae with *Alnus nitida* (Betulaceae) from Pakistan. *Mycotaxon*, 120: 181-188.
- Aučina, A., M. Rudawska, T. Leski, A. Skridaila, E. Riepšasand and M. Iwanski. 2007. Growth and Mycorrhizal Community Structure of *Pinus sylvestris* Seedlings following the Addition of Forest Litter. *App. Environ. Microbiol.*, 73(15): 4867-4873.
- Becerra, A.G., L. Beenken, K. Pritsch, G. Daniele, M. Schlöter and R. Agerer. 2005. Morphological description and molecular characterization of *Lactarius aff. omphaliiformis*, *Russula alnijorullensis* and *Cortinarius tucumanensis* ectomycorrhizas on *Alnus acuminata*. *Mycologia*, 97: 1047-1057.
- Berg, B. 1989. Charakterisierung und Vergleich von Ektomykorrhizen gekalkter Fichtenbestände. Diss Univ München.
- Cline, E.T., J.E. Ammirati and R.L. Edmonds. 2005. Does Proximity to Mature Trees Influence Ectomycorrhizal Fungus Communities of Douglas Fir Seedlings? *New Phytol.*, 166(3): 993-1009.
- Danielson, R.M. and M. Pruden. 1989. The mycorrhizal status of urban spruce. *Mycologia*, 81(3): 335-341.
- Fassi, B. and E. de Vecchi. 1962. Ricerche sulle micorrize ectotrofiche del Pino strobo in vivaio. I. Descrizione di alcune forme più diffuse in Piemonte. *Allionia*, 8: 133-152.
- Gamble, J.S. 1972. *A Manual of Indian Timbers*. S. Low, Marston & Co. Ltd., London.
- Gardes, M. and T. Bruns. 1993. ITS primers with enhanced specificity for *Basidiomycetes*, application to the identification of mycorrhizae and rusts. *Mol. Ecol.*, 2: 113-118.
- Gupta, B.L. 1945. *Forest Flora of Chakrata, Dehra Dun and Saharanpur*. Forest Research Institute Press.
- Hanif, M. 2012. PhD Thesis. *Aphylllophorales and their morphotypes from Pakistan*. Department of Botany, University of the Punjab, Lahore, Pakistan.
- Hanif, M., A.N. Khalid and S. Sarwar. 2012. Additions to the ectomycorrhizae associated with himalayan cedar (*Cedrus deodara*) using rDNA-ITS. *Int. J. Agri. Biol.*, 14: 101-106.
- Hansen, K. and D.H. Pfister. 2006. Systematics of the *Pezizomycetes* - the operculate discomycetes. *Mycologia*, 98: 1031-1041.
- Haug, I. and K. Pritsch. 1992. Ectomycorrhizal types of spruce (*Picea abies* (L.) Karst.) in the Black Forest, a microscopical atlas. Kernforschungszentrum, Karlsruhe.
- Haug, I., R. Weber, F. Oberwinkler and J. Tschén. 1994. The mycorrhizal status of Taiwanese trees and the description of some ectomycorrhizal types. *Trees*, 8: 237-253.
- Hynson, A.N., M. Weiß, K. Preiss, G. Gebauer and K.K. Treseder. 2013. Fungal host specificity is not a bottleneck for the germination of *Pyroleae* species (*Ericaceae*) in a Bavarian forest. *Mol. Ecol.*, doi: 10.1111/mec.12180.
- Ingleby, K., P.A. Mason, F.T. Last and L.V. Fleming. 1990. *Identification of ectomycorrhizas*. ITE Research Publication no. 5. HMSO, London.
- Jakucs, E. and R. Agerer. 2001. *Tomentella subtestacea* Bourdot & Galzin + *Populus alba* L. *Descr. Ectomycol.*, 5: 213-219.
- Jakucs, E., G.M. Kovacs, R. Agerer, C. Romsics and Z. Erős-Honti. 2005. Morphological-amatomical characterization and molecular identification of *Tomentella stiposa* ectomycorrhizae and related anatomotypes. *Mycorrhiza*, 15(4): 247-258.
- Kennedy, P.G. and L.T. Hill. 2010. A molecular and phylogenetic analysis of the structure and specificity of *Alnus rubra* ectomycorrhizal assemblages. *Fungal Ecol.*, DOI:10.1016/j.funeco.2009.08.005.
- Khan, M.K., N. Muhammad, N. Uddin, N. Ali, M. Umer and S. Ullah. 2020. Genetic diversity in threatened plant species *Alnus nitida* (Spach.) Endel". *Plant Sc. Today*, 7(3): 314-318.
- Kilpeläinen, J., A. Barbero-López, B. Adamczyk, P.J. Aphalo and T. Lehto. 2019. Morphological and ecophysiological root and leaf traits in ectomycorrhizal, arbuscular-mycorrhizal and non-mycorrhizal *Alnus incana* seedlings. *Plant Soil*, 436: 283-297.
- Kjøller, R. and E.C. Karina. 2008. The impact of liming on ectomycorrhizal fungal communities in coniferous forests in Southern Sweden, *Rapport*, pp. 1-70.
- Læssøe, T. and K. Hansen. 2007. Truffle trouble: what happened to the *Tuberales*, *Mycol. Res.*, 111: 1075-1099.
- Margit, B., Z. Margit and P. Ursula. 2010. Ectomycorrhizal status of *Larix decidua*-, *Picea abies*- and *Pinus cembra*-nursery plants in South Tyrol. *Forest Obser.*, 5: 3-30.
- Martin, F., A. Aerts, D. Ahrén, A. Brun, E.G.J. Danchin, F. Duchaussoy, J. Gibon, A. Kohler, E. Lindquist and V. Pereda. 2008. The genome of *Laccaria bicolor* provides insights into mycorrhizal symbiosis. *Nature*, 452: 88-92.
- McBurney, K.G., Erica T. Cline, J.D. Bakker and G.J. Ettl. 2017. Ectomycorrhizal community composition and structure of a mature red alder (*Alnus rubra*) stand. *Fungal Ecol.*, 27: 47-58.
- McCormick, M.K., D.F. Whigham and J. O'Neill. 2004. Mycorrhizal Diversity in Photosynthetic Terrestrial Orchids. *New Phytol.*, 163(2): 425-438.
- Miller, S.L., C.D. Koo and R. Molina. 1991. Characterization of red alder ectomycorrhizae: a preface to monitoring belowground ecological responses. *Can. J. Bot.*, 69: 516-531.
- Miller, S.L., C.D. Koo and R. Molina. 1992. Early colonization of red alder and Douglas-fir by ectomycorrhizal fungi and Frankia in soils from the Oregon Coast Range. *Mycorrhiza*, 2: 53-61.
- Molina, R. 1979. Pure culture synthesis and host specificity of red alder mycorrhizae. *Can. J. Bot.*, 57: 1223-1228.
- Montecchio, L., S. Rossi, A. Grendene and R. Causin. 2002. *Amphinema byssoides* (Pers.: Fr.) J. Erikss. + *Quercus ilex* L. *Descr. Ectomyc.*, 6: 1-6.
- Morris, M.H., M.E. Smith, D.M. Rizzo, M. Rejmánek and C.S. Bledsoe. 2008. Contrasting Ectomycorrhizal Fungal Communities on the Roots of Co-Occurring Oaks (*Quercus* spp.) in a California Woodland. *New Phytol.*, 178(1): 167-176.
- Nagina and M. Ibrar. 2018. *In vivo* antinociceptive potential of ethanolic extract of leaves, catkin and cone of *Alnus nitida* (Spach.) Endl. *Int. J. Biosci.*, 13(4): 256-261.
- Nand, K. and S. Naithani. 2018. Ethnobotanical uses of wild medicinal plants by the local community in the Asi Ganga sub-basin, Western Himalaya. *J. Compl. Med. Res.*, 9(1): 34-46. 10.5455/jcmr.20180507034822



- O'Donnell, K., E. Cigelnik, N.S. Weber and J.M. Trappe. 1997. Phylogenetic relationships among ascomycetous truffles and false morels inferred from 18S and 28S ribosomal DNA sequence analysis. *Mycologia*, 89: 48-65.
- Ostonen, I., L. Tedersoo, T. Suvi and K. Lohmus. 2009. Does a fungal species drive ectomycorrhizal root traits in *Alnus* species? *Can. J. For. Res.*, 39: 1787-1796.
- Pritsch, K. 1996. Untersuchungen zur Diversität und Ökologie von Mykorrhizen der Schwarzerle (*Alnus glutinosa* (L.) Gaertn.) [Doctoral dissertation]. Tübingen: Univ Tübingen Press. 197 p.
- Pritsch, K., J.C. Munch and F. Buscot. 2000. Identification and differentiation of mycorrhizal isolates of black alder by sequence analysis of the ITS region. *Mycorrhiza*, 10: 87-93.
- Qian, X.M., I. Kottke and F. Oberwinkler. 1998. Influence of liming and acidification on the activity of the mycorrhizal communities in a *Picea abies* (L.) Karst. stand. *Plant & Soil*, 199: 99-109.
- Raidl, S. 1997. Studien zur Ontogenie an Rhizomorphen von Ektomykorrhizen. *Bibl. Mycol.*, 169: 1-84.
- Rana, D., A. Bhatt and B. Lal. 2019. Ethnobotanical knowledge among the semi-pastoral Gujjar tribe in the high altitude (Adhwari's) of Churah subdivision, district Chamba, Western Himalaya. *J. Ethnobiol. Ethnomed.*, 15:10 <https://doi.org/10.1186/s13002-019-0286-3>.
- Rochet, J., P.A. Moreau, S. Manzi and M. Gardes. 2011. Comparative phylogenies and host specialization in the alder ectomycorrhizal fungi *Alnicola*, *Alpova* and *Lactarius* (*Basidiomycota*) in Europe. *B.M.C. Evo. Biol.*, 11: 40.
- Ryberg, M., E. Larsson and U. Molau. 2009. Ectomycorrhizal Diversity on *Dryas octopetala* and *Salix reticulata* in an Alpine Cliff Ecosystem. *Arct. Antarct. Alp Res.*, 41(4): 506-514.
- Selosse, M.A., S. Setaro, F. Glatard, F. Richard, C. Urcelay and M. Weiß. 2007. *Sebacinales* are common mycorrhizal associates of *Ericaceae*. *New Phytol.*, 174(4): 864-878.
- Setaro, S., M. Weiß, F. Oberwinkler and I. Kottke. 2006. *Sebacinales* form Ectendomycorrhizas with *Cavendishia nobilis*, a Member of the Andean Clade of *Ericaceae*, in the Mountain Rain Forest of Southern Ecuador *New Phytol.*, 169(2): 355-365.
- Smith, M.E., G.W. Douhan and D.M. Rizzo. 2007. Ectomycorrhizal Community Structure in a Xeric *Quercus* Woodland Based on rDNA Sequence Analysis of Sporocarps and Pooled Roots. *New Phytol.*, 174(4): 847-863.
- Takahide, A.I., K. Nara and T. Hogetsu. 2007. Host effects on ectomycorrhizal fungal communities: insight from eight host species in mixed conifer-broadleaf forests. *New Phytol.*, 174(2): 430-440.
- Tapwal, A., K.S. Kapoor, Y. Thakur and A. Kumar. 2021. Ectomycorrhizal fungi associated with *Pinus gerardiana* in Kinnaur district of Himachal Pradesh, India. *Stud. Mycol.*, 6(1): 425-436.
- Taylor, A.F.S. and F. Brand. 1992. Reaction of the natural Norway spruce mycorrhizal flora to liming and acid irrigation. In: Read DJ, Lewis DH, Fitter AH, Alexander IJ, eds. *Mycorrhizas in Ecosystems*. Wallingford: C.A.B. International, 404.
- Taylor, A.F.S. and R.D. Finlay. 2003. Effects of liming and ash application on below ground ectomycorrhizal community structure in two Norway spruce forests. *Water Air & Soil Poll.*, 3: 63-76.
- Tedersoo, L., U. Collar, N. Hallenberg and K.H. Larsson. 2003. Fine scale distribution of ectomycorrhizal fungi and roots across substrate layers, including coarse woody debris in a mixed forest. *New Phytol.*, 159(1): 153-165.
- Tedersoo, L., K. Hansen, B.A. Perry and R. Kjoller. 2006. Molecular and morphological diversity of pezizalean ectomycorrhiza. *New Phytol.*, 170: 581-596.
- Tedersoo, L., P. Pellet, U. Kõljalg and M.A. Selosse. 2007. Parallel evolutionary paths to mycoheterotrophy in understory *Ericaceae* and *Orchidaceae*: ecological evidence for mixotrophy in *Pyroleae*. *Oecologia*, 151: 206-217.
- Tedersoo, L., T. Summer, T. Jairus, I. Ostonen and S. Pölme. 2009. Revisiting ectomycorrhizal fungi of *Alnus*: the differential host specificity, diversity and determinants of the fungal community. *New Phytol.*, 182: 727-735.
- Thiem, D., A. Piernik and K. Hryniewicz. 2018. Ectomycorrhizal and endophytic fungi associated with *Alnus glutinosa* growing in a saline area of central Poland. *Symbiosis*, 75: 17-28. DOI 10.1007/s13199-017-0512-5.
- Trappe, J.M. 1964. Mycorrhizal host and distribution of *Cenococcum graniforme*. *Lloydia*, 27: 100-106.
- Urban, A., M. Weiß and R. Bauer. 2003. Ectomycorrhizas involving sebacinoid mycobionts. *Mycol. Res.*, 107(1): 3-14.
- Weiss, M. 1988. Ektomykorrhizen von *Picea abies*. Synthese, Ontogenie und Reaktion auf Umweltschadstoffe. Diss Univ München.
- Weiss, M. 1989. *Amphinema byssoides*. In: (Ed.): Agerer, R. Colour Atlas of Ectomycorrhizae, plate 23, Einhorn-Verlag, Schwäbisch Gmünd.
- Weiss, M. 1990. Studien an Ektomykorrhizen XXV. Untersuchungen zur Ontogenie von Ektomykorrhizen an *Picea abies*. *Nova Hed.*, 50: 361-393.
- Weiss, M. and R. Agerer. 1988. Studien an Ektomykorrhizen XII. Drei nichtidentifizierte Mykorrhizen an *Picea abies* (L.) Karst. aus einer Baumschule. *Eur. J. For. Path.*, 18: 26-43.
- Weiß, M., Z. Sýkorová, S. Garnica, K. Riess and F. Martos. 2011. *Sebacinales* Everywhere: Previously Overlooked Ubiquitous Fungal Endophytes. *PLoS One*, 6(2):e16793. doi:10.1371/journal.pone.0016793.

(Received for publication 21 December 2021)