

GENOME SIZE ESTIMATION AND PHYSIOLOGICAL RESPONSES IN WHEAT (*TRITICUM AESTIVUM* L. VAR. YECORA ROJO) UNDER SALT STRESS FOR IN VITRO SELECTION OF TOLERANT LINES

MOHAMMAD NADEEM*, FAHAD AL-QURAINY, SALIM KHAN, ABDULRAHMAN AL-HASHIMI, MOHAMED TARROUM, HASSAN O. SHAIKHALDEIN, ABDALRHAMAN M. SALIH, ABDEL-RHMAN Z. GAA FAR, ARWA ALI AL-HARBI AND NAWAL F. AL-RASHEEDI

Department of Botany and Microbiology, College of Science, King Saud University, P.O. Box 2455, Riyadh 11451, Saudi Arabia

*Corresponding author's email: mnadeem@ksu.edu.sa

Abstract

In vitro callus culture of wheat (*Triticum aestivum* L. var. Yecora rojo) was established on MS medium supplied with different concentrations of 2, 4-D, and NAA (0.5-10 μ M). The maximum callus growth was observed in MS media with 5.0 μ M 2,4-D on a fresh weight basis. The effects of salt stress were studied on the callus of wheat. We analyzed biochemical markers and the response of the antioxidant system in callus. The callus fresh and dry weight were found to be maximum (fresh wt. 1.01 g and dry wt. 0.14 g on MS+50.0 mM NaCl). As the concentration of salt increased, the content of soluble protein, proline, and TBARS were higher in comparison to control. However, all these parameters decreased at the highest and enhanced salt concentration (200 mM NaCl) as of control. Similar observations were made in the case of antioxidant enzymes under salinity stress which were found to be highest at 150 mM NaCl as of control. The flow cytometry analysis exhibited no significant differences in genome size in treated and control calli, hence it was genetically stable with their genome size (2C DNA content) like control. The high salt tolerance of calli was evident from the accumulation of solute in the cell as of control.

Key words: Salinity stress, Callus, Biochemical markers, Genetic fidelity, Genome size

Introduction

Wheat (*Triticum aestivum* L.) is one of the most significant cereal crops globally, cultivated across diverse regions (Briggle & Curtis, 1987). It has earned the moniker "King of Cereals" due to its extensive cultivation, high yield, and prominent role in international grain trade (Anon., 2005). According to the NCBI Taxonomy Browser, wheat belongs to the family Poaceae, subfamily Poodeae, and tribe Triticeae.

A pressing issue that is likely to intensify in the future is salt stress resulting from soil salinization. This problem is primarily caused by over-fertilization, poor irrigation practices, and other factors. Statistics from the Food and Agriculture Organization (Anon, 2008) indicate that over 800 million hectares of land worldwide are currently affected by salinization, encompassing both saline and sodic soils, which constitute more than 6% of the global land area. The ongoing salinization of arable land is anticipated to have a substantial global impact, leading to a projected loss of 30% of agricultural land within the next 25 years and a 50% loss by 2050.

Wheat (*Triticum aestivum* L.) plays a crucial role in global food security, providing approximately 30% of the world's total cereal harvest and ranking second only to rice in terms of human calorie provision (Anon., 2006). Beyond its significant contribution of carbohydrates, fats, minerals, and vitamins, wheat offers a more fiber- and nutrient-rich dietary staple compared to meat-based diets (Šramková *et*

al., 2009). With the global population projected to approach 9 billion within the next 40 years (DESA UN, 2015), the demand for wheat is expected to increase substantially. While wheat is cultivated across diverse agroecosystems, its productivity is frequently constrained by a range of biotic and abiotic stresses. The area planted for wheat in Saudi Arabia has varied over time. According to the International Grains Council (IGC), Saudi Arabia's wheat production in 2019-20 will be 700,000 tons, up from 500,000 tons in 2018-19. Saudi Arabia's per capita wheat consumption is currently estimated to be 107 grams per day or approximately 39 kilograms per year. According to the Saudi Arabia Grains Organization (SAGO), Saudi Arabia consumed approximately 3.5 million tons of wheat in 2017-18. In addition, in 2017-18, 2.8 million tons of wheat flour were consumed. 2020 (World Grain). Saudi Arabia is a net wheat importer, purchasing an average of 1.1 million tons per year. Yecora rojo Aksada 59, and Aksada 67 are the most cultivated varieties.

Therefore, it is highly recommended to develop stress-tolerant and high-yielding varieties capable of growing in the shortest possible time. Tissue culture and gene transformation technology used in crop improvement. Overall, it is estimated that at least 3 ha of farming land is lost every minute worldwide because of soil salinization (Kundzewicz *et al.*, 2007). The total area of salinized soils worldwide is projected to be about 76.3 million ha, of which 41.5 ha are believed severely degraded (Oldeman *et al.*, 1990). Tissue culture procedures have made extensive

use of both mature and immature embryos; mature embryos were a better option than immature embryos (Šramková *et al.*, 2009). When it comes to regeneration, immature embryos are a better source for explants; however, they need time and growth facilities (Kundzewicz *et al.*, 2007), while mature embryos are available all year round. Mature embryos can be used directly (Šramková *et al.*, 2009) or dissected (Oldeman *et al.*, 1990). Plant tissue culture techniques provide a potential and feasible approach to developing salt-tolerant plants (Dracup, 1991) and (Tal, 1994) opted for *in vitro* selection of salt-tolerant cell lines for various species. Although studies on *in vitro* selection for salt tolerance in wheat have been carried out using mainly somaclonal variants (Barakat & Abdel-Latif, 1996; Karadimova & Djambova, 1993). A very few studies have been conducted on genotypic evaluation for callus production and *in vitro* selection of salt tolerance.

Wheat plants undergo many biochemical, molecular, and physiological mechanisms to adjust salinity stress at the tissue, cell, and whole plant to optimize the growth and yield by antagonizing the adverse saline environment. The present investigation was carried out to evaluate the effect of salt on wheat calli raised on MS medium using biochemical and molecular approaches.

Material and Methods

The experiments were performed in the growth room and plant tissue culture laboratory, at the Department of Botany and Microbiology, College of Science, King Saud University, Riyadh, KSA.

Plant material: Mature seeds of a bread wheat variety (*Triticum aestivum* L. var. Yecora rojo) were used as a starting explant.

Induction of callus from mature embryo: Fully developed embryos were extracted from the seeds of the selected bread wheat variety. The culture of mature embryos was carried out following the methods of (Salama *et al.*, 2013). Initially, the seeds underwent sterilization with 70% ethanol for a duration of 5 minutes. Subsequently, they were disinfected with Clorox, a commercial bleach solution containing 5% sodium hypochlorite, for 15 minutes. This disinfection process was enhanced by the addition of a few drops of Tween 80 to the solution. All procedures were conducted under sterile conditions within a Laminar Air Flow cabinet. Following disinfection, the seeds were thoroughly rinsed three times with sterile distilled water to eliminate any residual bleach and detergent. The seeds were then immersed in sterile distilled water and maintained at room temperature (22-25°C) in complete darkness for an overnight period. The following day, embryos were carefully excised from the soaked seeds using sterile scalpels and forceps. These embryos were then cultured on a variety of callus induction media to facilitate further experimentation. The callus induction media was MS salts and vitamins (Murashige & Skoog, 1962) supplemented with 3% (w/v) sucrose, solidified using 7.5 g l⁻¹ agar, and three replicates of different 2,4-dichlorophenoxyacetic acid (2,4-D) concentrations ranging from 1.0 to 10 µM for callus induction. After adjusting all media to a pH of 5.7±0.1, they were autoclaved for 20 minutes at 121°C. Each Petri plate

contained five embryos, which were then subcultured every four weeks while being kept in the dark at 25±2°C for a month. Weighing fresh callus tissue after four weeks of culture allowed us to calculate callus fresh weight (CFW), which is a measure of callus growth. The following formula is used to calculate growth, which is expressed as relative growth (RG): $W - W_0 / W_0 \times 100 = RG$ (Chen *et al.*, 2006; Meredith, 1978). W_0 indicates the callus's initial fresh weight, and W denotes the fresh weight after the culture passage.

Salt stress on callus cultures: One-month-old calluses were employed to initiate *in vitro* experiments for the selection of salt tolerance. For this purpose, the optimal callus induction medium was identified and supplemented with four concentrations of sodium chloride (NaCl) at 50, 100, 150, and 200 mM, with three biological replicates per treatment. Cultures were maintained in complete darkness at a temperature of 25 ± 2°C for one month. Callus fresh weight (CFW) was measured before and after salt treatment to evaluate the effects of salt stress on callus growth. Additionally, callus necrosis (CN) and *in vitro* callus tolerance (INTOL) were assessed using the following formula:

$CN = (\text{number of necrotic calli} / \text{total number of calli}) \times 100$ (Briggle & Curtis 1987) $CN = (\text{total number of calli} - \text{number of necrotic calli}) / \text{total number of calli} \times 100$ (Bouiamrine & Diouri, 2012). $INTOL = RGR_{\text{treatment}} / RGR_{\text{control}}$ (Al-Khayri & Al-Bahrany, 2004). Where RGR, the relative growth rate captured by the formula: $CRG = [LnW_2 - LnW_1] / GP$ (Birsin & Ozgen, 2004). Where W_1 and W_2 are the initial and final callus weights, respectively, and GP is the growth period.

Biochemical parameters

Proline content estimation: The proline content of the callus (0.25 g) was estimated using the technique adopted by (Hanson *et al.*, 1979). In five milliliters of 3% aqueous sulphosalicylic acid, the fresh samples were ground. The samples were centrifuged for 10 minutes at 5,000 rpm, and the supernatant was gathered to estimate the amount of proline. To the 2 ml of supernatant that was produced in the previous step, 2 ml of each of the acids ninhydrin and acetic acid were added. The mixture was incubated for one hour in boiling water before the reaction was halted in an ice bath. The mixture above was vortexed after 4 milliliters of toluene was added. Toluene containing the chromatophore was separated from the proline-containing mixture. By measuring the absorbance of a chromatophore containing toluene at 520 nm, the proline content of the samples was determined (Model UB-1800, Shimadzu, Japan).

Thiobarbituric acid reactive substances (TBARS):

The Cakmak and Horst method was used to evaluate the amount of TBARS in new calli (Cakmak & Horst, 1991). The 0.25 g callus samples were homogenized in 0.1% (w/v) trichloroacetic acid (TCA). To determine TBARS, the supernatant was extracted at 5,000 rpm. TBA was added to the above mixture at a rate of 0.5% (w/v) and left in a water bath at 90°C for 30 minutes. After stopping the reaction with ice, it was centrifuged for 10 minutes at

8,000 rpm. After collecting the supernatant, TBARS was measured using a spectrophotometer to measure absorbance at 600 and 532 nm wavelengths.

$$\text{TBARS (nmol g}^{-1} \text{fw)} = \frac{(A_{532} - A_{600}) \times V \times 100}{155 \times \text{extinction coefficient} \times w \times 1}$$

A532 represents absorbance at 532 nm, A600 represents absorbance at 600 nm, V = extraction volume, and W = fresh weight of tissue.

Estimation of antioxidant enzymes: Catalase was estimated to be one of the antioxidant enzymes in fresh calli. (EC 1.11.1.6) (Aebi, 1984) Superoxide dismutase (EC 1.15.1.1) (Dhindsa *et al.*, 1981), Ascorbate peroxidase (EC 1.11.1.11) (Nakano & Asada, 1981), and Glutathione reductase (EC 1.6.4.2) (Rao, 1992). The CAT, SOD, APX, and GR were recorded at 240nm, 550 nm, 290 nm, and 340 nm, according to a UV-vis spectrophotometer. The unit of enzyme was measured in mg⁻¹ protein min⁻¹.

Assessment of genetic stability: By using flow cytometry to measure the genome size (2C DNA content) of calli under salinity stress and comparing it to the genome size of control calli, the genetic stability of the organisms was evaluated.

Nuclei isolation Buffers: MB01 buffer developed by (Sadhu *et al.*, 2016) was used for the extraction of nuclei. It consists of different components including 20 mM MOPS; 25 mM Na₂ EDTA; 0.7 mM spermine.4HCl; 80 mM KCl; 20 mM NaCl; 1% (w/v) PVP; 0.5% (v/v) β-mercaptoethanol; 0.2% (v/v) and Triton X-100. The young calli was used to extract nuclei. Young leaves from the germination of the *Vicia faba* seed were used as an external reference. The seeds were a kind gift from Prof. J. Dolezel ~ (Laboratory of Molecular Cytogenetics and Cytometry, Institute of Experimental Botany, Olomouc Research Center, Czech Republic).

Extraction of nuclei: Nuclei were extracted from the fresh callus of wheat using the protocol reported by (Sadhu *et al.*, 2016). The calli (25 mg) were sliced with a sharp blade in a 600 µl in cold nuclei MB01 extraction buffer. The suspension of nuclei was pipetted slowly to prevent the formation of bubbles and then filtered through a 20 µm double nylon mesh filter. A 50 µg/ml PI stain was applied to the nuclei suspension (Propidium iodide, Sigma, USA) for 10 min and stored on ice before analysis.

Flow cytometric analysis: The amount of nuclear DNA in the wheat calli was determined using the method of Dolezel ~ *et al.*, (2007). The flow cytometer FACS Muse cell analyzer (Sigma, USA) was used to estimate the fluorescence of at least 5,000 propidium iodide-stained nuclei. The wheat genome size was calculated using the external reference *Vicia faba* (2C = 26.9 pg). The capillary's minimum flow rate was set at 0.12 µl/s. Muse cell analyzer software packages (Muse 1.8 analyses, USA) were used to compute the produced histograms. The following formula was used to determine the DNA content of the sample: 2C DNA content of *Triticum* =

$\frac{\text{Fluorescence mean intensity of Triticum}}{\text{Fluorescence mean intensity of standard}} \times 2C \text{ DNA content of standard}$. According to Bennett & Smith (1976), the equivalent of 1 pg DNA = 965 megabase pairs was used to determine the number of base pairs per haploid genome.

Statistical Analysis

SPSS software was used to statistically analyze the data. To ensure the accuracy of the results, the data was used in triplicate. They were then subjected to a one-way ANOVA and examined by Duncan's test. Different alphabet letters were used to indicate significant difference levels at $p \leq 0.05$.

Results

Studies on callus induction were conducted by testing two plant growth regulators, 2,4-D and NAA, at varying concentrations ranging from 0.5 to 10 µM. The effects of 2,4-D and NAA were assessed by culturing mature embryos of *Triticum aestivum* var. Yecora rojo on MS medium. Embryo growth and callus were initiated after a week of inoculation on all the PGRs and concentrations. Induction of callus and callus growth was calculated after 6 weeks of culture. Maximum induction and callus growth on a fresh weight basis (FW=0.95g) as compared to control (FW=0.22g) was observed on MS medium containing 5.0 µM 2,4-D (Fig. 1) whereas in other concentrations slow growth was noted and increasing the concentration reduced the callus proliferation (Plate 1 A, B, C). In the case of NAA treatment, maximum FW (0.75g) was increased and noted at 5 µM concentration and lowest in control FW (0.16g) (Fig. 2). The growth of calluses was found to be inhibitory at higher concentrations of both the PGRs evaluated. The callus cultures were maintained on an MS medium supplied with 5 µM of 2,4-D for subsequent investigations.

The effect of salinity stress was studied on calli raised on MS media containing 5.0 µM of 2,4-D under different concentrations of salt (50, 100, 150, and 200 mM NaCl). The impact of salt stress was investigated through biochemical and molecular approaches. Maximum fresh and dry weights were recorded at varying concentrations of NaCl (Plate 1D-I, Fig. 3). The proline and TBARS contents were found to be significantly higher at 150 mM NaCl compared to the control (Figs. 4 and 5). However, both parameters declined as the NaCl concentration increased further in the MS medium. At the highest salt concentration (200 mM NaCl) on MS media, all studied parameters (soluble protein, TBARS, proline, and antioxidant enzymes, Figs. 5-10) were decreased, however, all these parameters remained higher than the controls significantly. The addition of 50 and 100 mM NaCl to the media resulted in a higher accumulation of soluble protein in the calli compared to the control, although the difference between the two treatments was non-significant. The activities of CAT, SOD, APX, and GR peaked at 150 mM NaCl but declined as the NaCl concentration increased further to 200 mM in the MS medium. CAT activity at 50 mM NaCl showed no significant difference compared to the control. Similarly, GR activity at 100 and 150 mM NaCl also exhibited non-significant results.




Plate 1. Embryo culture, callus induction and salt stress to callus of wheat **A)** Embryo culture on MS media as control; **B)** Callus on 5 μ M NAA; **C)** Callus on 5 μ M 2,4-D; **D)** Callus on 5 μ M 2,4-D with different salt concentration; **E)** Callus on 5 μ M 2,4-D control for salt stress; **F)** Callus on 50 mM NaCl; **G)** Callus on 100 mM NaCl; **H)** Callus on 150 mM NaCl; **I)** Callus on 200 mM NaCl

The genetic stability was assessed in the calli under salinity stress by measuring the genome size (2C DNA content) with flow cytometry following the protocol of Sadhu *et al.*, (2006). The nuclei were extracted from both treated and untreated calli along with reference plants for genome size (2C DNA content) estimation (Fig. 11). The extracted nuclei with MB01 buffer showed a sharp peak as shown in the generated histogram (Fig. 11). The estimated genome size (2C DNA content) in treated and untreated callus was almost similar (Fig. 12).

Discussion

The overall impact of 2,4-D on the observed outcomes suggests that increasing concentrations of 2,4-D or NAA negatively affect callus growth and associated processes. This phenomenon may be linked to the accumulation of supra-optimal levels of NAA and 2,4-D within the tissues, which can impair growth performance. Consequently, MS media supplemented with 2,4-D yielded the highest mean values for callus induction. This observation may be explained by the mechanism through which it influences callus induction; for example, (Irfan Hafeez *et al.*, 2012) and (Ashraf & Osama, 2005) It has been reported that the significant challenge of inducing and sustaining callus in wheat, in particular, has been partially addressed by employing 2 mg/l of 2,4-D. In terms of callus induction and callus weight, mature embryos demonstrated a higher

frequency of callus formation and produced larger calluses (Tyankova & Zagorska, 2001). In mature embryo culture, the embryos exhibited a greater capacity to form callus structures. The increased hormone concentration correlated with enhanced callus production in mature tissue cultures (Kintzios *et al.*, 1996; Stals & Inzé, 2001).

In general, the induction of wheat callus in tissue cultures is influenced by several key factors, including the composition of the culture medium, the type of explant used, and the genetic characteristics of the genotype. (He *et al.*, 1989; Maddock *et al.*, 1983; Özgen *et al.*, 1998). In this study, our findings demonstrate that 2,4-D alone was effective in inducing yellow compact callus from mature embryos of wheat. The efficacy of 2,4-D for wheat callus induction has also been supported by earlier research (Abdrabou & Moustafa, 1993). Moreover, in other studies (Arzani & Mirodjagh, 1999; Barro *et al.*, 1998) 2,4-D has been widely recognized as the most commonly used growth regulator for callus induction and maintenance in wheat. Among the various concentrations of 2,4-D tested in this study, 5.0 μ M was found to yield the optimal results in terms of callus weight (Table 1). These outcomes align with the conclusions of (Yu *et al.*, 2008). It has been reported that the frequency of wheat callus induction reached its maximum with 2.0 mg/l of 2,4-D in the induction medium. In contrast, the response to NAA exhibited an inverse relationship, as higher NAA concentrations generally resulted in reduced performance across the traits studied.




Fig. 1. Variation in callus weight at different concentration of 2, 4 -D after six weeks of culture.




Fig. 2. Variation in callus weight at different concentration of NAA after six weeks of culture.




Fig. 3. Callus fresh and dry weight on different concentrations of salt after six weeks of culture.




Fig. 4. Proline content in calli under different concentrations of salt and control.




Fig. 5. TBARS content in calli under different concentrations of salt and control.




Fig. 6. Soluble protein content in calli under different concentrations of salt C (control), T-50 (50 mM NaCl), T-100 (100 mM NaCl), T-150 (150 mM NaCl), T-200 (200 mM NaCl).




Fig. 7. CAT activity in calli under different concentrations of salt C (control), T-50 (50 mM NaCl), T-100 (mM NaCl), T-150 (150 mM NaCl), T-200 (200 mM NaCl).




Fig. 8. SOD activity in calli under different concentrations of salt C (control), T-50 (50 mM NaCl), T-100 (mM NaCl), T-150 (150 mM NaCl), T-200 (200 mM NaCl).




Fig. 9. APX activity in calli under different concentrations of salt C (control), T-50 (50 mM NaCl), T-100 (mM NaCl), T-150 (150 mM NaCl), T-200 (200 mM NaCl).




Fig.10. GR activity in calli under different concentration of salt C (control), T-50 (50 mM NaCl), T-100 (mM NaCl), T-150 (150 mM NaCl), T-200 (200 mM NaCl).

Regarding the explant, these results are consistent with findings reported by several researchers, as (Zale *et al.*, 2004) who also demonstrated that mature embryos are capable of generating a sufficient number of regenerated plants. These findings are further supported by (Özgen *et al.*, 1998) according to whom mature embryos exhibit a high frequency of callus induction and strong regeneration potential. Consequently, due to their year-round availability, they are highly suitable for use in wheat tissue culture (Delparte *et al.*, 2001) utilized mature embryos successfully to produce calli, which aligns with the findings of this study. However, rather than employing whole seeds or embryos, they used fragmented embryo sections. (Yu *et al.*, 2008) also employed mature embryos in wheat tissue culture, achieving results consistent with those reported by numerous researchers (Ahmet & Adak, 2007; Bi *et al.*, 2007; Patnaik *et al.*, 2006).

Salinity stands as one of the most critical abiotic stress factors, significantly limiting crop productivity by adversely affecting plant growth. It disrupts the activity of antioxidant enzymes essential for nucleic acid and protein metabolism. Previous studies have established a strong correlation between reactive oxygen species (ROS) and plant responses to abiotic stresses. Consequently, understanding plant responses to salt stress, both *in vivo* and *in vitro*, is crucial. Under salinity stress, plants generate free radicals that inflict cellular damage. This stress impacts plant growth by modifying lipid composition, cell membrane integrity, and protein structure, while also inducing osmotic and ionic stress. (Cominelli *et al.*, 2013). In response to salinity stress, plants produce certain solutes that counteract free radicals and safeguard cells from damage. This study aimed to evaluate the impact of salt stress on wheat

callus by analyzing morphological, molecular, and biochemical markers. Morphological markers, such as callus size and weight, provide insights into growth rates under salinity stress. Variations in fresh and dry weight of calli were observed across different NaCl concentrations (50–200 mM). Changes in morphological traits of calli under varying salt concentrations may be attributed to the generation of free radicals, which disrupt cellular physiology. The highest biomass of calli, measuring 0.14 g, was observed under 50 mM NaCl treatment, surpassing both other treatments and the control (0.11 g, without salt). Callus growth on MS medium supplemented with 50 mM NaCl was superior to that observed at higher concentrations (100–200 mM NaCl). The reduced growth at 200 mM NaCl is likely due to the production of free radicals, which negatively impacted callus development.

Proline and Thiobarbituric Acid Reactive Substances (TBARS) levels in plants are helpful indicators of their ability to withstand abiotic stressors. Maintaining osmotic balance and shielding the cell structure from stress are the primary roles of osmolytes. (Polash *et al.*, 2019). In comparison to the control, the wheat callus displayed a greater accumulation of different solutes, such as soluble protein and proline contents, as supported by the results of other researchers where these parameters increased under salinity stress (Al-Qurainy *et al.* 2021; Aharaf & Foolad 2007). The accumulation of these compounds helps in maintaining the osmotic balance of the cell and also removes the free radicals from the cell, which are produced under salinity stress. Our outcome is validated by (Hannachi *et al.*, 2021) who conducted experiments on *Solanum melongena* callus under salinity stress, reported that lipid peroxidase activity increased in calli tolerant to 40 and 80 mM NaCl, but no such increase was observed in callus tolerant to 120 mM NaCl. (Al-Qurainy *et al.*, 2021) demonstrated that the accumulation of TBARS content was higher at 100 mM NaCl compared to 150 mM NaCl in date palm cultivars. Numerous studies have also indicated that proline levels increase significantly in plants subjected to salt stress (Calzone *et al.*, 2019; Vanlalruati *et al.*, 2019). Under salinity stress, the proline levels in *B. monnieri* cultured in tissue culture were six times higher compared to the control (Ali *et al.*, 1999). Proline accumulation is primarily driven by the enzymatic conversion of Pyrroline-5-carboxylate (P5C) to proline, catalyzed by Pyrroline-5-carboxylate reductase (P5CR) (Nazar *et al.*, 2015). Proline plays a crucial role in maintaining the structural integrity of macromolecules, particularly enzymes, by stabilizing hydration levels within the cell cytoplasm. (Foyer & Noctor, 2005).

A similar pattern was observed in the activities of antioxidant enzymes, such as SOD, CAT, APX, and GR, which mirrored the trend of osmolyte accumulation under salinity stress in wheat calli. The levels of these enzymes increased in a concentration-dependent manner, except at 200 mM NaCl, where a decline was noted compared to lower salt concentrations. The antioxidant enzymes

produced under abiotic stress detoxify the reactive oxygen species (ROS) from the cell (Gill & Tuteja 2010) and help in maintaining the growth and weight of the plant. Each of these enzymes plays a distinct role in protecting plant cells from reactive oxygen species (ROS) under salinity stress. SOD serves as a critical early indicator of plant sensitivity to ROS, acting as a first line of defense. In numerous NaCl-tolerant plants, increased SOD activity has been reported as a key adaptive response. (Gomez *et al.*, 2004). The antioxidant enzyme, GR plays an important role to ameliorate the tolerance of plants against environmental stresses (Hasanuzzaman *et al.*, 2020). The level of GR was reported to be high under salinity stress (Manai *et al.*, 2014). APX and CAT both enzymes overcome the toxic level of H₂O₂ in plants (Foyer & Noctor, 2005; Gomez *et al.*, 2004). The ROS inhibition system and plants' ability to withstand alkaline and salinity stressors are both significantly impacted by APX. The plant's high CAT activity demonstrated its tolerance as a result of increased H₂O₂ detoxification. (Mittova *et al.*, 2002). The CAT activity depends on many parameters such as metabolic status and growth of the plant, time, and intensity of stresses (Sriniegn *et al.*, 2015). The CAT and SOD activity was increased in NaCl-tolerant calli of *Solanum melongena* under salinity stress (Hannachi *et al.*, 2021).

Regardless of the information that is encoded, it is believed that the size of the genome affects plant characteristics at the subcellular and organismal levels. (Bennett, 1971) and can interact with environmental stressors to affect both plant responses and traits (Suda *et al.*, 2015) (Bennett & Leitch, 2005). In our study, the genome size (2C DNA content) was observed to be the same in treated and untreated calli and no significant difference was to be detected. The histograms generated from treated and untreated calli using the MB01 buffer were almost similar (Figs. 11 and 12). Thus, the applied concentration of salt on calli grown on MS media has not been observed any genotoxic effect on genome size (2C DNA content). However, the growth of calli at the highest concentration of salt (200 mM NaCl) was reduced along with variation in biochemical parameters, and hence the growth of calli was inferior to the control and in other salt treatments.




Fig. 11. Histogram generated from untreated calli by flow cytometry using the MB01 buffer.




Fig. 12. Histogram generated from calli under different concentrations of salt treatment (a: 50 mM NaCl; b: 100 mM NaCl; c: 150 mM NaCl; d: 200 mM NaCl) by flow cytometry using the MB01 buffer.

Conclusion

In comparison to NAA, MS media supplemented with 2,4-D demonstrated superior callus growth, as evidenced by higher fresh weight gain after one month of culture. Salt stress experiments were subsequently conducted on this vigorously growing callus. The results revealed that salinity stress in the culture medium significantly restricted callus growth, as evaluated through morphological, biochemical, and molecular markers. All analyzed parameters, including soluble protein content, TBARS, proline levels, and antioxidant enzyme activities, were elevated compared to the control. Genetic fidelity in the callus under salinity stress was assessed by measuring genome size (2C DNA content) using flow cytometry, which showed no significant variation across salinity treatments compared to the control callus. These findings highlight the need for further research to evaluate salinity tolerance in wheat calli, which could contribute to the development of salt-tolerant wheat varieties in future studies.

Funding

The authors extend their appreciation to the Ongoing Research Funding program (ORF-2025-73), King Saud University, Riyadh, Saudi Arabia.

References

- Abdrabou, R.T and R. A.K. Moustafa. 1993. Effect of 2, 4-D concentrations and two levels of sucrose on callus induction and plantlet formation in two wheat genotypes. *Ann. Agric. Sci. Cairo*. 1 (special issue):41-46.
- Aebi, H. 1984. Catalase in vitro. In: *Methods in enzymology* (Vol. 105, pp. 121-126). Elsevier.
- Ahmet, H. and M.S. Adak. 2007. Callus induction and plant regeneration in some Iraqi common wheat varieties. *Tarim Bilimleri Dergisi*, 13 (3): 285-292.
- Ali, G., P. Srivastava and M. Iqbal. 1999. Proline accumulation, protein pattern and photosynthesis in *Bacopa monniera* regenerants grown under NaCl stress. *Biol. Plant.*, 42: 89-95.
- Al-Khayri, J. and A. Al-Bahrany. 2004. Growth, water content, and proline accumulation in drought-stressed callus of date palm. *Biol. Plant.*, 48(1): 105-108.
- Al-Qurainy, F., S. Khan, M. Nadeem, M. Tarroum and A.R.Z. Gaafar. 2017. Antioxidant System Response and cDNA-Scot Marker Profiling in *Phoenix dactylifera* L. Plant Under Salinity Stress. *Int. J. Genom.*, 2017(1): 1537538.
- Al-Qurainy, F., S. Khan, S. Alansi, M. Nadeem, A. Alshameri, A.-R. Gaafar, M. Tarroum, H. O. Shaikhaldein, A.M. Salih and N.A. Alenezi. 2021. Impact of Phytomediated Zinc Oxide Nanoparticles on Growth and Oxidative Stress Response of In Vitro Raised Shoots of *Ochradenus arabicus*. *Biomed Res Int.*, Dec 6:2021:6829806. doi: 10.1155/2021/6829806. eCollection 2021.
- Anonymous. 2005. Crop Production Statistics. Food and Agricultural Organization of the United Nation.

- Anonymous. 2006. Supply C, Brief D. Food and Agriculture Organization of the United Nation. <http://www.fao.org/worldfoodsituation/csdb/en>.
- Anonymous. 2008. Land and plant nutrition management service. Food and Agricultural Organization of the United Nations. <http://www.fao.org/ag/agl/agll/spush>.
- Arzani, A. and S.S. Mirodjagh. 1999. Response of durum wheat cultivars to immature embryo culture, callus induction and *In vitro* salt stress. *Plant Cell Tiss. Org.*, 58: 67-72.
- Ashraf, F. and E.S. Osama. 2005. Improvement of plant regeneration from long-term callus cultures of two Egyptian wheat cultivars. *Arab J. Biotech.*, 8(1): 177-188.
- Ashraf, M. and M.R. Foolad. 2007. Roles of glycine betaine and proline in improving plant abiotic stress resistance. *Environ. Exp. Bot.*, 59(2): 206-216.
- Barakat, M. and T. Abdel-Latif. 1996. *In vitro* selection of wheat callus tolerant to high levels of salt and plant regeneration. *Euphytica*, 91: 127-140.
- Barro, F., M. Cannell, P. Lazzeri and P. Barcelo. 1998. The influence of auxins on transformation of wheat and tritordeum and analysis of transgene integration patterns in transformants. *Theor. Appl. Genet.*, 97: 684-695.
- Bennett, M.D. 1971. The duration of meiosis. *Proceedings of the Royal Society of London. Series B. Biol. Sci.*, 178(1052): 277-299.
- Bennett, M.D. and I.J. Leitch. 2005. Genome size evolution in plants. In: *The evolution of the genome* (pp. 89-162). Elsevier.
- Bi, R., M. Kou, L. Chen, S. Mao and H. Wang. 2007. Plant regeneration through callus initiation from mature embryo of Triticum. *Plant Breeding*, 126(1): 9-12.
- Birsin, M.A. and M. Ozgen. 2004. A comparison of callus induction and plant regeneration from different embryo explants of triticale (x Triticosecale Wittmack). *Cell Mol. Biol. Lett.*, 9(2): 353-362.
- Bouiamrine, E. and M. Diouri. 2012. Response of durum wheat (*Triticum durum* Desf.) callus culture to osmosis-induced drought stress caused by polyethylene glycol (PEG). *Ann. Biol. Res.*, 3(9):4555-4563.
- Briggle, L.W. and B.C. Curtis. 1987. Wheat worldwide. In: (Ed.): E.G. Heyne. *Wheat and wheat improvement*, 2nd ed. Agronomy 13: 1-32
- Cakmak, I. and W.J. Horst. 1991. Effect of aluminium on lipid peroxidation, superoxide dismutase, catalase, and peroxidase activities in root tips of soybean (*Glycine max*). *Physiol. Plant.*, 83(3): 463-468.
- Calzone, A., A. Podda, G. Lorenzini, B.E. Maserti, E. Carrari, E. Deleanu, Y. Hoshika, M. Haworth, C. Nali and O.Bade. 2019. Cross-talk between physiological and biochemical adjustments by Punica granatum cv. Dente di cavallo mitigates the effects of salinity and ozone stress. *Sci. Total Environ.*, 656: 589-597.
- Chen, J.Y, R.Q. Yue, H.X. Xu and X.J. Chen. 2006. Study on plant regeneration of wheat mature embryos under endosperm-supported culture. *Agri. Sci. China*, 5(8): 572-578.
- Cominelli, E., L. Conti, C. Tonelli and M. Galbiati. 2013. Challenges and perspectives to improve crop drought and salinity tolerance. *New Biotechnol.*, 30(4): 355-361.
- Delporte, F. O. Mostade and J. Jacquemin. 2001. Plant regeneration through callus initiation from thin mature embryo fragments of wheat. *Plant Cell Tiss. Org.*, 67:73-80.
- Dhindsa, R.S., P. Plumb-Dhindsa and T.A. Thorpe. 1981. Leaf senescence: correlated with increased levels of membrane permeability and lipid peroxidation, and decreased levels of superoxide dismutase and catalase. *J. Exp. Bot.*, 32(1): 93-101.
- Dracup, M. 1991. Increasing Salt Tolerance of Plants Through Cell Culture Requires Greater Understanding of Tolerance Mechanisms. *Funct. Plant Biol.*, 18(1): 1-15.
- Foyer, C.H. and G. Noctor. 2005. Oxidant and antioxidant signalling in plants: a re-evaluation of the concept of oxidative stress in a physiological context. *Plant Cell Environ.*, 28(8): 1056-1071.
- Gill, S.S. and N. Tuteja. 2010. Reactive oxygen species and antioxidant machinery in abiotic stress tolerance in crop plants. *PPB*, 48(12): 909-930.
- Gomez, J. A. Jimenez, E. Olmos and F. Sevilla. 2004. Location and effects of long-term NaCl stress on superoxide dismutase and ascorbate peroxidase isoenzymes of pea (*Pisum sativum* cv. Puget) chloroplasts. *J. Exp. Bot.*, 55(394): 119-130.
- Hafeez, I., B. Sadia, H. Sadaqat, R. I Kainth, M. Iqbal and I. Khan. 2012. Establishment of efficient *in vitro* culture protocol for wheat land races of Pakistan. *Afr. J. Biotechnol.*, 11(11): 2782-2790.
- Hannachi, S., S. Werbrouck, I. Bahrini, A. Abdelgadir, H.A. Siddiqui and M.C. Van Labeke. 2021. Obtaining salt stress-tolerant eggplant somaclonal variants from *in vitro* selection. *Plants*, 10(11): 2539.
- Hanson, A., C. Nelsen, A. Pedersen and E. Everson. 1979. Capacity for proline accumulation during water stress in barley and its implications for breeding for drought resistance. *Crop Sci.*, 19(4): 489-493.
- Hasanuzzaman, M., M.B. Bhuyan, F. Zulfiqar, A. Raza, S.M. Mohsin, J.A. Mahmud, M. Fujita and V. Fotopoulos 2020. Reactive oxygen species and antioxidant defense in plants under abiotic stress: Revisiting the crucial role of a universal defense regulator. *Antioxidants*, 9(8): 681.
- He, D., Y. Yang and K. Scott. 1989. The effect of macroelements in the induction of embryogenic callus from immature embryos of wheat (*Triticum aestivum* L.). *Plant Sci.*, 64(2): 251-258.
- Karadimova, M. and G. Djambova. 1993. Increased NaCl-tolerance in wheat (*Triticum aestivum* L. and *T. durum* Desf.) through *in vitro* selection. *In Vitro Cell Dev-Pl*, 29: 180-182.
- Kintzios, S.E., M. Triantafyllou and J. Drossopoulos. 1996. Effect of genotype and different growth regulator treatments on callus induction, proliferation and plant regeneration from mature wheat embryos. *Cereal Res. Comm.*, 147-153.
- Kundzewicz, Z.W., L.J. Mata, N.W. Arnell, P. Döll, P. Kabat, B. Jiménez, K.A. Miller, T. Oki, Z. Sen and I.A. 2007. Freshwater resources and their management. Climate Change 2007: Impacts, Adaptation and Vulnerability. Contribution of Working Group II to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change, M.L. Parry, O.F. Canziani, J.P. Palutikof, P.J. van der Linden and C.E. Hanson, Eds., Cambridge University Press, Cambridge, UK (PDF) *Freshwater Resources and their Management*. Available from: https://www.researchgate.net/publication/263557304_Freshwater_Resources_and_their_Management#fullTextFileContent [accessed Jun 15 2025].
- Maddock, S., V. Lancaster, R. Risiott and J. Franklin. 1983. Plant regeneration from cultured immature embryos and inflorescences of 25 cultivars of wheat (*Triticum aestivum*). *J. Exp. Bot.*, 34(7): 915-926.
- Manai, J., H. Gouia and F.J. Corpas. 2014. Redox and nitric oxide homeostasis are affected in tomato (*Solanum lycopersicum*) roots under salinity-induced oxidative stress. *J. Plant. Physiol.*, 171(12): 1028-1035.
- Meredith, C.P. 1978. Response of cultured tomato cells to aluminum. *Plant Sci. Lett.*, 12(1): 17-24.
- Mittova, V., M. Tal, M. Volokita and M. Guy. 2002. Salt stress induces up-regulation of an efficient chloroplast antioxidant system in the salt-tolerant wild tomato species *Lycopersicon pennellii* but not in the cultivated species. *Physiol. Plant.*, 115(3): 393-400.
- Murashige, T. and F. Skoog. 1962. A revised medium for rapid growth and bio assays with tobacco tissue cultures. *Physiol. Plant.*, 15(3): 473-497.

- Nakano, Y., and K. Asada. 1981. Hydrogen peroxide is scavenged by ascorbate-specific peroxidase in spinach chloroplasts. *Plant Cell Physiol.*, 22(5): 867-880.
- Nazar, R., S. Umar, N. Khan and O. Sareer. 2015. Salicylic acid supplementation improves photosynthesis and growth in mustard through changes in proline accumulation and ethylene formation under drought stress. *S. Afr. J. Bot.*, 98: 84-94.
- Oldeman, L., V. Van Engelen and J. Pulles. 1990. The extent of human-induced soil degradation. *Annex*, 5: 1992-1993.
- Özgen, M., M. Türet, S., Altınok and C. Sancak. 1998. Efficient callus induction and plant regeneration from mature embryo culture of winter wheat (*Triticum aestivum* L.) genotypes. *Plant Cell Rep.*, 18: 331-335.
- Patnaik, D., D. Vishnudasana and P. Khurana. 2006. Agrobacterium-mediated transformation of mature embryos of *Triticum aestivum* and *Triticum durum*. *Curr. Sci. India*, 307-317.
- Polash, M.A.S., M.A. Sakil and M.A. Hossain. 2019. Plants responses and their physiological and biochemical defense mechanisms against salinity: A review. *Trop. Plant Res*, 6: 250-274.
- Rao, M. 1992. Cellular detoxifying mechanisms determine the age dependent injury in tropical trees exposed to SO₂. *J. Plant Physiol.*, 140(6): 733-740.
- Sadhu, A., S. Bhadra and M. Bandyopadhyay. 2016. Novel nuclei isolation buffer for flow cytometric genome size estimation of Zingiberaceae: a comparison with common isolation buffers. *Ann. Bot. London*, 118(6): 1057-1070.
- Salama, E., A. Abido, A. Khaled and N. Abdelsalam. 2013. Embryo callus induction and regeneration of some Egyptian wheat cultivars. *Res J. Agri. Biol. Sci.*, 9(2): 96-103.
- Šramková, Z., E. Gregová and E. Šturdík. 2009. Chemical composition and nutritional quality of wheat grain. *Acta Chim Slov*, 2(1): 115-138.
- Sriniegn, K., T. Saisavoey and A. Karnchanatat. 2015. Effect of salinity stress on antioxidative enzyme activities in tomato cultured in vitro. *Pak. J. Bot.*, 47(1): 1-10.
- Stals, H. and D. Inzé. 2001. When plant cells decide to divide. *Trends Plant Sci.*, 6(8): 359-364.
- Suda, J., L.A. Meyerson, I.J. Leitch and P. Pyšek. 2015. The hidden side of plant invasions: the role of genome size. *New Phytol.*, 205(3): 994-1007.
- Tal, M. 1994. In vitro selection for salt tolerance in crop plants: theoretical and practical considerations. *In Vitro-Plant*, 30: 175-180.
- Tyankova, N. and N. Zagorska. 2001. Genetic control of in vitro response in wheat (*Triticum aestivum* L.). *In Vitro Cell Dev-Pl*, 37: 524-530.
- Vanlalruati, V., G. Kumar and A. Tiwari. 2019. Effect of saline stress on growth and biochemical indices of chrysanthemum (*Chrysanthemum morifolium*) germplasm. *Ind. J. Agr. Sci.*, 89(1): 41-45.
- Yu, Y., J. Wang, M.L. Zhu and Z.M. Wei. 2008. Optimization of mature embryo-based high frequency callus induction and plant regeneration from elite wheat cultivars grown in China. *Plant Breed.*, 127(3): 249-255.
- Zale, J.M., H. Borchardt-Wier, K.K. Kidwell and C.M. Steber. 2004. Callus induction and plant regeneration from mature embryos of a diverse set of wheat genotypes. *PCTOC*, 76: 277-281.