

NUTRITIVE VALUE OF FREE RANGELAND GRASSES OF NORTHERN GRASSLANDS OF PAKISTAN

JAVED IQBAL SULTAN¹, INAM-UR-RAHIM¹, MUHAMMAD YAQOOB²,
HAQ NAWAZ AND MANSOOR HAMEED³

¹Institute of Animal Nutrition and Feed Technology, University of Agriculture, Faisalabad-38040 Pakistan

²Department of Livestock Management, University of Agriculture, Faisalabad-38040 Pakistan

³Department of Botany, University of Agriculture, Faisalabad-38040, Pakistan

Corresponding author e-mail: dr_haq_nawaz@yahoo.com

Abstract

A study was conducted in the valley of Chaghcharzai in Bunair district lying in the north Trans-Himalayan moist zone occupying Malakand Division, North Western Frontier Province (NWFP), Pakistan to determine the nutritive value of locally available free rangeland grasses. The study area lies between 34.42 to 34.66 degree latitude and 72.62 to 72.78 degree longitude, having a humid subtropical to temperate environment. The annual precipitation varies from 600 to 1000 mm, mainly during summer and spring. Ten free rangeland grasses were identified and analyzed for DM, OM, ash, CP, NDF, ADF, hemi-cellulose, and lignin. The mean percentage values for dry matter (DM), organic matter (OM), ash, crude protein (CP), neutral detergent fiber (NDF), acid detergent fiber (ADF), hemi-cellulose and lignin at early bloom stage were 33.1 ± 0.69 , 30.6 ± 0.55 , 7.4 ± 0.42 , 7.8 ± 0.33 , 54.7 ± 2.08 , 24.7 ± 0.89 , 30.0 ± 2.11 and 3.9 ± 0.22 , respectively. The mean percentage values for DM, OM, ash, CP, NDF, ADF, hemi-cellulose and lignin at mature stage were 43.6 ± 1.03 , 41.4 ± 0.86 , 7.1 ± 0.42 , 5.5 ± 0.25 , 61.9 ± 1.44 , 29.4 ± 1.16 , 31.5 ± 2.14 and 4.7 ± 0.17 , respectively. In free grazing rangeland grasses the highest ($p < 0.05$) potential intake rate (PIR) was observed for *Heteropogon contortus* (53.80 ± 15.82 g/4 minute) and lowest for *Cymbopogon schoenanthus* (35.8 ± 12.16 g/4 minute). However, the highest ($p < 0.05$) relative preference (RP) was noted for *Dichanthium annulatum* ($81.15 \pm 0.61\%$) and lowest for *Cymbopogon schoenanthus* ($19.33 \pm 1.84\%$). The mean *In vitro* dry matter digestibility (IVDMD) and metabolizable energy (ME) of free rangeland grasses at early bloom stage were $55.3 \pm 1.86\%$ and 7.57 ± 0.25 MJ/kg DM, respectively, whereas, the mean IVDMD and ME at mature stage were $44.0 \pm 2.11\%$ and 5.86 ± 0.29 MJ/kg DM, respectively. The chemical and structural composition, IVDMD, RP and PIR values indicate that free grazing rangeland grasses be fed to livestock with some supplementation for different levels of production and types of livestock.

Introduction

Livestock grazing represents a system of land management in non-agricultural marginal areas, whereas, on rangeland livestock grazing represents the most suitable land use (Jones & Martin, 1984). Rangelands support 30 million herds of livestock, which contribute US \$ 400 million to Pakistan's annual export earnings (Anon., 2006). Past policies have often favored crops over livestock production, resulting in misuse of land having economically inefficient production potentials. Good pastures are being converted into croplands leaving increasingly poorer lands for livestock production (Pratt *et al.*, 1997), without thinking about the conservation of soil. There was no appreciation of the value of grasses and their ability to hold the soil against destructive erosion (Heath *et al.*, 1985). In the northern areas of Pakistan, livestock contributes nearly 55% to the gross provincial income by the agriculture sector. The mostly hilly terrain (73%) of the province has little land for crop agriculture; hence, dependence on livestock is relatively

high, particularly for rural subsistence (Anon., 1998). To match the maintenance requirements of livestock, there is a need of 13.5 and 110.3 million tons of crude protein (CP) and total digestible nutrients (TDN), respectively (Anon., 2006). However, present feed resources provide 40% CP and 75% TDN to the livestock (Younas & Yaqoob, 2005). The deficiency of nutrients leads to under nourishment, low productivity and predisposes the livestock to parasitism, epidemics and breeding problems. The improper utilization of rangelands has resulted in great changes in their ecosystem. The more palatable grass species are becoming extinct and are replaced by less palatable weeds (Humphreys, 1984). An indicator to range deterioration, in NWFP is the decline in range dependent sheep and goat population by 10.43% and 39.23%, respectively during 1986 and 1996 (Anon., 1996).

There is a handsome share of various grass species to the feeding regimens of animals during scarcity periods. For prolonged winter scarcity, the grasses are harvested from protected hillside rangelands and stored as hay. Grasses from fertile cropland sides and adjacent uneven areas are also cut several times during summer and are fed to livestock. Free rangeland grasses are, however, still the main way of procuring feed. Nutritive value of locally available free rangeland grass species have never been explored. Therefore, this study was conducted to evaluate the nutritive value of the free rangeland grasses of Northern grasslands of Pakistan.

Materials and Methods

Study area: The study was conducted in the valley of Chaghcharzai in Bunair district lying in the North Trans-Himalayan moist zone occupying Malakand Division, North Western Frontier Province (NWFP), Pakistan to determine the nutritive value of locally available free rangeland grasses. The study area lies between 34.42 to 34.66 degrees latitude and 72.62 to 72.78 degree longitude, having a humid subtropical to temperate environment. It extends from "Budhal" foothills (up to 800 meters altitude) in the south to "Burha Banr" in the north and northeast (more than 2000 meters altitude). Total surface area is approximately 209 square kilometer, with north and south parts divided by a mountain ridge known as "Sar Qala". The annual precipitation varies from 600 to 1000 mm, mainly during summer and spring. In general the irrigated agriculture in the area is confined to the narrow valley bottom and adjoining gentle slopes. The gentle slopes on mid hills are mainly used for rain fed (Barani) agriculture. The medium to steep hill slopes near the residences and gentle to medium hill slopes away from the residences are protected during wet summer season for forage harvest. The forage is harvested at a mature stage during early autumn and fed as hay to the wintering livestock. The steeper hill slopes near the residences are generally used for free grazing by settled livestock throughout the year. The steeper slopes away from residences are generally scrub/bush lands and natural forest mostly grazed by nomadic sheep and goat flocks. The lower elevation rangelands are grazed during winter season and by nomadic flocks, while tracking to and fro of alpine pastures graze the upper elevation rangelands.

Identification and sampling of grass species: A questionnaire was prepared and the farmers of various social groups at three elevations i.e., upper, middle and lower elevations were interviewed. They were asked about the names of grasses in local language (Pashtu), their season of use, location, elevations and aspects of their

availability, species of animal fed and trends in their frequency. Ninety farmers (30 x 3, at each elevation) were interviewed in 9 villages (3 x 3, villages at each elevation). Based on the information generated by the farmers through the questionnaire, the samples of different grass species were collected and their specimens were sent to Pakistan Forest Institute, Peshawar, Pakistan for botanical identification. Ten available rangeland grass species were selected for detailed study. Samples of grass species harvested at early bloom and mature stages of growth were chopped to 2 to 3 cm in length, dried in air and stored in polythene bags for further analysis.

Nutritional value of free rangeland grasses

Chemical and structural constituents: The air-dried samples were further dried in a forced draught oven at 60°C and were analyzed for dry matter (DM), organic matter (OM), ash and nitrogen (Anon., 1990). Grass samples were also analyzed for neutral detergent fiber, (NDF) acid detergent fiber (ADF) hemi-cellulose and lignin (Van Soest & Robertson 1990).

Palatability of free rangeland grasses: Four mature local sheep of 2.5 years age (average body weight 40 kg) were purchased from the local livestock market and were drenched for internal parasites. The experiments for palatability measurement were conducted at Civil Veterinary Dispensary, "Deewana Baba" in District Buner of Malakand Division. The sheep were adapted to the grass hay, trained in the experimental procedure by offering the test samples to them alone or in pairs daily. It took about five weeks to accustom them with the hay and to train them in the experimental procedure before any measurement was taken. During preliminary periods each animal was fed a diet 800 g/day. The diet consisted of 80% mixed grass hay and 20% concentrate mixture. The diet contained 10% crude protein (CP) and 8.37 MJ/kg metabolizable energy (ME). The mineral elements were also added in the diet to meet the sheep requirement (Anon., 1985). The potential intake rate (PIR) for different forages was determined through the procedure adopted by Rehman (1995). Each forage species was offered to sheep for a set of comparisons consisting of four consecutive periods of one minute duration each at ten minutes interval. Sheep were offered forages in suitable plastic containers ensuring that part of forage was left over after one minute of intake. Relative preference (RP) was also evaluated in 95 days through the procedure used by Rehman (1995). The preference ranking of forages within each group was determined by offering forages in pairs, initially with the forage having highest PIR and then with other forages, until all possible combination within a group was studied and similarly all forage groups were studied. Like PIR, RP test also consisted of a set of four consecutive period of one-minute duration each at ten minutes interval. There was one-hour gap before a new set of comparisons was started with a maximum of four sets in a day. The containers of both forages in a pair were reversed for each successive comparison to avoid left or right hand bias. The preference for particular forage was determined by the standard procedure developed by Bell (1959) for two choice tests, as the intake of forage expressed as a percentage of the combined intake of both test and standard forage.

$$\text{Relative Preference (RP) \%} = \frac{\text{Amount of test forage eaten}}{\text{Amount of test + standard forage eaten}} \times 100$$

In vitro dry matter digestibility (IVDMD) and metabolizable energy (ME) MJ/kg DM: For the IVDMD determination, oven dried ground samples were incubated at $37\pm1^{\circ}\text{C}$ for 48 hours between pH 6.7-7.0 in an all glass system using 45 ml of inoculums. The inoculums comprised of 36 ml of McDougal's artificial saliva and 9.0 ml of strained fresh rumen liquor from grass fed sheep. After incubation and centrifugation, the residue was then treated for 48 hours with pepsin in weak acid (pH 2.0). The final residue was composed of undigested plant cell wall and bacterial debris and yield values were supposed to be comparable to *in vivo* apparent digestibility (Tilley & Terry, 1963). The IVDMD was used for calculating the metabolizable energy (ME) of grass species by the following equation (Anon., 1984).

$$\text{ME (MJ/kg DM)} = 0.15 \text{ IVDMD\%}$$

$$\text{IVDMD\%} = 0.98 \text{ IVDMD\%} - 4.8$$

The average values for chemical composition, structural constituents, IVDMD and ME were integrated to develop a matrix of correlation (Steel *et al.*, 1997).

Results and Discussion

Identification of range grass species and their use pattern: The free grazing rangeland grasses included *Heteropogon contortus*, *Chrysopogon aucheri*, *Panicum antidotale*, *Dichanthium annulatum*, *Chrysopogon gryllus*, *Cymbopogon jwarancusa*, *Chrysopogon montanus*, *Themeda anathera*, *Aristida adscensionis* and *Cymbopogon schoenanthus*. The *Chrysopogon aucheri*, *Panicum antidotale*, *Cymbopogon jwarancusa*, *Cymbopogon schoenanthus* and *Aristida adscensionis* were commonly available at lower and middle elevation. The *Chrysopogon gryllus* was characteristically a high elevation grass. The *Dichanthium annulatum*, *Heteropogon contortus* and *Themeda anathera* were available at all elevations. Among the free grazing rangeland grasses found mainly on rocky slopes and shallow sites included *Heteropogon contortus*, *Chrysopogon aucheri*, *Chrysopogon gryllus*, *Chrysopogon montanus*, *Themeda anathera* and *Cymbopogon jwarancusa*. Whereas, the grass species found on less rocky and less shallow sites included *Panicum antidotale*, *Dichanthium annulatum* and *Cymbopogon schoenanthus*.

The results of present study were in line with the findings of Leede, (1998) who reported that in the subtropical zone of Leganai range (Bunair) of Malakand division, the free grazing rangelands grasses include *Cymbopogon jwarancusa*, *Heteropogon contortus* and *Chrysopogon aucheri* in Topdarra range of Bunair, *Chrysopogon*, *Themeda* and *Heteropogon* were the common grass species. It was observed that *Dichanthium annulatum* and *Cymbopogon schoenanthus* were the most common grasses covering field boundaries at low elevation and on deep soils having higher clay contents. On the other hand *Themeda anathera*, *Chrysopogon montanus*, *Chrysopogon aucheri* and *Heteropogon contortus* were commonly found on shallow soil hill slopes.

Chemical and structural constituents of grasses: The chemical constituents of free grazing rangeland grasses at early bloom and maturity are presented in Table 1. The DM contents in free grazing rangeland grasses at early bloom varied from 30.4% (*Themeda anathera*) to 36.4% (*Cymbopogon schoenanthus*) and the average was $33.1\pm0.69\%$. At mature stage, the DM contents in these grasses varied from 39.9% (*Chrysopogon gryllus*) to

47.5% (*Cymbopogon aucheri*) and the mean was $43.6 \pm 1.03\%$. The OM in free grazing rangeland grasses at early bloom stage ranged from 28.4% (*Panicum antidotale*) to 33.2% (*Cymbopogon schoenanthus*) and the mean was $30.6 \pm 0.55\%$. At maturity stage the lower and upper OM values in these grasses were 37.4% (*Chrysopogon gryllus*) to 48.0% (*Cymbopogon schoenanthus*) and the mean was $41.4 \pm 0.86\%$. At early bloom stage, the ash contents of free grazing rangeland grasses varied between 6.1% (*Themeda anathera*) to 8.8% (*Cymbopogon schoenanthus*) and the mean was $7.4 \pm 0.42\%$. The ash contents of free rangeland at mature stage varied from 5.1% (*Heteropogon contortus*) to 9.2% (*Aristida adscensionis*, *Cymbopogon schoenanthus*) and the mean value was $7.1 \pm 0.42\%$. The CP contents at early bloom stage in these grasses ranged from 6.4% (*Dichanthium annulatum*) to 9.8% (*Chrysopogon montanus*) and the mean was $7.8 \pm 0.33\%$. At mature stage the range for CP was 4.3% (*Dichanthium annulatum*) to 7.1% (*Aristida adscensionis*) and the mean was $5.5 \pm 0.25\%$. These results supported the findings of Distel *et al.*, (2005) who reported that CP contents in different grass species declined with time.

The structural constituents of free grazing rangeland grasses at early bloom and maturity are presented in Table 2. The NDF contents in free grazing rangeland grasses at early bloom varied from 43% (*Themeda anathera*) to 67% (*Cymbopogon schoenanthus*) and mean was $54.7 \pm 2.08\%$. At mature stage, the NDF ranged from 56% (*Themeda anathera*) to 70% (*Cymbopogon schoenanthus*), and mean $61.9 \pm 1.44\%$. The ADF contents at early bloom varied from 20% (*Themeda anathera*) to 31% (*Dichanthium annulatum*) and the mean was $24.7 \pm 0.89\%$. At mature stage, ADF varied from 22% (*Aristida adscensionis*) to 34% (*Dichanthium annulatum*) and the mean was $29.4 \pm 1.16\%$. The percentage of hemi-cellulose contents in the free grazing rangeland grasses at early bloom stage varied between 21% (*Chrysopogon aucheri*) to 43% (*Cymbopogon schoenanthus*) and the mean was $30.0 \pm 2.11\%$. At mature stage hemi-cellulose contents varied from 24% (*Chrysopogon aucheri*) to 43% (*Cymbopogon schoenanthus*) and the mean was $31.5 \pm 2.14\%$. The lignin contents in free grazing rangeland grasses at early bloom stage varied between 3.1% (*Themeda anathera*) to 5.2% (*Cymbopogon schoenanthus*) and the mean was $3.9 \pm 0.22\%$. At mature stage, lignin contents varied from 3.6% (*Chrysopogon gryllus*) to 5.6% (*Cymbopogon schoenanthus*) and the mean was $4.7 \pm 0.17\%$.

The structural constituents (NDF, ADF, hemi-cellulose and lignin) increased in grasses from early bloom to maturity stage. An increase in fiber and lignin contents over time had been reported in grasses by Distel *et al.*, (2005). According to Cherney *et al.*, (1993), the tropical grasses generally showed an increase in structural constituents with increasing maturity. Bourquin *et al.*, (1994) reported 72.4% NDF and 43.8% ADF in the orchard grass on DM basis. Sanderson *et al.*, (1989) observed a difference of 31.4 to 66.8% in NDF contents of alfalfa in two different years. In the stem of alfalfa grass the NDF concentration ranged from 21 to 68%. According to Cherney *et al.*, (1990) NDF and ADF tended to be lower in inflorescence than in other morphological components. The ADF was also reported higher in stem than in blade and sheath of leaves. The findings of this study were in line with those of Cherney *et al.*, (1993) who also reported an increase in all fiber constituents with increasing maturity. They reported that lignin was proportionally higher in stems than other parts of the plant. Brown *et al.*, (1984) reported that the soil fertility could also influence grass lignin concentration.

Palatability of rangeland grasses: Palatability refers to the relish with which feed is consumed as stimulated by the sensory impulses (Heath *et al.*, 1985). The potential intake rate (PIR) and relative preference (RP) are considered the main indicators for palatability (Rehman, 1995). The PIR and RP of ten free grazing rangeland grasses are presented in Table 3. The average PIR (grams consumed during 4 minutes per sheep) was the highest for *Heteropogon contortus* (53.80 ± 15.82 g/4 minute) followed by *Chrysopogon aucheri* (51.5 ± 12.5 g/4 minute), while the lowest PIR value was observed for *Cymbopogon schoenanthus* (35.8 ± 12.16 g/4 minute). The RP value was the highest for *Dichanthium annulatum* (81.15 ± 0.61 %) followed by *Panicum antidotale* (80.22 ± 1.41 %). The lowest RP value was observed for *Cymbopogon schoenanthus* (19.33 ± 1.84 %).

Among the free grazing rangeland grasses *Cymbopogon jwarancusa* despite of its higher PIR values showed relatively lower RP. This might be due to the presence of an essential oil (piperitone) that made it less palatable. This essential oil is responsible for the peculiar smell and taste of the grass (Rehman, 1995) and it might decrease the relative preference of the grass despite of its higher PIR. The PIR was more strongly affected by the degree of tenderness and stage of growth while the RP seemed to be more affected by the intrinsic chemical factors, hence when the grass was offered in pair with other forages having no such repellent essential oil, the other forage showed higher RP value despite of their lower PIR.

Correlation of palatability with chemical constituents, structural constituents and IVDMD is presented in Table 5. The RP and PIR were positively correlated ($r=0.86$) with each other across all grasses studied. These results supported the findings of Rehman (1995) who suggested PIR as a useful indicator of preference. The results of present study revealed that although the PIR and RP correlated with each other, their major determinant affects both the parameters differently. The PIR was influenced by the degree of tenderness, while RP was influenced by chemical factors. The present study suggests that more precise prediction equations can be developed if concentrations of sugars and chemical factors were integrated.

Kenney & Black (1984) reported that when effects of taste and odor were removed, sheep preferred diet having faster intake rates. The IVDMD is positively correlated with both RP and PIR across all the grasses ($r=0.31$, 0.04). These results supported the findings of Provenza *et al.*, (1996) who reported that lambs preferred food having higher CP contents. The CP contents had positive correlation with RP ($r=0.02$) but negative correlation with PIR ($r= -0.23$). The ADF showed positive correlation with RP ($r=0.33$) and PIR ($r=0.2$), whereas NDF, hemi-cellulose and lignin contents showed negative correlation with RP and PIR (Table 5). The results of present study were in agreement with Rehman (1995) who reported that NDF and lignin had negative correlation with both RP and PIR across all the species. The rapid accumulation of cell wall contents, fast lignification of cell wall and rapid reduction in CP levels may allow the unpalatable grass to avoid grazing since an early stage of re-growth. Fibrousness reduced intake rate because of the associated reduction in bite size to properly sever forage and the associated increase in chewing time necessary to adequately process the forage (Laca *et al.*, 2001). Lignin and CP contents interfered with the digestion of structural carbohydrates, the former by acting as a physical barrier to rumen microbial enzymes (Moore & Jung, 2001) and the later by limiting rumen microbial growth (Orskov, 1982). The factors affecting palatability need more investigation (Pratt *et al.*, 1997).

Table 3. Potential intake rate (PIR) and relative preference (RP) of free grazing rangeland grasses of Northern grasslands of Pakistan.

S. No	Grass name	PIR (g/4 minute) \pm SE*	(%) RP \pm SE**
1.	<i>Heteropogon contortus</i>	53.8 \pm 15.82 ^a	72.38 \pm 0.93 ^b
2.	<i>Chrysopogon aucheri</i>	51.5 \pm 12.50 ^a	55.03 \pm 0.84 ^c
3.	<i>Panicum antidotale</i>	48.8 \pm 16.92 ^{ab}	80.22 \pm 1.41 ^a
4.	<i>Dichanthium annulatum</i>	48.5 \pm 12.47 ^{ab}	81.15 \pm 0.61 ^a
5.	<i>Chrysopogon gryllus</i>	45.3 \pm 11.66 ^{ab}	71.20 \pm 0.68 ^b
6.	<i>Cymbopogon jwarancusa</i>	44.3 \pm 10.22 ^{ab}	28.50 \pm 0.53 ^e
7.	<i>Chrysopogon montanus</i>	42.5 \pm 11.35 ^{ab}	57.78 \pm 1.08 ^c
8.	<i>Themeda anathera</i>	41.8 \pm 8.42 ^{abc}	44.97 \pm 1.82 ^d
9.	<i>Aristida adscensionis</i>	40.5 \pm 10.69 ^{abc}	42.53 \pm 2.287 ^d
10.	<i>Cymbopogon schoenanthus</i>	35.8 \pm 12.16 ^{bc}	19.33 \pm 1.84 ^f

SE = Standard error

*Each figure represent mean (\pm standard error of the mean) of 4 samples.Figures having different letters are significant ($p<0.05$) within the same forage grass.**Each figure represent mean (\pm standard error of the mean) of 48 comparisons.Figures having different letters are significant ($p<0.05$) within the group.**Table 4. Digestibility and metabolizable energy of free grazing rangeland grasses of Northern grasslands of Pakistan.**

S. No.	Grass name	Early bloom		Maturity	
		IVDMD %	MJ/kg DM	IVDMD %	MJ/kg DM
1.	<i>Heteropogon contortus</i>	48.6	6.40	37.3	4.77
2.	<i>Chrysopogon aucheri</i>	58.5	7.87	43.5	5.69
3.	<i>Panicum antidotale</i>	49.4	6.53	38.9	4.98
4.	<i>Dichanthium annulatum</i>	51.4	6.82	39.5	5.11
5.	<i>Chrysopogon gryllus</i>	62.5	8.45	53.9	7.20
6.	<i>Cymbopogon jwarancusa</i>	50.4	6.70	35.0	4.44
7.	<i>Chrysopogon montanus</i>	57.2	7.70	48.5	6.40
8.	<i>Themeda anathera</i>	68.1	10.96	51.5	6.86
9.	<i>Aristida adscensionis</i>	58.3	7.87	52.6	7.03
10.	<i>Cymbopogon schoenanthus</i>	48.7	6.44	38.9	4.98
Mean \pm SE		55.3\pm1.86	7.57\pm0.25	44.0\pm2.11	5.86\pm0.29

Table 5. Correlation matrix among relative preference, potential intake rate, digestibility, crude protein and structural constituents of free grazing rangeland grasses of Northern grasslands of Pakistan.

.	PIR	IVDMD	CP	NDF	ADF	Hemi-cellulose	Lignin
RP %	+0.86	+0.31	+0.02	-0.48	+0.33	-0.70	-0.50
PIR	-	0.04	-0.23	-0.72	+0.20	-0.70	-0.66
IVDMD	-	-	+0.36	-0.08	-0.42	-0.39	-0.61

In vitro dry matter digestibility (IVDMD) and metabolizable energy (ME) of grasses: The IVDMD for free grazing rangeland grasses at early bloom stage varied from 48.6 (*Heteropogon contortus*) to 68.1% (*Themeda anathera*) and the mean was 55.3 \pm 1.86% (Table 4). The ME value at early bloom stage varied from 6.40 MJ/Kg DM (*Heteropogon contortus*) to 10.96 MJ/Kg DM (*Themeda anathera*) and mean was 7.57 \pm 0.25 MJ/Kg DM. The IVDMD value of free grazing rangeland grasses at maturity varied from 35.0% (*Cymbopogon jwarancusa*) to 53.9% (*Chrysopogon gryllus*) and the

mean was $44.0 \pm 2.11\%$ (Table 4). The derived ME value at maturity had a range of 6.70 MJ/Kg DM (*Cymbopogon jwarancusa*) to 8.45 MJ/Kg DM (*Chrysopogon gryllus*) with a mean of 5.86 ± 0.29 MJ/Kg DM. The IVDMD was positively correlated with CP ($r=0.36$) and negatively correlated with NDF ($r= -0.08$), ADF ($r= -0.42$), hemi-cellulose ($r= -0.39$) and lignin ($r= -0.61$). The IVDMD decreased with increasing maturity of the plants and similar findings were reported by Skerman & Riveros (1990) who found a fall of 0.1 to 0.2% DM digestibilities per day with maturity of pasture grasses. Buxton (1989) reported that the proportion of stem in a grass approaching maturity was the main morphological factor determining the digestibility. Gabrielsen *et al.*, (1990) and Van Soest (1965) reported that NDF, ADF and lignin concentration increased with maturity while IVDMD and CP declined. Revell *et al.*, (1994) reported a positive correlation between CP and digestibility, whereas, Cherney *et al.*, (1990) observed negative correlation of IVDMD with NDF, ADF and lignin. The results of our study were in line with those of Van Soest (1978) who reported poor relationship of NDF with digestibility. Lichtenberg & Hemken (1985) also reported that per unit increase in lignin often resulted in a three to four unit decrease in DM digestibility. It had been reported that cell wall component, NDF, ADF and lignin were negatively correlated with IVDMD in tree leaves (Mowatt *et al.*, 1969, Kundu & Sherma, 1988; Perveen, 1998).

References

Anonymous. 1984. *Energy Allowances and Feeding Systems for Ruminants*. Ministry of Agriculture Food and Fisheries, Her Majesty Stationery Office, London.

Anonymous. 1985. *Nutrient Requirements of Sheep*: Sixth Revised Edition. National Academy Press, Washington, DC.

Anonymous. 1990. *Official methods of analysis*. 13th Ed. Association of Official Analytical Chemists (AOAC), Washington, DC.

Anonymous. 1996. *Pakistan census of livestock*. Agriculture census organization, Statistics Division, Islamabad, Pakistan.

Anonymous. 1998. *Background paper on the sustainable management of grazing lands in the NWFP*. IUCN, The World Conservation Union, Sarhad Program Office Pakistan.

Anonymous. 2006. *Economic Survey, Government of Pakistan, Finance Division, Islamabad, Pakistan*.

Archer, S. and F.E. Smeins 1993. Ecosystem level processes In: *Grazing management: An ecological perspective*. (Eds.): R.K. Heitschmidt and J.W. Stuth. Timber Press, Portland Oregon, USA.

Bell, F.R. 1959. Preference thresholds for taste determination in goats. *J. Agric. Sci.*, 52: 125-29

Bourquin, L.D., E.C. Tibgemeyer, J.V. Milgen and J.C. Fahey Jr. 1994. Forage level and particle size effects on orchard grass digestion by steers: II Ruminal digestion kinetics of cell wall components. *J. Anim. Sci.*, 72: 759-67.

Brown, P.H., R.D. Graham and D.G.D. Nicholas. 1984. The effect of manganese and nitrate supply on the level of phenolics and lignin in young wheat plant. *Plant Soil.*, 81: 437-440.

Buxton, D.R. 1989. *In Vitro* digestion kinetics of temperate perennial forage legume and grass stems. *Crop Sci.*, 29: 213 -219.

Cherney, D.J.R., D.R. Mertens and J.E. Moore. 1990. Intake and digestibility by withers as influenced by forage morphology at three levels of forage offering. *J. Anim. Sci.*, 68(12): 4387-4399.

Cherney, D.J.R., J.H. Cherney and R.F. Lucey. 1993. *In Vitro* digestion kinetics and quality of perennial grasses as influenced by forage maturity *J. Dairy Sci.*, 76: 790-797.

Distel, R.A., N.G. Didoné and A.S. Moretto. 2005 Variations in chemical composition associated with tissue aging in palatable and unpalatable grasses native to central Argentina. *J. Arid Environ.*, 62: 351-357.

Gabrielsen, B.C., K.P. Vogel, B.E. Anderson and J.K. Ward. 1990. Alkali labile cell wall phenolics and forage quality in Switch grasses selected for different digestibility. *Crop Sci.*, 30(6): 1313-1320.

Heath, M.E., R.F. Barnes and D.S. Metcalfe. 1985. *Forages: The science of grassland agriculture*. Iowa State University Press. Ames, Iowa, USA.

Humphreys, L.R. 1984. *Tropical pastures and fodder crops*. Longman Group U.K. Ltd. Longman House, Burnt Mill, Harlow Essex CM 20 2J. E. England.

Jones, G.E. and Martin. 1994. *Eco zone suite of model, for FAO training service*. Eco zone Gough SAC Edinburgh Policy Analysis Division, Rome.

Kenney, P.A. and J.L. Black. 1984. Factors affecting diet selection by sheep I. Potential intake rate and acceptability of feed. *Aust. J. Agric. Res.*, 35: 55 1-63.

Kundu, S.S. and V. Sharma. 1988. Chemical composition and *in vitro* dry matter digestibility of certain tree leaves. *Indian J. Anim. Nutr.*, 5(1): 23-27.

Laca, E.A., L.A. Shipley and E.D. Reid. 2001 Structural anti-quality characteristics of range and pasture plants. *J. Range Manage.*, 54: 413-419.

Leede, B. 1998. *Vegetation production in hill-side plantation in Bunair and upper Swat. Environmental Rehabilitation Project*. Technical Report # 2.7 (WFP 4659).

Lichtenberg, V.L. and R.W. Hemken. 1985. Hay Quality. In: *Grazing management: An ecological perspective*. Timber Press, Portland, Oregon USA.

Moore, K.J. and H.G. Jung. 2001. Lignin and fiber digestion. *J. Range Management*, 54: 420-430.

Mowatt, D.N., M.L. Kwain and J.E. Wind. 1969. Lignification and *in vitro* cell wall digestibility of plant parts. *Can. J. Plant Sci.*, 49: 499-504.

Orskov, E.R. 1982. *Protein Nutrition in Ruminants*. Academic Press, London.

Perveen, S. 1998. *Nutritive evaluation of some fodder tree leaves through In vitro digestibility techniques*. Technical Paper, NWFP Agricultural University, Peshawar.

Pratt, D.J., F.L. Gall and C.D. Haan. 1997. *Investing in Pastoralism*. The International Bank of Reconstruction and Development. The World Bank 1818 H Street, NW Washington, D.C. 20433, USA.

Provenza, F.D., C.B. Scott, T.S. Phy and J.J. Lynch. 1996. Preference of sheep for foods varying in flavor and nutrients. *J. Anim. Sci.*, 74(10): 2355-61.

Rehman, A. 1995. *The potential for the use of salt bush in sheep grazing systems during summer and autumn in a Mediterranean climate*. Ph.D. Thesis, University of Western Australia.

Revell, D.K., S.K. Baker and B.B. Purser. 1994. Estimates of the intake and digestion of nitrogen by sheep grazing a Mediterranean pasture as it matures and senesces. *Proc. Aust. Soc. of Anim. Prod.*, 20: 217-20.

Sanderson, M.A., J.S. Hornstein and W.F. Wedin. 1989. Alfalfa morphological stage and its relation to *in-situ* digestibility of detergent fiber fractions of stem. *Crop Sci.*, 29:13 15-19.

Skerman, P.J and F. Riveros. 1990. *Tropical grasses*. FAO Plant Production and Protection Series, No. 23. FAO Rome

Steel, R.G.D., J.H. Torrie and J.H. Dickey. 1997. *Principles and Procedures of Statistics a Biometrical Approach*. 3rd Ed. McGraw Hill Book Co. Inc., New York, USA.

Tilley, J.M.A. and R.A. Terry. 1963. A two stage technique for *in vitro* digestion of forage crops. *Grass Forage Sci.*, 18: 104-111.

Van Soest, P.J. 1965. Voluntary intake in relation to intake and digestibility. Symposium on Factors influencing the voluntary intake by ruminants. *J. Anim. Sci.*, 24: 834-43.

Van Soest, P.J. 1978. Evaluation of forages and feedstuffs in the laboratory. *Paper presented at the 6th International Veterinary Congress, Buenos Aires, Argentina*.

Van Soest, P.J. and J.B. Robertson. 1990. Methods of dietary fiber, NDF and non starch poly saccharides. *Symposium paper ADSA June 1990. Cornell University, Dept. of Animal Science. Ithaca N.Y.*

Younas, M and M. Yaqoob. 2005. Feed resources of livestock in the Punjab, Pakistan. *Livest. Res. Rural Devel.* <http://www.cipav.org.co/lrrd/lrrd17/2/youn17018.htm>

(Received for publication 15 October 2007)