

SEED BORNE FUNGI ASSOCIATED WITH BOTTLE GOURD (*LAGENARIA SICERARIA* (MOL.) STANDL.)

NASREEN SULTANA AND A. GHAFFAR*

*Crop Diseases Research Institute, PARC,
Karachi University Campus, Karachi-75270, Pakistan*

**Department of Botany, University of Karachi, Karachi-75270, Pakistan.*

Abstract

Using ISTA techniques, the seed borne fungi of bottle gourd (*Lagenaria siceraria* was studied. A total of 22 genera and 45 species of fungi were isolated, of which 35 have not hitherto been recorded from seeds of bottle gourd in Pakistan. Both blotter and deep-freezing methods yielded quantitatively as well as qualitatively more fungi than agar plate method. *Lasiodiplodia theobromae*, *Fusarium semitectum* *Macrophomina phaseolina* and *Fusarium oxysporum* were most frequently isolated from 33, 91, 50 and 66 % seed samples of bottle gourd respectively.

Introduction

Bottle-gourd (*Lagenaria siceraria* (Mol.) Standl.), a paratropical species of Asian and African origin is cultivated throughout Pakistan all the year round for its young and tender fruits eaten as popular domestic vegetable called Lauki and Kaddu. The pulp has cooling and antibilious affect. The seed oil is applied externally in headache (Nazimuddin & Naqvi, 1984). There are reports of few diseases of bottle gourd eg., powdery mildew, downy mildew, fruit rot, anthracnose, root rot, root knot, insect pest and viral diseases (Kamal & Moghal, 1968; Maholay, 1989; Hafiz, 1986; Zitter *et al.*, 1996). Some of the fungi reported from seeds of bottle gourd are *Altenaria alternata*, *Aspergillus flavus*, *A. niger*, *Botryodiplodia theobromae*, *Chaetomium* sp., *Curvularia lunata*, *Drechslera tetramera*, *Fusarium equiseti*, *F. moniliforme*, *F. solani*, *Macrophomina phaseolina*, *Myrothecium roridum*, *Rizoctonia solani*, *Sclerotium rolfsii* and *Trichoderma* sp., (Manthachitra, 1971; Maholay & Sohi, 1976; Richardson, 1979; Maholay, 1989; Shakir & Mirza, 1992). The present study describes seed borne fungi of bottle gourd.

Materials and Methods

Using ISTA techniques (Anon., 1976), 10 bottle gourd seed samples collected from different places of Sindh, Baluchistan and Punjab were examined for the seed borne mycoflora. For standard blotter and deep freezing methods, seeds before and after treatment with 2 % NaOCl₂ for 2 minutes, were placed on three layers of moistened blotters, 10 seeds per Petri dish. The dishes were incubated at 24°C in 12 h alternating cycle of light and darkness for 7 days. In deep freezing method the treated and untreated seeds were incubated for 1 day each at 20°C and -20°C followed by 5 days incubation at 24°C. Fungi growing on seeds were identified after reference to Barnett & Hunter (1977), Booth (1971), Ellis (1971), Nelson *et al.*, (1983) and Raper & Fennell (1965).

Result and Discussion

Using blotter method, 22 genera and 45 species were isolated from 24 samples of bottle gourd seed collected from different parts of Pakistan followed by deep-freezing and agar plate methods where respectively 44 and 31 fungal species were detected (Table 1). Out of 45 fungal species, 35 fungal species viz., *Alternaria raphani* Groves & Skorko, *A. tenuissima* Kunze ex Pers., *Aspergillus candidus* Link, *A. quadrilineatus* Thom & Raper, *A. terreus* Thom, *A. wentii* Wehmer., *Chaetomium bostrychoides* Zope., *C. funicola* Cooke, *C. globosum* Kunze ex Fr., *C. olivaceum* Cook & Ellis, *C. tortile* Bainier, *Cladosporium cladosporioides* (Fr.) de Vries, *C. sphaerospermum* Penz., *Curvularia clavata* Jain, *C. lunata* (Wakker) Boedijn, *C. tuberculata* Jain, *Doratomyces stemonitis* (Pers. ex Fr.) Morton & Smith, *D. halodes* (Drechsler) Subram. & Jain, *D. hawaiiensis* (Bugn.) Subram. & Jain, *D. rostrata* (Drechsl) Richardson & Frasier, *D. state* of *Cochliobolus spicifer* Nelson, *Epicoccum purpurascens* Ehrenb. ex Schlechi, *Fusarium oxysporum* Schlecht emend. Snyd. & Hans., *F. semitectum* Berk & Rav., *Gliocladium roseum* Bainier, *Memnoniella echinata* (Riv.) Galloway, *Myrothecium verrucaria* (Alb. & Schw.) Ditm. ex Fr., *Nigrospora oryzae* (Berk & Br.) Petch, *Penicillium purpurogenum* Stoll, *Sordaria fimicola* (Rob.) Ces. & De Not., *S. tetraspore* Winter, *Stachybotrys atra* Corda, *Trichurus spiralis* Hasselring, *Ulocladium atrum* Preuss, and *U. botrytis* Preuss isolated do not appear to have been recorded from seeds of bottle gourd (Anon., 1990-2007; Noble & Richardson, 1968; Richardson, 1979; 1992; Wahid, 1985; Wahid *et al.*, 1991; Mathur, 1990; Ahmad, 1993). The average percent incidence and the range of occurrence of fungi in seed samples tested revealed that *Lasiodiplodia theobromae*, *Fusarium semitectum* *Macrophomina phaseolina* and *Fusarium oxysporum* were most frequent and isolated from 33, 91, 50 and 66 % seed samples of bottle gourd respectively. Most of the seeds infected by *L. theobromae* were dark coloured showed thick mycelial fragments on their surface. Seeds infected by *M. phaseolina* were discoloured with minute black sclerotia and most of them were of reduced size. *L. theobromae* and *M. phaseolina* have been reported to be seed borne on bottle gourd, squash and muskmelon causing black rot of fruit and blackening of seeds (Maholay & Sohi, 1982; Maholay, 1988, 1989).

Of the *Fusarium* spp., *F. semitectum* and *F. oxysporum* were frequently isolated followed by *F. solani* and *F. moniliforme*. *Fusarium* is a highly pathogenic fungus and its different species have been reported to cause seed rot, seedling blight and wilt in a number of cucurbitaceous crops (Booth, 1971). Similarly among *Drechslera* spp., *Drechslera* state of *Cochliobolus spicifer* and *D. halodes* showed seed rot and brown rot symptoms in seedlings. *Alternaria alternata* and *Curvularia lunata* caused delay or reduction in seed germination and is due to decay of seeds. Similar results have been reported by Mishra & Prakash (1975).

Use of 2% NaOCl₂ as seed disinfectant helped appreciably in minimizing the incidence of superficial and fast growing as well as common seed borne fungi like *Aspergillus* spp., *Chaetomium* spp., *Cladosporium* spp., *Rhizopus* spp., *Doratomyces stemonitis*, *Cephaliophora irregularis*, *Epicoccum purpurascens*, *Trichurus spiralis*, *Memnoniella echinata* and *Stachybotrys atra*. Whereas other seed borne fungi like *Curvularia* spp., *Drechslera* spp., *Fusarium* spp., *Myrothecium* spp., *Sodaria* spp., *Ulocladium* spp., and *Lasiodiplodia theobromae* were detected in greater frequency in seeds treated with 2% NaOCl₂. These observations are in conformity with the findings of Limonard (1968) and Khan *et al.*, (1988).

Both blotter and deep-freezing methods yielded quantitatively as well as qualitatively more fungi from seeds of bottle gourd. The standard blotter method yielded maximum number of fungi. Such similar results have been observed from the detection of seed borne fungi in rice (Khan *et al.*, 1988), cotton (Bhutta, 1988) and sunflower (Dawar, 1994). Begum & Momin (2000) reported that blotter method was found useful for detection of most infectious fungi of cucurbits. Deep-freezing method was found most suitable for detection of deep-seated as well as slow growing seed borne fungi like *Drechslera halodes*, *D. rostrata*, *Fusarium moniliforme*, *F. oxysporum*, *F. solani*, *Macrophomina phaseolina*, *Myrothecium* spp., *Penicillium purpurogenum* and *Sordaria* spp. These findings corroborate the reports that the deep-freezing method is more suitable for deeply seated seed borne fungi especially *Fusarium* spp., (Mathur *et al.*, 1975; Khan *et al.*, 1988; Diekmann & Assend, 1987; Dawar, 1994).

Disinfection of the seeds with 2% NaOCl₂ lowered the incidence of *Aspergillus* spp., *Cladosporium* spp., and *Rhizopus* sp., whereas these ubiquitous fungi were isolated in high percentage on agar plate method. Agar plate was found most suitable for isolation of *Curvularia* spp., and *Drechslera* state of *Cochliobolus spicifer* where disinfected seeds of bottle gourd were used. Khan *et al.*, (1988) preferred the use of agar plate method over the blotter method for isolation of *Curvularia* spp., and *Drechslera* spp., from disinfected seeds of rice. However, in the present studies *Drechslera rostrata* and *D. halodes* were isolated more frequently from bottle gourd seeds by deep-freezing method.

Presence of *Aspergillus* spp., especially *A. niger* and *A. flavus* on seeds of bottle gourd in higher frequencies and its association with ungerminated seeds of bottle gourd confirmed the findings that species of *Aspergillus* though occur as saprophytes may cause low germination in seeds (Christensen, 1967; Shakir & Mirza, 1992; Dawar, 1994). *Macrophomina phaseolina*, a virulent fungus causing charcoal rot and stem rot disease in a number of crops, was present in higher percentage in bottle gourd. *M. phaseolina* has been isolated from seeds of various cucurbitaceous crops and found associated with diseased seeds (Shakir & Mirza, 1992; Wahid, 1985; Shakir *et al.*, 1995; Maholay, 1988; 1989). *Lasiodiplodia theobromae* has been reported to cause seed borne disease in watermelon, squash and bottle gourd (Sohi & Maholay, 1974; Maholay & Sohi, 1976; Sultana & Ghaffar 1992). In the present study *Fusarium oxysporum* and *F. solani* were most consistently isolated fungi from bottle gourd seeds and varied in the nature and severity of the symptoms that they induced in emerging seeds and seedlings. *F. solani* has been reported on seeds of bottle gourd, sponge gourd (Shakir *et al.*, 1992; 1995; Wahid, 1985; Wahid *et al.*, 1991), *Benincasa cerifera* Sav., and *Lagenaria vulgaris* Ser., (Mirza & Qureshi, 1978). *F. oxysporum* has been isolated in high frequencies from seeds of water melon (McLaughlin & Martyn, 1982) and sponge gourd (Shakir *et al.*, 1995).

References

Ahmad, I., S. Iftikhar and A.R. Bhutta. 1993. *Seed-Borne Micro-organism in Pakistan*. Pakistan Agriculture Research Council, Islamabad. pp. 32.

Anonymous. 1990-2007. CAB Abstract, *Seed pathology and Microbiology*. Vol. 1-18, GISP,CABI Publishing ISSN 0959-9592.

Anonymous. 1976. International Rules of Seed Testing. *Proc. Int. Seed Test Assoc.*, 4: 3.49.

Barnett, H.I. and B.B. Hunter. 1972. *Illustrated Genera of Imperfecti Fungi*. Burgess Publishing Co., Minnesota, pp. 241.

Begum, H.A. and A. Momin. 2000. Comparison between two detection techniques of seed-borne pathogens in cucurbits in Bangladesh. *Pak J. Sci. & Inds. Res.*, 43: 244-248.

Bhutta, A.R. 1988. Comparison of cotton seed health testing method and their economics. *Pakistan Cotton*, 32: 146-153.

Booth, C. 1971. *The Genus Fusarium*. Commonwealth Mycological Inst, Kew, Surrey, England. 237 pp.

Charya, M.A.S. and S.M. Ready. 1979. Studies on seed mycoflora of *Cajanus cajan*. *Geobios*, 6: 299-301.

Christensen, C.M. 1967. Germinability of seeds free and invaded by storage fungi. *Proc. Assoc. of Seed Analyt. N. Am.*, 57: 141-143.

Dawar, S. 1994. *Studies on the seed-borne fungi associated with sunflower*. Ph.D. Thesis. Dept. Bot., Univ. Karachi, Pakistan, pp. 213.

Diekmann, M. and S. Assend. 1989. Comparison of agar and deep freezing blotter test for detection of *Fusarium* spp., in seed of lentil, chickpea and barley. *Zeitschrift fur pflanzenkrankheiten und pflanzenschutz*, 96:134.

Ellis, M.B. 1971. *Dematiaceous Hyphomycetes*. CMI, Kew, Surrey, England, pp. 608.

Hafiz, A. 1996. *Plant Disease*. Pakistan Agric. Res. Coun. Islamabad, 552 pp.

Kamal, M. and S.M. Moghal. 1968. *Studies on plant diseases of South West Pakistan*. Agric. College Tandojam, Pakistan, 207 pp.

Khan, S.A.J., A.K. Khanzada, N. Sultana and M. Aslam. 1988. Evaluation of seed health testing techniques for the assessment of seed borne mycoflora of rice. *Pak. J. Agric. Res.*, 9: 502-505.

Limonard, T. 1968. Ecological aspect of seed health testing. *Proc. Int. Seed Test. Assoc.*, 33: 343-513.

Manthachitra, P. 1971. Investigations on seed-borne fungi of some vegetable crops of Thailand. *Summaries of research projects (1967-1988)*. S.B. Mathur. 1990. *Danish Govt. Inst. Seed Path. Dev. Countries, Denmark*. 18 pp.

Mclaughlin, R.J. and R.D. Martyn. 1982. Identification and pathogenicity of *Fusarium* species isolated from surface disinfected watermelon seed. *J. Seed Tech.*, 7: 97-107.

Maholay, M.N. 1989. Seed borne diseases of cucurbits.III. Bottle gourd (*Lagenaria siceraria* (Mol.) Standl. *Seed and Farm*, 15: 30-31.

Maholay, M.N. 1988. Seed borne disease of cucurbits. Muskmelon (*Cucumis melo* L.). *Seed and Farm*, 14: 11-12.

Maholay, M.N. 1989. Seed borne diseases of cucurbits.III. Bottle gourd (*Lagenaria siceraria* (Md.) Standl. *Seed and Farm*, 15: 30-31.

Maholay, M.N. and H.S. Sohi.1976. Studies on *Botryodiplodia theobromae* rot of *Dolichos biflorus*. *Indian J. Mycol. Plant Pathol.*, 6: 126-129.

Maholay, M.N. and H.S. Sohi. 1982. *Botryodiplodia* seed rot of bottle gourd and squash. *Indian J. Mycol. Plant Pathol.*, 12: 32-36.

Mathur, S.B. 1990. *Summaries of Research Project 1967-1988*. Danish Govt. Inst. of Seed Path. for Dev. Countries, Denmark. 111 pp.

Mirza, J.H. and M.S.A. Qureshi. 1978. *Fungi of West Pakistan*. Dept. Plant Pathol., Uni. Agri., Faisalabad. 311 pp.

Mishra, B. and O. Prakash. 1975. *Alternaria* leaf spot of soybean from India. *Indian J. Mycol. & Pl. Pathol.*, 5: 95.

Nazimuddin, S. and S.S. Naqvi. 1984. *Flora of Pakistan. No.154, Cucurbitaceae*. Deptt. Bot. Univ. Karachi. 56 pp.

Noble, M. and M.J. Richardson. 1968. *An annotated list of seed-borne disease*. 191 pp. Commonwealth Mycological Institute, Kew, Surrey, England.

Raper, K.E. and D.I. Fennel. 1965. *The genus Aspergillus*. Williams and Wilkins Co. Baltimore, pp 686.

Richardson, M.J. 1979. *An Annotated List of Seed borne Diseases*. Supplement 1. Int. Seed Test. Assoc. Zurich, Switzerland. 320 pp.

Richardson, M.J. 1990. *An Annotated list of Seed borne Diseases*. Int. Seed Test. Assoc. Zurich, Switzerland. 320 pp.

Shakir, A.S. and J.H. Mirza. 1992. Seed-borne fungi of Bottle gourd from Faisalabad and their control. *Pak. J. Phytopathol.*, 4: 54-57

Shakir, A.S., J.H. Mirza, S.T. Sahi and F. Ahmad. 1995. Detection of seed-borne fungi associated with sponge gourd {*Luffa cylindrica* (L.) Roem.}, their location in different seed components and their control. *Pak. J. Phytopathol.*, 7: 140-144.

Sultana, N. and A. Ghaffar. 1992. Seed-borne fungi associated with pumpkin (*Cucurbita pepo* L.) *Proceeding of National Symposium Status of plant pathology in Pakistan*, 117-123. In: Proceedings of National Symposium held at the Department of Botany, University of Karachi, 3-5 December, 1991. (Eds.): Abdul Ghaffar & Saleem Shahzad.

Wahid, A. 1985. Seed-borne pathogens of vegetable crops grown in Pakistan. A Report submitted to the Danish Govt. Institute of Seed Pathology for Developing Countries. Copenhagen, Denmark. 36p.

Wahid, A., S. Ali and A.S. Shakir. 1991. Seed-borne mycoflora of sponge gourd in Punjab. *Pak. J. Agri. Res.*, 12: 151-152.

Zitter, D.L., D.L. Hopkins and C.E. Thomas. 1996. *Compendium of Cucurbit Diseases*. APS Press, St. Paul, MN. pp. 732.

(Received for publication 17 September 2008)