

**C₄ PHOTOSYNTHETIC CHARACTERISTICS
AND ANTIOXIDATIVE PROTECTION OF C₃ DESERT
SHRUB *HEDYSARUM SCOPARIUM* IN NORTHWEST CHINA**

**CHUN-MEI GONG^{*}, XIAO-WEI GAO^{*}, DONG-LIANG CHENG^{*}
AND GEN-XUAN WANG^{**}**

*Key Laboratory of Arid and Grassland Ecology at Lanzhou University,
Ministry of Education, Lanzhou 730000, China and ^{**}College of Life Science, Zhejiang
University, Hangzhou 310027, China*

Abstract

To understand the adaptive mechanisms, and the strategy and direction of evolution of exploiting plants for living in oasis-desert transitional areas where there was a water gradient, mechanisms of C₄ photosynthesis and antioxidative protection were studied in leaves and assimilating shoots of C₃ desert shrub *Hedysarum scoparium* under water gradient accompanied with high irradiance and high temperature in simulated water gradient trial site that lies in the desert. In this study, C₄ pathway was detected in the leaves and assimilating shoots of *H. scoparium* and the antioxidative enzymatic system was found to be very important in the protection against oxidants, both increasing considerably antioxidative enzymes and C₄ photosynthetic enzymes under increasing water stress in C₃ plant. The results indicated that the C₄ photosynthetic response of assimilating shoots was more positive than that of leaves with water condition exacerbate. It was proposed that the direction of evolution of organs of C₄ photosynthesis and positive interaction between antioxidative enzymes and C₄ photosynthetic key enzymes in C₃ desert species were important factors for plant growth, survival and reproduction in water gradient environment.

Introduction

Hedysarum scoparium (Leguminosae), a desert pioneer plant, is famous for its capacity to resist aridity and fixate dune. It is a small-leaf desert shrub, distributing very widely in the arid regions of China with dry air, excess sunshine, and annual precipitation less than 130 mm (Li *et al.*, 1995). *H. scoparium* makes use of leaves and assimilating shoots performing photosynthesis as different photosynthetic organs. Some reports have been shown on the photosynthetic attribution in leaves of *H. scoparium* (Chen *et al.*, 1961; Li *et al.*, 1995), but less is known about its leaf-axis-like assimilating shoots. Su *et al.*, (2003) confirmed *H. scoparium* was C₃ plant by its characteristics of $\delta^{13}\text{C}$ value (-26.38 ‰) and gas exchanges (high CO₂ compensation point, 90 $\mu\text{mol} \cdot \text{mol}^{-1}$), but studies on photosynthetic enzymes were absent. In general, C₄ plants originated from tropical arid areas, which possess higher photosynthetic rates, lower photorespiration and better capacity to endure leanness than that of C₃ plants. The desert plants are often exposed to different stressful conditions such as high irradiance, extreme temperature, water deficit and air-dryness. Especially, water deficit is perhaps one principal factor of

Author for Correspondence:
Tel. / Fax: +86-931-891-2893;
E-mail: gcm228@163.com (C.M. Gong), wanggx@lzu.edu.cn (G.X. Wang)

impacts on plants growth and development, and causes physiological and biochemical responses. On the one hand, cellular malfunction caused by active oxygen species and lipid peroxidation was mitigated by antioxidative enzymes such as SOD (EC 1.15.1.1), POD (EC 1.11.1.7), CAT (EC 1.11.1.6). Antioxidant enzymes are known to increase in response to drought (Menconi *et al.*, 1995; Zhang & Kirkham, 1996; Sairam & Saxena, 2000; Sairam & Srivastava, 2001), high temperature (Upadhyaya *et al.*, 1990; Sairam *et al.*, 2000), external calcium under water stressed conditions (Li *et al.*, 2003). On the other hand, the reproductive parts of wheat (C_3 plant), chickpea (*Cicer arietinum*) and rapeseed (*Brassica campestris*) contain high activities of C_4 enzymes (Singal *et al.*, 1987; Singh, 1993), such as PEPCase and NADP-ME in the pods, which were purified and characterized by Das *et al.*, (1986) and Singal & Singh (1986) respectively. Voznesenskaya *et al.*, (2001) showed C_4 photosynthesis could function within a plant that its carbon isotope composition was like that of C_4 , but lacked Kranz anatomy. Hibberd & Quick (2002) also reported that tobacco and celery (a typical C_3 plant) showed characteristics of C_4 photosynthesis in cells of stems and petioles surrounding the xylem and phloem. Moreover, the enzymatic activities of C_4 pathway in soybean have positive relations to net photosynthetic rate (Li *et al.*, 2000). These results implied that some C_3 species might extend their photosynthetic types in different assimilating organs without Kranz anatomy along the increasing environmental stress.

Fryer *et al.*, (1998) studied the relationships between CO_2 assimilation, photosynthetic electron transport and antioxidant enzyme activities in field-grown maize and found that the donation of electrons to oxygen by the photosynthetic electron transport chain was elevated by growth at low temperatures (Fryer *et al.*, 1998). There was a positive correlation between the photosystem stoichiometries and active oxygen protective mechanisms (Yamazaki & Kamimura, 2001). Further research on the correlation between adaptable changes in photosynthetic pathway and antioxidative protection in different photosynthetic organs of desert plant is essential to understand the adaptive mechanisms and the strategy and direction of evolution of exploiting plants for living in oasis-desert transitional areas since the adaptation of desert vegetation across oasis-desert transitional areas was important for the oasis to maintain its stability. Water availability was the major limiting environmental factor for the growth of desert plants within desert ecosystem (Mittler *et al.*, 2001). So, the experiment is conducted in a simulated trial site in the desert area with a water gradient, attempting to investigate the following hypotheses: (1) C_3 dominant desert species *Hedysarum scoparium* extends its photosynthetic pathway to improve its survival and adaptation; (2) The positive interactions between antioxidative enzymes and C_4 photosynthetic key enzymes in C_3 desert species are important factors for the growth, survival and reproduction; (3) Assimilating shoots become an adaptable strategy and direction of evolution with water condition exacerbate in many of these species.

Materials and Methods

Habitat of trial site: The study was carried out near the town of Minqin, northwest of China, which is located in a joint region of Tengger Desert and Badain Jaran Desert ($38^{\circ}38'N$, $103^{\circ}05'E$). The trial site lies in the Plant Transpiration Consume Water Observation Station of the Desert Plantation of Gansu Provincial Institute of Desert Control Research. There are many sand dunes in a protected area in nature, correlating to *Nitraria tangutorum* and *Reaumuria soongorica* communities.

The average annual precipitation is 115 mm with the largest precipitation in July, August and September. The average annual evaporation is 2643.9 mm, 23 times of the average annual precipitation. The annual average temperature is 7.8°C with the highest temperature of 38.1°C and the lowest temperature of – 28.8°C. There are 165 days of non-frost days, 139 days of blowing sand and 37 days of sandstorm. The soil is sandy soil with the depth of ground water of 18 m (Zhao *et al.*, 2003a).

The growth conditions and plant materials procedures: *Hedysarum scoparium* was growing in the Plant Transpiration Consume Water Observation Station which consisted of 66 cultivated pools filling with sandy soil, establishing in a natural environment for plant growth. Cultivated pools were 1 m in both length and width with different depth of 1.6 m, 2.6 m and 3.6 m. The cobble and crushed stones were 11 cm in depth acting as a layer for water storage, where there was dune soil as the nutrient soil layer. The water that plants need was supplied by the groundwater eternally through the compensatory apparatus, which could stabilize water storage level. The Plant Transpiration Consume Water Observation Station has 3 depths of ground water (DGW), i.e., 1.4 m, 2.4 m and 3.4 m (Guo & Zhao, 1988). *H. ammodendron* and *H. scoparium* of two years old were transplanted to cultivated pools in 1995 (Zhao *et al.*, 2003a). Every water gradient and every plant have 4 repeats. Because the level of water storage of transpiration and the permeation instrument (were made up of cultivated pools and groundwater eternal compensatory apparatus) were stable, soil water contents of every layer of cultivated pools were almost constant, which created a more stable soil water environment for plants growth (Zhao *et al.*, 2003b).

The samples were all collected at growing and developing stage in August 2003. The collected leaves and assimilating shoots were frozen in liquid nitrogen immediately after sampling.

Determination of PFD and T_{Air}: We measured the photon flux density (PFD) and air temperature (T_{Air}) with portable photosynthetic gas analysis systems (CIRAS-1, PP Systems Corporation in Britain) before sampling on sunny and clear days.

Determination of soil water content: Water contents of the soil were determined by the fresh and drying weight of the soil. Soil samples were randomly collected from the ground every 20 cm in depth. This procedure was repeated 3 times.

The relative water content (RWC) of leaves and assimilating shoots were estimated by the fresh mass (FM) and saturated mass (SM). 0.5 g fresh leaf samples first were kept in water for 24 h and then dried in hot air oven till constant dry mass (DM) was achieved according to Barrs & Weatherley (1962).

$$\text{RWC}(\%) = ((\text{FM}-\text{DM})/(\text{SM}-\text{DM})) \times 100\%$$

Antioxidative enzymatic assays: Leaf tissue of 0.5 g was frozen in liquid nitrogen and ground to fine powder with a pre-cold mortar and pestle. The powder was homogenized in 1: 5 (w/v) 50 mmol L⁻¹ phosphate buffer (pH 7.8) containing 1 % PVP (MW 40,000) and 10 mmol L⁻¹ sulfhydryl alcohol. The homogenate was centrifuged at 15,000 g for 20 min and the supernatant obtained was used immediately for assay as enzyme extract. All steps in the preparation of the enzyme extract were carried out at 4°C. Determination of all enzyme activities was conducted through UV-751 ultraviolet spectrophotometer. An aliquot of the extract was used to determine the protein content according to Bradford (1976) using bovine serum albumin as a standard.

SOD (EC 1.15.1.1) activity was measured by its ability to inhibit the photochemical reduction of nitro blue tetrazolium (NBT) using the method of Beauchamp & Fridovich (1971) as modified by Dhindsa & Matowe (1981). The reaction mixture (3 mL) contained 50 mmol L⁻¹ phosphate buffer (pH 7.8) 0.1 mmol L⁻¹ EDTA, 13 mmol L⁻¹ methionine, 75 µmol L⁻¹ nitroblue tetrazolium (NBT), 2 µmol L⁻¹ riboflavin and 50 µL of the supernatant which had been diluted 5 times. The production of blue formazan was followed by monitoring the increase of the absorbance at 560 nm. A non-irradiated reaction mixture did not develop color and served as control.

POD (EC1.11.1.7) activity was assayed as the increase in absorbance due to the formation of tetra-guaiacol recorded at 470 nm (Castillo *et al.*, 1984). The 3 mL reaction mixture contained 16 mmol L⁻¹ guaiacol, 2 mmol L⁻¹ H₂O₂, 70 mmol L⁻¹ phosphate buffer (pH 6.1) and 20 µL enzyme extract which had been diluted twice.

CAT (EC 1.11.1.6) was assayed by measuring the initial rate of disappearance of hydrogen peroxide (H₂O₂) at 240 nm according to the method of Chance & Maehly (1955) as modified by Dhindsa & Matowe (1981) and Cakmak & Marschner (1992). The 3 mL reaction mixture contained 50 mmol L⁻¹ phosphate buffer (pH 7.0) 15 mmol L⁻¹ H₂O₂, and 50 µL enzyme extract which had been diluted 10 times. Enzyme solution containing hydrogen peroxide-free phosphate buffer was used as control.

Photosynthetic enzymatic extraction and assay: Enzymes were extracted from 0.1 g tissue which was ground at 4°C in a pre-cold mortar and pestle with 1 mL of grinding buffer containing 50 mmol L⁻¹ Tris-HCl (pH 7.0), 20 mmol L⁻¹ MgCl₂, 2 mmol L⁻¹ MnCl₂, 1 mmol L⁻¹ EDTA, 5 mmol L⁻¹ DTT, 10% glycerol and 1% PVP (MW 40,000). The crude extract was immediately centrifuged at 15,000×g for 15 min at 4°C and then used to assay different enzymatic activities at 25°C.

PEPCase, NADP-ME and NAD-ME activities were monitored spectrophotometrically by the change of pyridine nucleotide at 340 nm. PEPCase (EC 4.1.1.31) activity was assayed as described by Ting & Osmond (1973). The buffer for PEPC (1 mL) contained 50 mmol L⁻¹ Tris-HCl (pH 8.0), 5 mmol L⁻¹ MgCl₂, 2 mmol L⁻¹ DTT, 1 mmol L⁻¹ NaHCO₃, 0.2 mmol L⁻¹ NADH, 0.1 mmol L⁻¹ EDTA, 3 units malate dehydrogenase and 20 µL extract. The reaction was initiated with 2 mmol L⁻¹ PEP.

NADP-ME (EC 1.1.1.40) and NAD-ME (EC 1.1.1.38) activities were assayed as described by Sayre *et al.*, (1979). Assays of NADP-ME activity were carried out in 1 mL buffer containing 50 mmol L⁻¹ Tris-HCl (pH 8.0), 1 mmol L⁻¹ MgCl₂, 1 mmol L⁻¹ MnCl₂, 1 mmol L⁻¹ EDTA, 0.33 mmol L⁻¹ NADP, 5 mmol L⁻¹ L-malate and 20 µL extract. NAD-ME activity were assayed in 1 mL buffer containing 25 mmol L⁻¹ Tris-HCl (pH 7.4), 5 mmol L⁻¹ MnCl₂, 5 mmol L⁻¹ L-malate, 2 mmol L⁻¹ NAD, 7.5 mmol L⁻¹ MgSO₄, 5 mmol L⁻¹ DTT, 0.2 mmol L⁻¹ EDTA and 20 µL extract.

Protein content of the crude enzyme extract was detected according to Bradford (1976) using bovine serum albumin as a standard.

Results

Influence of PFD, T_{Air}, soil moisture and RWC to desert plants: The diurnal variations of PFD and T_{Air} were shown on sunny days in the early ten days of August 2003 in Minqin (Fig. 1). PFD reached its highest value (1,638 µmol m⁻² s⁻¹) at 13:00 and the highest T_{Air} is 42.4°C at 15:50. While the values of PFD and T_{Air} were all much higher at 15:00, they reached 1,568 µmol m⁻² s⁻¹ and 40°C respectively.




Fig. 1. Diurnal changes of photon flux density PFD ($\mu\text{mol m}^{-2} \text{s}^{-1}$) and air temperature T_{Air} ($^{\circ}\text{C}$) on sunny days in the early ten days of August 2003 in Minqin, northwest China




Fig. 2. Soil water moisture at different soil depths for three depths of ground water (DGW) which were 1.4 m, 2.4 m, and 3.4 m, respectively. Each value represents mean (\pm S.E.) of three replications ($p < 0.05$)

Soil moisture gradually increased with the increase of soil depths for the 3 depths of ground water, resulting in different effects on plant growth and development at different DGW (Fig. 2).

As shown in Table 1, RWC in leaves and assimilating shoots of *H. scoparium* were very low. Slightly higher RWC was recorded in the 1.4 m DGW with the lowest in the 3.4 m at 15:00. RWC was lower in the leaves than in the assimilating shoots and the changes of RWC in leaves were gentler under the same condition.

Table 1. Effects of different depths of ground water (DGW) on the relative water content (RWC, %) in leaves and assimilating shoots of *Hedysarum scoparium* at 15:00. Each value represents mean (\pm S.E.) of three replications.

Organs	DGW		
	1.4 m	2.4 m	3.4 m
Assimilating shoots	50.0 \pm 3.85	45.78 \pm 2.89	42.43 \pm 3.09
Leaves	47.58 \pm 2.05	42.62 \pm 2.10	38.87 \pm 2.39

Antioxidant enzymes system of *H. scoparium*: In order to compare the activities of antioxidant enzymes in leaves and in assimilating shoots, the changes in activities of antioxidative enzymes were investigated at different DGW at 15:00 when plants were exposed to high irradiance and temperature. SOD activity in leaves and assimilating shoots gradually increased with the increase of DGW. Among the enzymes responsible for the scavenging of active oxygen species, SOD activity was obviously much higher in leaves than in assimilating shoots of *H. scoparium* at 3 DGW (Fig. 3A), and the ratio of SOD activity of leaves/assimilating shoots gradually decreased.

Some different trend in POD activity in leaves of *H. scoparium* was observed (Fig. 3B). POD activity in leaves increased slightly from 1.4 m to 2.4 m but declined at 3.4 m DGW. POD activity in assimilating shoots gradually increased with the increase of DGW, as well as the ratio of POD activity of leaves / assimilating shoots gradually decreased with the increase of DGW.

CAT activity was found to be higher in leaves than in assimilating shoots of *H. scoparium* at 3 DGW (Fig. 3C). However, increased CAT activity was observed either in leaves or in assimilating shoots from 1.4 m to 2.4 m but declined at 3.4 m DGW. CAT activity increased subjected to mild stress but declined during severe water stress.

Activities of C₄ photosynthetic key enzymes: The activities of C₄ enzymes in *H. scoparium* were lower than that in C₄ plant. The activities of C₄ photosynthetic key enzymes were investigated in *Hedysarum scoparium* and *Haloxylon ammodendron* (a C₄ desert shrub) at 2.4 m DGW where represents moderate water stress (Table 2).

The activities of C₄ photosynthetic key enzymes in leaves were all higher than that in assimilating shoots of *H. scoparium* at different DGW. The activity of PEPCase in leaves reached the top at 15:00 (ca. 7 μ mol PEP min⁻¹mg⁻¹ protein) while it kept at a lower level in assimilating shoots (ca. 3 μ mol PEP min⁻¹mg⁻¹ protein). The activity of PEPCase in leaves increased slightly with the increase of DGW, and the activity of PEPCase in assimilating shoots increased in the same time, even the ratio of PEPCase activity of leaves / assimilating shoots was more and more lower (Fig. 4A), which was consistent with the changes of RWC. NADP-ME were also detected in leaves and in assimilating shoots of *H. scoparium*, NADP-ME activity in leaves gradually declined, but the changes of NADP-ME activity in assimilating shoots was similar to the changes of PEPCase. The activity of NAD-ME in leaves and assimilating shoots increased gradually with the increase of DGW, and the activity of NAD-ME was about 3 times higher in leaves than in assimilating shoots (Fig. 4B, C). The activity of NADP-ME was lower than that of NAD-ME.




Fig. 3. Variations of SOD, POD and CAT activities in leaves and assimilating shoots of *Hedysarum scoparium* at different DGW of 1.4 m, 2.4 m, and 3.4 m at 15:00. Each value represents mean (\pm S.E.) of three replications

Table 2. Comparisons of PEPCase, NADP-ME and NAD-ME activities in *Hedysarum scoparium* and *Haloxylon ammodendron* at 2.4 m DGW where water stress was moderate. Each value represents mean (\pm S.E.) of three replications.

Species	Organs	PEPCase ($\mu\text{mol PEP min}^{-1} \text{mg}^{-1}$ protein)	NADP-ME ($\mu\text{mol min}^{-1} \text{mg}^{-1}$ protein)	NAD-ME ($\mu\text{mol min}^{-1} \text{mg}^{-1}$ protein)
<i>Hedysarum scoparium</i> (C ₃)	Assimilating shoots	3.40 \pm 0.35	0.75 \pm 0.06	1.38 \pm 0.21
	Leaves	6.98 \pm 0.55	1.24 \pm 0.06	3.38 \pm 0.53
<i>Haloxylon ammodendron</i> (C ₄)	Assimilating shoots	53.39 \pm 1.91	24.21 \pm 1.52	7.37 \pm 0.70

Discussion

Antioxidant protection in *Hedysarum scoparium*: Protection might be performed by the increased activities of enzymes involved in the destruction of free radicals and oxidants, but their responses depended on the water status of the plant. RWC either in leaves or in assimilating shoots of *H. scoparium* was very low, which corresponded with the water use efficiency (WUE) of *H. scoparium* (Su 2003). Lower RWC of *H. scoparium* suggested that water availability was the key environmental factor. Lower RWC and gentler change were found in leaves than in assimilating shoots of *H. scoparium* under the same condition, indicating that assimilating shoots played an important role in water storage.

On exposure to high light, there was an increase in synthesis of protective antioxidant enzymes (Mishra *et al.*, 1993; Sharma & Singhal, 1992). They also observed that water stress altered the membrane organization and the absorbance/fluorescence properties of chloroplasts (Singh & Singhal, 1984). Among the enzymes responsible for the scavenging of active oxygen species, the changes of antioxidant enzymes were all undulated violently either in leaves or assimilating shoots in process of a day (data not shown). Activities of antioxidative enzymes were higher at noon and lower in the early morning and evening. In general, the activities of antioxidative enzymes were higher in leaves than in assimilating shoots. SOD activity was obviously much higher in leaves than in assimilating shoots of *H. scoparium* at 3 DGW (Fig. 3A), and SOD activity increased gradually both in leaves and in assimilating shoots, depending upon the increase of water stress. The ratio of SOD activity of leaves / assimilating shoots gradually decreased with the increase of DGW, demonstrating that SOD activity in assimilating shoots played a more significant role than that in leaves when plants grow in the prolonged water stress and explaining why the appearance of assimilating shoots of *H. scoparium* was essential to defend oxidative stress in exacerbated water environment. Lyshede (1979) pointed out the formation of assimilating shoots was the summit of evolution under drought condition. Exogenous antioxidants stimulated shoot organogenesis (Gupta & Datta, 2003). Maybe the evolution of shoot was supplemented and complemented each other with the increasing stress.

As SOD activity increased, the quantity of H₂O₂ formed through SOD catalysis might also increase. The action modes of these two enzymes were essentially different. CAT catalyzed the dismutation of two molecules of H₂O₂ to water and molecular O₂ at near diffusion-controlled rates, whereas POD used the substrates to reduce H₂O₂ to water. APx had a much higher affinity for H₂O₂ than CAT, but CAT had a much higher V_{max} (Asada, 1992). There was a similar effect in POD activity of *H. scoparium* with in SOD activity of *H. scoparium*. Induction of antioxidant enzymes, such as SOD, POD and CAT, could lead to the increase in antioxidant protection and the decrease in oxidative




Fig. 4. Variations of PEPCase, NADP-ME and NAD-ME activities in ■ leaves and □ assimilating shoots of *Hedysarum scoparium* at different DGW of 1.4 m, 2.4 m, and 3.4 m at 15:00. Each value represents mean (\pm S.E.) of three replications

damage. High activity of antioxidative enzymes could be correlated to the process of differentiation that occurred during shoot induction (Thakar & Bhargava, 1999). At the more drought-tolerant 3.4 m DGW, the lower POD activity in leaves and the higher in assimilating shoots were found, suggesting that POD might not greatly contribute to drought tolerance mechanism in leaves at 3.4 m DGW. In scavenging H₂O₂, maybe APX played a more important role in leaves (Willekens *et al.*, 1995). Therefore, further studies on other antioxidative enzymes are needed.

CAT was the enzyme capable of detoxifying H₂O₂ in the cell (Lin & Kao, 2000). We considered the performance of some enzyme activities from the perspective of H₂O₂ scavenging. So, we measured CAT activity. There was only minor losses of CAT activity occurred in assimilating shoots of *H. scoparium* at 3 DGW, but a marked apparent water deficit inactivation of CAT was observed in leaves of *H. scoparium* in 3.4 m DGW. CAT activity was suppressed in severe water stress, which was an indication of a reduced capacity of the leaves to decompose H₂O₂. This result was consistent with CAT activity in stems of *R. raetam* and in leaves of *A. halimus* in natural daylight (Streb *et al.*, 1997). The present findings further confirmed that CAT activity was subjected not only to high irradiance but also to water deficit. Furthermore, mild water stress could stimulate CAT activity and severe water stress inactivated CAT activity, supporting that drought stress inactivated the key enzyme activity of light respiration-ethanol acid oxidative enzyme 44% (Morgan *et al.*, 1994). The inactivation of the key enzyme activity resulted in the accumulation of more NADPH and ATP, and the increase of production of O₂⁻, H₂O₂ in chloroplast. More NADPH and ATP may supply more energy to recapture CO₂ released by respiration performing C₄ photosynthesis (Ku *et al.*, 1991; Li *et al.*, 2000) in C₃ plant. The result was also supported by the changes of C₄ enzymatic activities (Fig. 4). So, it was concluded that PSII might avoid photoinhibition by supplying more energy to ensure performance of C₄ pathway at high temperature and irradiance under severe water stress.

C₄ photosynthesis in *Hedysarum scoparium*: The variations in C₄ photosynthetic key enzymes were consistent with the changes of protein content of enzymes (data not shown), indicating that the increase of enzymatic activities was the synthesis of enzymatic protein but not the activation of enzymes.

The intrinsic link between photosynthesis and biomass production suggested that photosynthetic response to drought probably played a major role in determining the ability of these species to persist in drought areas (Lambers *et al.*, 1998; Gulás *et al.*, 2002), also in consideration of global change. Water availability was an important factor affecting photosynthetic activity of plant species (Llorens *et al.*, 2003). The lack of water that occurred during the dry season was the important limiting environmental factor that triggered the plant to change their photosynthetic strategy. Liu *et al.* (2004) suggested that the occurrence of C₄ species was common in the more arid region of China, so it was considered that environmental conditions played significant roles in the distribution and ecophysiological features of different photosynthetic types and even changed the photosynthetic pathways. Wang *et al.* (2001) reported the presence of PEPCase in all green organs of wheat. Furthermore, this activity was significantly higher in each non-leaf organ (Xu *et al.*, 2003). As PEPCase activity was positively and negatively regulated by metabolites (Kai *et al.*, 2003), the ubiquity of these C₄ cycle enzymes in C₃ plants strongly indicated that these 'C₃ isoforms' served as the starting point for the evolution of the C₄ genes (Monson, 1999). C₄ enzymatic activities were measured on *H. scoparium* and demonstrated the existence of C₄ photosynthesis both in leaves and assimilating

shoots, although lacking Kranz-anatomy in desert areas. Meanwhile, determination of enzymatic activity showed that leaves and assimilating shoots of *H. scoparium* had the properties of NAD-ME biochemical subtypes. C₄ plants of NAD-ME subtypes were known to be more resistant to stress than the species of the NADP-ME subtypes (Pyankov & Vakhrusheva, 1989, Gamaley *et al.*, 1992), suggesting the trend of evolution of C₄ pathway in C₃ desert plant was more resistant to arid stress.

The activity of PEPCase in leaves had a minor elevation, while the activity of NADP-ME was declined faintly with the increase of water stress. The activity of NAD-ME in leaves increased and was higher than that of NADP-ME. However, both of them in assimilating shoots were all increased with the increase of water stress, which was consistent with the variations of antioxidative enzymes in assimilating shoots. In addition, photosynthetic key enzymatic activities of leaves and assimilating shoots almost increased with the increase of water stress, suggesting that photosynthetic latent capacity may be advanced in *H. scoparium* with the increase of water stress. Although the photosynthetic rate of *H. scoparium* declined and stomatal conductance decreased in midday (Su, 2003), the activities of C₄ photosynthetic key enzymes increased, suggesting that the decline of photosynthetic rate was caused by stomata factor but not the activities of photosynthetic enzymes. Furthermore, plants may perform C₄ pathway by supplied malate from root systems (Hibberd & Quick, 2002) to improve photosynthetic inactivation at midday.

Water, heat and light as stress factors interacted to modulate photosynthesis (Jagtap *et al.*, 1998). C₄ photosynthetic key enzymes and antioxidative enzymes in leaves were always higher than those in assimilating shoots. The presence of the more severe water stress in 3.4 m DGW (Table 1) and the relative high C₄ photosynthetic carbon assimilating activities (Fig. 4) in leaves might lead to the production of active oxygen species and induce the high activities of active oxygen detoxifying enzymes. High irradiance results in an increase of high respiration rate in leaves (Yamazaki *et al.*, 1999), higher CO₂ concentration in cellular result in an increase of PEPCase activity. It was emphasized that water sensitivity was different between the two photosynthetic organs, so an increase of the metabolic pathway and the degradation of many scavenging enzymes induced the accumulation of active oxygen species, and especially, the production of H₂O₂ during drought stress may also result from the catalytic activity of glycolate oxidase in peroxisomes during photorespiration (Osmond, 1981) inactivated enzymatic activities of C₃ pathway (Wang, 2001), became a survival reason for the occurrence of C₄ pathway under stress. But the changes of photosynthetic contribution in leaves and assimilating shoots were dramatically different. Although the activities of C₄ photosynthetic key enzymes in leaves were always higher than that in assimilating shoots of *H. scoparium*, indicating that leaves of *H. scoparium* were the main photosynthetic and antioxidative organs at actual water gradient. However, C₄ pathway was presented with enhanced activities as the increase of water stress in assimilating shoots of *H. scoparium*, it was proposed that C₄ photosynthetic response in assimilating shoots was more positive than that in leaves as water condition became deteriorated rapidly. It might be the adaptable mechanism and the evolutional direction of the organs of the C₃ desert species in exacerbated water circumstance in desert areas.

In conclusion, the findings demonstrated that the desert species of Leguminosae was characterized by performing C₄ biochemical pathways of CO₂ fixation, not only in leaves, but also in assimilating shoots. Both photosynthetic pathway types appeared adaptation to

desert environments and this species showed environmentally induced changes in their photosynthetic responses consistent with desert adaptation. The multiple independent origins of C₄ photosynthesis suggested that the evolution of a C₃ into a C₄ species must have been relatively easy in genetic terms (Westhoff & Gowik, 2004). Kellogg (1999) and Monson (1999) provided strong evidence that even within a single taxon, i.e., the Gramineae, the transition from C₃ to C₄ may have occurred more than once. The results indicated that the occurrence of C₄ photosynthetic pathway was an adaptation in many of these species. In addition, the organ diversity of C₄ photosynthesis in C₃ desert species was an important factor for plant growth, survival and reproduction in arid regions of China. Simultaneity, the antioxidative enzymes played an important role in the protection of plants from oxidants during the C₃-CAM shift induced by water stress (Castillo, 1996). This study demonstrated that antioxidative enzymatic system seemed to be very important in the protection against oxidants, both increasing considerably antioxidative enzymes and C₄ photosynthetic enzymes under increasing water stress in C₃ plant.

Abbreviations: **APX**, ascorbate peroxidase; **CAT**, catalase; **DGW**, depth of ground water; **DM**, dry mass; **DTT**, 1,4-dithiothreitol; **FM**, fresh mass; **NAD-ME**, NAD⁺-malate enzyme; **NADP-ME**, NADP⁺-malate enzyme; **NBT**, nitroblue tetrazolium; **PEPCase**, phosphoenolpyruvate carboxylase; **PFD**, photon flux density; **POD**, peroxidase; **PVP**, polyvinylpyrrolidone; **RWC**, relative water content; **SM**, saturated mass; **SOD**, superoxide dismutase; **WUE**, water use efficiency.

Acknowledgements

We are grateful for the funding supported by National Natural Scientific Foundation of China (90102015, 30170161) and International Project of Cooperation in China and Greece (2003DFB00034). We thank J.M. Deng for assistance in the field and C.M. Zhao for data support. Comments from Professor M. Ashraf, the reviewer and H.M. Yang have helped to improve the manuscript.

References

- Asada, K. 1992. Ascorbate peroxidase - a hydrogen peroxidescavenging enzyme in plants. *Physiologia Plantarum*, 85: 235-241.
- Barrs, H.D. and P.E. Weatherley. 1962. A re-examination of the relative turgidity technique for the estimating of water deficits in leaves. *Australian Journal of Biology Science*, 15: 413-428.
- Beauchamp, C. and I. Fridovich. 1971. Superoxide dismutase: improved assays and an assay applicable to acrylamide gels. *Analytical Biochemistry*, 44: 276-287.
- Bradford, M.M. 1976. A rapid and sensitive method for the quantification of microgram quantities of protein utilizing the principle of protein-dye binding. *Analytical Biochemistry*, 72: 248-254.
- Cakmak, I. and H. Marschner. 1992. Magnesium deficiency and high light intensity on enhance activities of superoxide dismutase, ascorbate peroxidase and glutathione reductase in bean leaves. *Plant Physiology*, 98: 1222-1227.
- Castillo, F.I., I. Penel and H. Greppin. 1984. Peroxidase release induced by ozone in *Sedum album* leaves. *Plant Physiology*, 74: 846-851.
- Castillo, F.J. 1996. Antioxidative protection in the inducible CAM plant *Sedum album* L., following the imposition of severe water stress and recovery. *Oecologia*, 107: 469-477.
- Chance, B. and A.C. Maehly. 1955. Assay of catalases and peroxidases. *Methods in Enzymology*, 2: 764-817.

Chen, Q.C., G.L. Zhang and Y.W. Sun. 1961. A preliminary study on ecological and morphologic anatomical features of preponderant species of plant community along middle and lower reaches of Sule river. *Journal of Lanzhou university (natural science)*, 12: 61-96.

Das, S., S.K. Sawhney, D.R. Sood and R. Singh. 1986. Properties of NADP-malic enzyme from pod-walls of chickpea. *Physiologia Plantarum*, 68: 308-314.

Dhindsa, R.S. and W. Matowe. 1981. Drought tolerance in two mosses: correlated with enzymatic defense against lipid peroxidation. *Journal of Experimental Botany*, 32: 79-91.

Fryer, M.J., J.R. Andrews, K. Oxborough, D.A. Blowers and N.R. Baker. 1998. Relationships between CO₂ assimilation, photosynthetic electron transport and active O₂ metabolism in leaves of maize in the field during periods of low temperature. *Plant Physiology*, 116: 571-580.

Gamaley, Y.V., T.A. Glagoleva, K.G. Koltschevsky and M.V. Chulanovskaya. 1992. The C₄-syndrome types, ecology and evolution in connection with phylogeny of the Chenopodiaceae and Poaceae. *Botanicheskii Zhurnal*, 77: 1-12.

Gulías, J., J. Flexas, A. Abadía and H. Medrano. 2002. Photosynthetic response to water deficit in six Mediterranean sclerophyll species: possible factors explaining the declining distribution of *Rhamnus ludovici-salvatoris*, an endemic Balearic species. *Tree Physiology*, 22: 687-697.

Gupta, S.D. and S. Datta. 2003. Antioxidant enzyme activities during in vitro morphogenesis of gladiolus and the effect of application of antioxidants on plant regeneration. *Biologia Plantarum*, 47: 179-183.

Guo, Z.Z. and M. Zhao. 1988. Establishment of laboratory for evapotranspiration. *Journal of Desert Research*, 8: 48-56.

Hibberd, J.M. and W.P. Quick. 2002. Characteristics of photosynthesis in stems and petioles of C₃ flowering plants. *Nature*, 415: 451-454.

Jagtap, V., S. Bhargava, P. Streub and J. Feierabend. 1998. Comparative effect of water, heat and light stresses on photosynthetic reactions in *Sorghum bicolor* (L.) Moench. *Journal of Experimental Botany*, 49: 1715-1721.

Kai, Y., H. Matsumura and K. Izui. 2003. Phosphoenolpyruvate carboxylase: three dimensional structure and molecular mechanisms. *Archives of Biochemistry and Biophysics*, 414: 170-179.

Kellogg, F.A. 1999. Phylogenetic aspects of the evolution of C₄ photosynthesis. In: *C₄ Plant Biology*. (Eds.): R.F. Sage & R.K. Monson. Academic Press, San Diego. pp. 411-444.

Ku, M.S.B., J.R. Wu, Z.Y. Dai, R.A. Scott, C. Chu and G.E. Edwards. 1991. Photosynthetic and photorespiratory characteristics of *Flaveria* species. *Plant Physiology*, 96: 518-528.

Lambers, H., F.S. Chapin and T.L. Pons. 1998. *Plant Physiological Ecology*. Springer-Verlag, New York.

Li, G.Y., G.X. Gao, Y.L. Lu and X.Z. Ji. 1995. Observation on the morphology and anatomical structure of three bushes. *Research of soil and water conservation* 2: 141-145.

Li, W.H., Q.T. Lu, N.B. Hao, Q.Y. Ge, Q.D. Zhang, G.M. Jiang, W.G. Du and T.Y. Kuang. 2000. The relation between C₄ pathway enzymes and PSII photochemical function in soybean. *Acta Botanica Sinica*, 42: 689-692.

Li, M., G.X. Wang and J.S. Lin. 2003. Application of external calcium in improving the PEG-induced water stress tolerance in liquorice cells. *Botanical Bulletin of Academia Sinica*, 44: 275-284.

Lin, C.C. and C.H. Kao. 2000. Effect of NaCl on H₂O₂ metabolism in rice leaves. *Plant Growth Regulation*, 30: 151-155.

Liu, X.Q., R.Z. Wang and Y.Z. Li. 2004. Photosynthetic pathway types in rangeland plant species from Inner Mongolia, North China. *Photosynthetica*, 42: 339-344.

Llorens, L., J. Peñuelas and I. Filella. 2003. Diurnal and seasonal variations in the photosynthetic performance and water relations of two co-occurring Mediterranean shrubs, *Erica multiflora* and *Globularia alypum*. *Physiologia Plantarum*, 118: 84-95.

Lyshede, O.B. 1979. Xeromorphic features of three stem assimilating in relation to their ecology. *Botanica Journal of the Linnean Society*, 78: 85-98.

Menconi, M., C.L.M. Sgherri, C. Pinzino and F. Navari-Izzo. 1995. Activated oxygen production

and detoxification in wheat plants subjected to a water deficit programme. *Journal of Experimental Botany*, 46: 1123-1130.

Mishra, N.P., R.K. Mishra and G.S. Singhal. 1993. Changes in the activities of anti-oxidant enzymes during exposure of intact wheat leaves to strong visible light at different temperatures in the presence of protein synthesis inhibitors. *Plant Physiology*, 102: 903-910.

Mittler, R., E. Merquiol, E. Hallak-Herr, S. Rachmilevitch, A. Kaplan and M. Cohen. 2001. Living under a 'dormant' canopy: a molecular acclimation mechanism of the desert plant *Retama raetam*. *The Plant Journal*, 25: 407-416.

Monson, R.K. 1999. The origins of C₄ genes and evolutionary pattern in the C₄ metabolic phenotype. In: *C₄ Plant Biology*. (Eds.): R.F. Sage & R.K. Monson. Academic Press, San Diego. pp.377-410.

Morgan, J.A., H.W. Hunt, C.A. Monz and D.R. LeCain. 1994. Consequences of growth at two carbon dioxide concentrations and two temperatures for leaf gas exchange in *Pascopyrum smithii* (C₃) and *Bouteloua gracilis* (C₄). *Plant, Cell & Environment*, 17: 1023-1033.

Osmond, C.B. 1981. Photorespiration and photoinhibition: some implications for the energetics of photosynthesis. *Biochimica et Biophysica Acta*, 639: 77-98.

Pyankov, V.I. and D.V. Vakhrusheva. 1989. Pathways of primary CO₂ fixation in C₄-plants of the family Chenopodiaceae from the arid zone of Central Asia. *Soviet Plant Physiology*, 36: 178-187.

Sairam, R.K. and D.C. Saxena. 2000. Oxidative stress and antioxidants in wheat genotypes: Possible mechanism of water stress tolerance. *Journal of Agronomy and Crop Science*, 184: 55-61.

Sairam, R.K. and G.C. Srivastava. 2001. Water stress tolerance of wheat (*Triticum aestivum* L.): Variations in hydrogen peroxide accumulation and antioxidant activity in tolerant and susceptible genotypes. *Journal of Agronomy and Crop Science*, 186: 63-70.

Sairam, R.K., G.C. Srivastava and D.C. Saxena. 2000. Increased antioxidant activity under elevated temperature: a mechanism of heat stress tolerance in wheat genotypes. *Biological Plantarum*, 43: 245-251.

Sayre, R.T., R.A. Kennedy and D.J. Pringnitz. 1979. Photosynthetic enzyme activities and localization in *Mollugo verticillata* population differing on the leaves of C₃ and C₄ cycle operations. *Plant Physiology*, 64: 293-299.

Sharma, P.K. and G.S. Singhal. 1992. Influence of photoinhibition of photosynthesis on lipid peroxidation. *Journal of Photochemical Photobiology*, 13: 83-94.

Singal, H.R. and R. Singh. 1986. Purification and properties of PEP carboxylase from immature pods of chickpea. *Plant Physiology*, 80: 369-373.

Singal, H.R., I.S. Sheoran and R. Singh. 1987. Photosynthetic carbon fixation characteristics of fruiting structures of *Brassica campestris* L. *Plant Physiology*, 83: 1043-1047.

Singh, B.R. and G.S. Singhal. 1984. Effect of aliphatic alcohols on the temperature induced absorption changes in barley chloroplasts. *Photobiochem Photobiophys*, 8: 73-84.

Singh, R. 1993. Photosynthetic characteristics of fruiting structures of cultivated crops. In: *Photosynthesis: Photoreactions to Plant Productivity*. (Eds.): Y.P. Abrol, P. Mohanty and Govindjee. Kluwer Academic Publishers, Dordrecht, The Netherlands. pp. 289-415.

Streb, P., E. Tel-or and J. Feierabend. 1997. Light stress effects and antioxidative protection in two desert plants. *Functional Ecology*, 11: 416-424.

Su, P.X. 2003. Studies on Photosynthesis and Water Use Efficiency of Desert Plants *Haloxylon ammodendron* and *Calligonum mongolicum*. PhD dissertation, Cold and Arid Regions Environmental and Engineering Research Institute, the Chinese Academy of Sciences, Lanzhou, 40-47.

Su, P.X., H.S. Chen and Q.S. Li. 2003. Characteristics of $\delta^{13}\text{C}$ values of desert plants and their water utilization efficiency indicated by $\delta^{13}\text{C}$ values in the Desert of Central Hexi Corridor Region. *Journal of Glaciology and Geocryology*, 25: 597-602.

Thakar, J. and S. Bhargava. 1999. Seasonal variation in antioxidant enzymes and the sprouting response of *Gmelina arborea* Roxb. Nodal sectors cultured *in vitro*. *Plant Cell Tissue Organ Cultivation*, 59: 181-187.

Ting, I.P. and C.B. Osmond. 1973. Photosynthetic phosphoenolpyruvate carboxylase characteristics of alloenzymes from leaves of C₃ and C₄ plants. *Plant Physiology*, 51: 439-447.

Upadhyaya, A., T.D. Davis, M.H. Larsen, R.H. Waker and N. Sankhla. 1990. Uniconazole-induced thermo-tolerance in soybean seedling root tissues. *Plant Physiology*, 79: 78-84.

Voznesenskaya, E.V., V.R. Franceschi, O. Kuirats, H. Freitag and G.E. Edwards. 2001. Kranz anatomy is not essential for terrestrial C₄ plant photosynthesis. *Nature*, 414: 543-546.

Wang, A.G. 2001. Oxygen metabolism of plants. In: *Plant physiology and molecular biology*. Academic Press, Beijing, pp. 366-389.

Wang, Z.M., A.L. Wei and D.M. Zheng. 2001. Photosynthetic characteristics of non-leaf organs of winter wheat cultivars differing in ear type and their relationship with grain mass per ear. *Photosynthetica*, 39: 239-244.

Westhoff, P. and U. Gowik. 2004. Evolution of C₄ phosphoenolpyruvate carboxylase. Genes and proteins: a case study with the genus *Flaveria*. *Annals of Botany*, 93: 13-23.

Willekens, H., D. Inzé, M. Van Montagu and W. Van Camp. 1995. Catalases in plants. *Molecular Breeding*, 1: 207-228.

Xu, X.L., Y.H. Zhang and Z.M. Wang. 2003. Effect of heat stress during grain filling on phosphoenolpyruvate carboxylase and ribulose-1,5-bisphosphate carboxylase/oxygenase activities of various green organs in winter wheat. *Photosynthetica*, 42: 317-320.

Yamazaki, J. and Y. Kamimura. 2001. Relationship between photosystem stoichiometries and changes in active oxygen scavenging enzymes in natural grown rice seedlings. *Plant Growth Regulation*, 00: 1-8.

Yamazaki, J., Y. Kamimura and Y. Sugimura. 1999. Changes in photosynthetic apparatus in the juvenile rice canopy and a possible function of photosystem I in the bottom leaves. *Z. Naturforsch.*, 54c: 915-922.

Zhang, J. and M.B. Kirkham. 1996. Antioxidant responses to drought in sunflower and sorghum seedlings. *New Phytologist*, 132: 361-373.

Zhao, G.D., S.R. Liu and Q.L. Ma. 2003a. Ecophysiological responses of two xerophytes *atraphaxis frutescens* and *elaeagnus angustifolia* to the change of groundwater depth in arid area I. Changes in leaf nutrient, chlorophyll, soluble sugar and starch contents. *Acta Phytocologica Sinica*, 27: 228-234.

Zhao, M., Z.Z. Guo, Y.L. Wang, A.D. Li, D.K. Zhang and B.Q. Jia. 2003b. Study on the transpiration characteristics of 10 species of plants under the different depths of groundwater level. *Arid Zone Research*, 20: 286-291.

(Received for publication 29 September 2005)