

USE OF SEA WEED AND BACTERIA IN THE CONTROL OF ROOT ROT OF MASH BEAN AND SUNFLOWER

SHAHNAZ DAWAR, SADIA ARJUMUND,
MARIUM TARIQ AND M.J. ZAKI

*Department of Botany,
University of Karachi, Karachi-75270, Pakistan.*

Abstract

A significant increase in growth parameters in terms of shoot length, shoot weight and a significant reduction in infection of *Fusarium* spp, *Rhizoctonia solani* and *Macrophomina phaseolina* was observed where *Rhizobium meliloti* treated seeds of mash bean and sunflower were used in *Melanothanus afaqhusainii* amended soil. Combined use of *Pseudomonas aeruginosa* with sea weeds viz., *Melanothanus afaqhusainii*, *Padina tetrastromatica*, *Cystoclonium purpureum* and *Hypnea valentiae* significantly reduced the infection of *M. phaseolina* and *R. solani* on mash bean and sunflower followed by *R. meliloti* and *B. subtilis* in combination with *C. purpureum* and *M. afaqhusainii* respectively on mash bean and sunflower. Seed treatment was found more effective method in controlling the root infecting fungi as compared to soil treatment.

Introduction

Sea weeds have received the attention of plant scientists throughout the world (Atzman *et al.*, 1994, Staden *et al.*, 1995), since it contains greater potash and nitrogen as compared to farm yard manure (Chapman & Chapman, 1980) and thus provide better plant growth. Pratt *et al.*, (1951) reported antimicrobial activity of some sea weeds. The marine plants constitute a prodigious source of new and structurally complex secondary metabolites (Atta-ur-Rehman *et al.*, 1991). In many countries, commercial sea weed fertilizer has been marketed as soil additives and foliar spray (Paracer *et al.*, 1987). Many plant growth promoting rhizobacteria (PGPRS) have a beneficial effect on plants, besides biological control of soil borne pathogens, induce systemic resistance to plant pathogens, phytoharmonic productions and improvement of nutrition and water uptake of plants (Seuk Bae *et al.*, 2000). Of the *Pseudomonas* species *P. aeruginosa* has been found as an effective biocontrol agent of root rot pathogens (Izhar *et al.*, 1995). Similarly *Rhizobia*, the root nodulating bacteria, are also known to control soil borne root infecting fungi (Zaki & Ghaffar, 1987; Ehteshamul-Haque & Ghaffar, 1993; Siddiqui *et al.*, 1998). Experiments were therefore carried out to study the use of sea weeds and bacteria in the control of root rot of mash bean (*Vigna mungo* (L.) and sunflower (*Helianthus annus* L.).

Materials and Methods

Seaweeds like *Melanothanus afaqhusainii*, *Padina tetrastromatica*, *Cystoclonium purpureum* and *Hypnea valentiae* were collected from Buleji, Karachi, washed with tap water, dried under shade, powdered in an electric grinder and kept at room temperature. Cultures of *Rhizobium meliloti* (R5), *Pseudomonas aeruginosa* (P-58), *Bacillus subtilis* (B-35) were obtained from Department of Botany, University of Karachi. Soil used for the experiment was obtained from the experimental plots of the Department of Botany, University of Karachi and sieved through 2mm sieve to discard particles. The soil used

was sandy loam (Sand, Silt, Clay; 70, 19, 11%), pH range from 7.5 - 8.1 with moisture holding capacity (MHC) of 24.04 % (Keen & Raczkowski, 1922), total nitrogen 1.5 % (Mackenzie & Wallace, 1954), total organic matter 2.4 %. Soil had natural infestation of 1-3 sclerotia of *Macrophomina phaseolina* as found by wet sieving dilution technique (Sheikh & Ghaffar, 1975), 5-10% colonization of *Rhizoctonia solani* on sorghum seeds used as baits (Wilhelm, 1955) and 3000cfu of *Fusarium* spp., as assessed by soil dilution technique (Nash & Snyder, 1962). Soil was amended with seaweeds viz., *M. afaqhusainii*, *P. tetrastromatica*, *C. purpuraeum*, *H. valentiae* @ 1 % w/w and transferred in 8 cm diameter plastic pots, each pot containing 300 g soil and soil moisture adjusted and maintained at 50 % MHC (Keen & Raczkowski, 1922). Pots were kept for 2 weeks for the decomposition of the organic substrate. Soil was drenched with 20 ml water containing cell suspension of *R. meliloti* (R5) (7.48×10^8 cells/ml), *B. subtilis* (B-35) (12.93×10^8 cells/ml), *P. aeruginosa* (P-58) (6.23×10^8 cells/ml). Mash bean and sunflower were used as test plants. Eight seeds were sown in each pot. Pots without seaweed and bacterium served as control.

In another treatment, seeds of mash bean were treated with cell suspension of *B. subtilis* containing (5.68×10^7 cells/seeds), *R. meliloti* (8.98×10^7 cells/seeds) and *P. aeruginosa* (12.43×10^7 cells/seeds) whereas in sunflower cell suspension containing (9.28×10^7 cells/seeds), (13.46×10^7 cells/seeds) and (16.98×10^7 cells/seeds) respectively were used and sown in soil amended with seaweeds. There were three replicates of each treatment and pots were randomized on a green house bench. After 30 days, roots of sunflower and mash bean were washed in a running tap water, surface sterilized in 1% Ca (OCl)₂ and then five 1 cm long root pieces were transferred on PDA plates containing penicillin @100,000/litre and streptomycin @ 20mg/l. Petri dishes were incubated for 5 days at room temperature to confirm infection of root infecting fungi. After 30 days of growth, plants were uprooted. Plants growth parameters in terms of root length, shoot length and fresh weights of shoot, root and incidence of root infecting fungi viz., *M. phaseolina*, *R. solani* and *Fusarium* spp., were recorded. Data were analyzed and subjected to analysis of variance (ANOVA) following the procedure as given by Gomez & Gomez (1984).

Results and Discussion

Significant increase in shoot length in mash bean and sunflower ($p < 0.001$) was observed where *R. meliloti*, *B. subtilis* and *P. aeruginosa* were drenched in *P. tetrastromatica* amended soil ($p < 0.01$) (Table 1). There was a significant increase in shoot length ($p < 0.05$), shoot weight ($p < 0.01$), root length and root weight ($p < 0.01$) when *B. subtilis* were drenched in *C. purpuraeum* amended soil (Table 1) whereas combined use of *P. aeruginosa* with sea weeds viz., *M. afaqhusainii*, *P. tetrastromatica*, *C. purpuraeum* and *H. valentiae* significantly reduced the infection of *M. phaseolina* and *R. solani* on mash bean and sunflower followed by *R. meliloti* and *B. subtilis* in combination with *C. purpuraeum* and *M. afaqhusainii* respectively on mash bean and sunflower (Table 1) whereas *P. aeruginosa* in combination with *H. valentiae* significantly reduced the infection of *Fusarium* spp. Sunflower and mash bean treated seeds with *R. meliloti* when used in *M. afaqhusainii* amended soil significantly increased the growth parameters in terms of shoot length, shoot weight ($p < 0.05$) and significantly reduced the infection of *Fusarium* spp., *M. phaseolina* and *R. solani* ($p < 0.05$) whereas *P. aeruginosa* treated seeds of mash bean and sunflower in *C. purpuraeum* and *H. valentiae* treated soil showed a significant increase in shoot length ($p < 0.05$) and reduced the infection of *Fusarium* spp., and *M. phaseolina* ($p < 0.05$) (Table 2).

Antagonists applied to seeds not only have the potential for protecting the seeds but being the initial colonizer of the roots, may also provide protection against root infecting pathogens (Kommedahl & Windels, 1981). In the present study seed dressing with bacteria was found more effective in the control of root infecting fungi as compared to soil treatment. Seaweeds in combination with bacteria were also effective for growth of plants. Seed dressing and soil treatment with different strains of bacteria viz., *P. aeruginosa*, *R. meliloti* and *B. subtilis* showed promising results in the control of root infecting fungi. Raaijimaker *et al.*, (2002) and Weller *et al.*, (2002) reported that plant growth promoting rhizobacteria that colonize roots improve plant growth either through direct stimulation of the plant by producing growth regulators or by suppression of pathogens. The production of certain antibiotics (Leavy *et al.*, 1992) and siderophores (Buysens *et al.*, 1996) by *P. aeruginosa* has been regarded as one of the mechanism involved in antagonism. Siddiqui *et al.*, (2000) and Siddiqui & Ehtesham-ul-Haque (2001) reported that species of *Pseudomonas* are antagonistic to soil borne plant pathogens. During the present study, reduction in disease intensity and improved plant growth as compared to untreated plants was observed in mash bean and sunflower. Chapman & Chapman (1980) and Colwell (1983) reported that potash, equal nitrogen and role of active biocontrol principles in seaweed may cause disease suppression resulting in plant growth enhancement. The seaweeds increased the efficacy of bacteria in the control of *M. phaseolina* on mash bean, while sea weed *H. valentiae* with *P. aeruginosa* significantly reduced the infection of *M. phaseolina*, *R. solani* and *Fusarium* spp., on mash bean and sunflower. This work suggests that seaweed alongwith plant growth promoting bacteria used either as seed dressing or soil drenching have greater potential in controlling root infecting fungi on crop plants.

Reference

Atta-ur-Rehman, Z. Shah, M.I. Choudhary, S.A. Abbas and M. Shameel. 1991. Stockerine, a novel linear metabolite from *Stockeyia indica*. *Fitoterapia*, 62:77-80.

Atzman, N., J.V. Staden and S.J. Van. 1994. The effect of sea weeds concentration on the growth of *Pinus pinea* seedlings. *New Forests*, 81: 279-288.

Buysens, S., K. Heungens, J. Poppe and M. Hofte. 1996. Involvement of pyochelin and pyoverdin in suppression of *Pythium*-induced damping-off of tomato by *Pseudomonas aeruginosa* 7NSK2. *Appl. Environ. Microbiol.*, 62: 865-871.

Chapman, V.J. and D.J. Chapman. 1980. *Seaweeds and their uses*. 3rd ed. Chapman and Hall, New York. pp.334.

Colwell, R.R. 1983. Biotechnology in marine science. *Science*, 222: 19-24.

Ehteshamul-Haque, S and A. Ghaffar. 1993. Use of rhizobia in the control of root rot diseases of sunflower, okra, soyabean and mungbean. *J. Phytopath.*, 138: 157-163.

Gomez, K.A. and A.A. Gomez. 1984. *Statistical procedures for agriculture research* 2nd ed. Wiley, New York pp. 680.

Izhar, I.S., S. Ehteshamul-Hque, M. Javed and A. Ghaffar. 1995. Efficacy of *Pseudomonas aeruginosa* and *Bradyrhizobium* sp., in the control of root disease in chick pea. *Pak. J. Bot.*, 27: 451-455.

Keen, B.A and H. Rackowski. 1922. The relation between clay content and certain physical properties of soil. *J. Agric. Sci.*, 11:441- 449.

Kommedahl, T and C. Windels. 1981. Introduction to microbial antagonists to specific courts of infection: Seed, seedlings and wounds. In: *Biological control in crop production*. (Ed.): G.C. Papavizas. Beltsville symposia in Agricultural Research 5. Osmum publishing, Granada: Allenheld, 227-248.

Leavy, E., Z. Eyal, I. Chet and A. Hochman. 1992. Resistance mechanisms of *Septoria tritici* to antifungal products of *Pseudomonas*. *Physiol. Mol. Pl. Pathol.*, 40: 163-171.

Mackenzie, H.A. and H.S. Wallace. 1954. The Kjeldahl determination of nitrogen: A critical study of digestion conditions, temperature, catalyst and oxidizing agents. *Aust. J. Chem.*, 7: 55-70.

Nash, S.M. and W.C. Snyder. 1962. Quantitative estimation by plate counts of propagules of the bean root rot *Fusarium* in field soils. *Phytopathology*, 52: 567-572.

Paracer, S., C. Armin, A.C. Targan and L.M. Hodgson. 1987. Effective use of marine algal products in the management of plant parasitic nematodes. *J. Nematol.*, 19: 194-200.

Pratt, R., R.H. Mautner, G.H. Gardner, Y. Sha and J. Defrenoy. 1951. Report on antibiotic activity of seaweed extracts. *J. Am. Pharm. Assoc.*, 40: 575-579.

Raaijimaker, J.M., M. Vilami and J.T. de Sauza. 2002. Antibiotic production by bacterial biocontrol agents. *Antonie Van Leeuwenhoek*, 81: 537-547.

Sheikh, A.H and A. Ghaffar. 1975. Population study of sclerotia of *Macrophomina phaseolina* in cotton field. *Pak. J. Bot.*, 7: 13-17.

Seuk Bae, Yeoung, OK-Hee Choi, Kyung-Soek park, Sang-Bum Lee and Choong-Hoe Kim. 2000. *A useful method for functional analysis of plant growth promoting rhizobacteria in the development of cucumber root system*. Plant pathology Division, National Institute of Agricultural Science and Technology, Suwon 441-707, Korea.

Siddiqui, I.A, S. Ehteshamul-Haque and A. Ghaffar. 1998. Effect of fungicides on the efficacy of *Rhizobium meliloti* and *Bradyrhizobium* sp., in the control of root infecting fungi on chick pea. *Pak. J. Bot.*, 30: 69-74.

Siddiqui, I.A., S.A. Qureshi, V. Sultana, S. Ehteshamul-Haque and A. Ghaffar. 2000. Biological control of root rot and root knot disease complex of tomato. *Plant Soil*, 227: 163-169.

Siddiqui, I.A and S. Ehteshamul-Haque. 2001. Multiplication of *Pseudomonas aeruginosa* an talc-based formulation for use in the control of root rot and root knot disease in tomato. *Trop. Sci*, 41: 163-167.

Staden, J.V., R.P. Beckett, M.J. Rijkenberg and J.S. Van. 1995. Effect of seaweed concentration on the growth of seedlings of three species of *Eucalyptus*. *South-African. J. Bot.*, 61: 169-172.

Weller, D.M., J.M. Raaijimakers, B.B.M. Gardner and L.S. Thomashow. 2002. Microbial population responsible for specific soil suppressiveness to plant pathogens. *Annu. Rev. Phytopathol.*, 40: 309-348.

Wilhelm. S. 1955. Longevity of the *Verticillium* wilt fungus in the laboratory and field. *Phytopathology*, 45:180-181.

Zaki, M.J. and A.Ghaffar. 1987. Effect of *Rhizobium* spp., on *Macrophomina phaseolina*. *Pak. J. Sci & Ind. Res.*, 30: 305-306.

(Received for Publication 2 January 2007)