

PHYCOCHEMISTRY AND BIOACTIVITY OF TWO STONEWORT ALGAE (CHAROPHYTA) OF SINDH

B. GHAZALA, B. NAILA, MUSTAFA SHAMEEL, SALEEM SHAHZAD AND
SULTAN MAHMOOD LEGHARI¹

Department of Botany
University of Karachi, Karachi-75270, Pakistan

Abstract

Two stonewort green algae, *Chara contraria* A. Braun ex Kützinger and *Nitella flexilis* (Linnaeus) C. A. Agardh were collected from Kinjhar and Haleji lakes, Distt. Thatta, Sindh during October to December 1999. Their methanol extracts revealed the presence of a variety of saturated and unsaturated fatty acids (FAs) by GC-MS and β -sitosterol by EI-MS & ¹H-NMR. In *C. contraria*, 9 saturated, 5 mono- and 2 tri-unsaturated FAs were detected, and *N. flexilis* contained 5 saturated, 7 mono-, 2 di-, 2 tri- and 3 poly-unsaturated FAs; while 4 FAs remained unidentified. The C13:1, C14:0, C16:0, C16:3, C17:2, C17:3, C29:0 and C29:3 acids were most commonly present in appreciable quantities (8.053-20.423 %) in them. Their methanol extracts showed little antibacterial activity but a strong antifungal activity against 6 of the 10 tested fungal pathogens. They displayed no bioactivity in brine shrimp bioassay and insecticidal tests. *Chara contraria* exhibited appreciable phytotoxicity (6.66 %) and antitumour activity (31 %) but *N. flexilis* did not, they showed specific differences.

Introduction

A large number of green seaweeds growing at the seashore of Karachi and the adjacent coastal areas of Sindh have been investigated phycochemically (Usmanghani *et al.*, 1985; Qasim, 1986; Shameel, 1987, 1993; Aliya *et al.*, 1995; Aliya & Shameel 1999, 2003), and their bioactivities were studied (Usmanghani *et al.*, 1984; Naqvi *et al.*, 1992; Aliya *et al.*, 1994; Atta-ur-Rahman *et al.*, 1997; Rizvi & Shameel, 2003). Only a few studies were conducted on stonewort green algae of this area (Khaliq-uz-Zaman *et al.*, 1998, 2001; Shameel *et al.*, 1999). This paucity of knowledge initiated a research program to compare the observations made on green seaweeds with their freshwater counterparts of Sindh (Ghazala *et al.*, 2003, 2004a-d). The present investigation is a continuation of this program in which taxonomy, phycochemistry and bioactivity of two stonewort algae *Chara contraria* A. Braun ex Kützinger and *Nitella flexilis* (Linnaeus) C. A. Agardh have been studied.

Materials and Methods

Collection of material: Specimens were collected during October to December 1999 from the bottom of a shallow water pond at Hyderabad and from sandy bottom of Kinjhar and Haleji lakes of District Thatta (Sindh) at a depth of 10-40 cm.

Detection of fatty acids: The algae under investigation weighting 1 kg fresh weight (*C. contraria*) and 4811.2 g dry wt. (*N. flexilis*) were percolated with *n*-hexane: chloroform (1:1, v/v) in a respirator for two weeks. The extract so obtained was reduced under vacuum and partitioned between EtOAc and water (1:1, v/v), which on evaporation

¹Department of Freshwater Biology & Fisheries, University of Sindh, Jamshoro-76080, Pakistan.

yielded 15 g (*C. contraria*) and 10 g (*N. flexilis*) of residue. It was saponified, refluxed, esterified, methylated and finally subjected to GC-MS, as described in Ghazala *et al.*, (2004a).

Isolation of sterol & terpene: Thalli were thoroughly washed with tap water to remove debris and animal castings. Dried and chopped material weighing 650.5 g was soaked in methanol for almost three weeks. The MeOH fraction on evaporation under vacuum gave 38.4 g thick crude algal material, and the slurry was loaded on to silica gel column for chromatography eluting with *n*-hexane: chloroform and finally with methanol in increasing order of polarity by 2 % (v/v) respectively, as given by Ghazala *et al.*, (2003).

Bioactivity tests: The methodologies for bactericidal activity, antifungal bioassay, food poisoning and phytotoxic activities, brine shrimp bioassay and insecticidal activity were the same as have been described previously (Ghazala *et al.*, 2003, 2004a).

Antitumour activity: Potato tubers obtained from local markets were sterilized by immersion in sodium hypochlorite for 20 minutes. After removing their ends potatoes were soaked for 10 minutes or more in Clorox. A core of the tissue was extracted from each tuber with sterilized cork borer (1.5 cm) radius. The 2 cm pieces were removed from each end and discarded; the remainder of the cylinder was cut into 0.5 cm discs with a surface-sterilized scalpel and knife. The discs were then transferred to 1.5 % agar plate (1.5 g of Merk agar dissolved in 100 mL of distilled water, autoclaved and 20 mL of it poured into each sterile Petri plate). The extracts were dissolved in 2 mL of DMSO. Plates were incubated at 27 °C for 12-20 days after which the tumours produced were counted with the naked eye. Significant activity was indicated when two or more independent assays gave consistent 20 % or greater inhibition. Details of the procedure may be found in Parveen (1997).

Results

***Chara contraria* A. Braun ex Kützinger:** Thallus coarse, usually heavily encrusted, 20-30 cm high; stem up to 730 µm in diameter, stem cortex regularly diplostichous; stipulates in double whorls, paired with each leave; leaves 6-8 in a whorl, incurved or spreading; terminal cell of the stem and leaves ecorticated; sex organs monoecious, both organs produced at the same node, antheridia 311-470 µm in diameter, oogonia 800 µm long, spiral cells showing 13-15 convolutions (Fig. 1).

***Nitella flexilis* (Linnaeus) C.A. Agardh:** Thallus 30-40 cm high; stem and leaves uncorticated, encrusted with lime; stem up to 0.8 mm thick; internodes 4.5-9.0 cm long; leaves up to 6 in a whorl, forked; last segment unicellular; stipule-circle lacking; antheridia borne above oogonia; crown developed on oogonium by tips of 5 two-celled coiled tubes (Fig. 2).

Detection of fatty acids: The GC-MS of the methylated fatty acids (FAs) revealed the presence of 9 saturated and 7 unsaturated FA-methyl esters in *C. contraria* and 9 saturated and 14 unsaturated FA-methyl esters in *N. flexilis*, while 4 FAs remained unidentified (Tables 1 & 2).




Fig. 1. *Chara contraria*: **a.** branchlet with globule and nucule, **b.** main thallus (scale: 400 μ m).




Fig. 2. *Nitella flexilis* (Linnaeus) C.A. Agardh: **a.** part of thallus showing whorl of forked leaves, **b.** reproductive organs (x 50).

Extraction of sterol and terpene: Pure sterol compound was eluted in a mixture of *n*-hexane: chloroform (90:10) from column and purified on preparative thick layer silica gel glass plates in solvent system *n*-hexane: chloroform (80:20). Purity was then checked on TLC card in the above solvent system and by spraying with $\text{Ce}(\text{SO}_4)_2$ a pinkish red spot was found. After ascertaining the purity of the compound the isolated sterol was subjected to spectral analysis (EIMS and $^1\text{H-NMR}$). In this way β -sitosterol was identified, but *trans*-phytol could not be detected (Table 3).

Bioactivities: Methanol extract of *C. contraria* showed no bioactivity against any one of the 10 tested bacterial species but displayed a strong antifungal activity against 6 of the 10 tested fungal organisms (Table 5). It exhibited phytotoxicity against *Lemna minor* of low quality at the three concentrations and its cytotoxic activity against brine shrimp was observed to be non-significant. It revealed an appreciable antitumour activity, which enhanced from 10 to 31 % with increasing concentration of the extract. The extract of *N. flexilis* exhibited antibacterial activity against 3 of the 10 tested bacterial species and antifungal activity against 6 of the 10 tested fungal organisms (Table 3). Its cytotoxic and insecticidal activities were found to be non significant.

Table 1. Fatty acids detected as methyl esters in three fractions (A-C) of *Chara contraria* and four fractions (A-D) of *Nitella flexilis*.

Acid type	Systematic name	Common name	Molecular formula	Mol. wt.	Rel. % age
<i>Chara contraria</i>					
Fraction A					99.754
C14:0	<i>n</i> -Tetradecanoate	Myristate	C ₁₅ H ₃₀ O ₂	242	2.448
C15:0	<i>n</i> -Pentadecanoate	Pentadecylate	C ₁₆ H ₃₂ O ₂	256	1.53
C16:0	<i>n</i> -Hexadecanoate	Palmitate	C ₁₇ H ₃₄ O ₂	270	6.12
C17:3	5,8,11-Heptadecatrienoate	-	C ₁₈ H ₃₀ O ₂	278	38.862
C18:1	8-Octadecenoate	Oleate	C ₁₉ H ₃₆ O ₂	296	5.506
C18:0	9-Methyl-heptadecanoate	-	C ₁₉ H ₃₈ O ₂	298	2.448
C19:1	10-Nonadecenoate	Nonadecylenate	C ₂₀ H ₃₈ O ₂	310	1.836
C20:1	9-Eicosenoate	Gadoleate	C ₂₁ H ₄₀ O ₂	324	7.65
C22:0	Docosanoate	Behenate	C ₂₃ H ₄₆ O ₂	354	3.06
C29:3	Nonacosatrienoate	-	C ₃₀ H ₅₄ O ₂	446	7.956
-	Unidentified	-	-	330	7.65
-	Unidentified	-	-	410	9.18
-	Unidentified	-	C ₂₉ H ₅₂ O ₂	432	5.508
Fraction B					99.918
C20:1	9-Eicosenoate	Gadoleate	C ₂₁ H ₄₀ O ₂	324	4.697
C22:1	11-Docosenoate	Cetoleate	C ₂₃ H ₄₄ O ₂	352	14.518
C23:1	Tricosenoate	-	C ₂₄ H ₄₆ O ₂	366	3.843
C24:0	<i>n</i> -Tetracosanoate	Lignocerate	C ₂₅ H ₅₀ O ₂	382	17.934
C29:3	Nonacosatrienoate	-	C ₃₀ H ₅₄ O ₂	446	53.375
C29:0	<i>n</i> -Nonacosanoate	-	C ₃₀ H ₆₀ O ₂	452	5.551
Fraction C					99.735
C17:0	<i>n</i> -Heptadecanoate	Margarate	C ₁₈ H ₃₆ O ₂	284	9.81
C18:0	9-Methyl-heptadecanoate	-	C ₁₉ H ₃₈ O ₂	298	17.004
C29:1	10-Nonadecenoate	Nonadecylenate	C ₂₀ H ₃₈ O ₂	310	7.194
C21:0	<i>n</i> -Heneicosanoate	-	C ₂₂ H ₄₄ O ₂	340	23.871
C22:1	11-Docosenoate	Cetoleate	C ₂₃ H ₄₄ O ₂	452	41.856
<i>Nitella flexilis</i>					
Fraction A					99.99
C11:0	Undecanoate	Undecylate	C ₁₂ H ₂₄ O ₂	200	0.6711
C14:1	9-Tetradecenoate	Mysistoleate	C ₁₅ H ₂₈ O ₂	240	2.013
C15:3	3,7,11-Trimethyl-2,6,10-dodecatrienoate	-	C ₁₆ H ₂₆ O ₂	250	10.066
C15:0	<i>n</i> -Pentadecanoate	Pentadecylate	C ₁₆ H ₃₂ O ₂	256	6.039

Table 1 (Cont'd.)

Acid type	Systematic name	Common name	Molecular formula	Mol. wt.	Rel. % age
C16:3	Hexadecatrienoate	-	C ₁₇ H ₂₈ O ₂	264	32.212
C17:2	9,12-Heptadecadienoate	-	C ₁₈ H ₃₂ O ₂	280	42.279
C18:5	Octadecapentaenoate	-	C ₁₉ H ₂₈ O ₂	288	3.355
C18:2	5,8-Octadecadienoate	-	C ₁₉ H ₃₄ O ₂	294	3.355
Fraction B					99.99
C14:1	9-Tetradecenoate	Mysistoleate	C ₁₅ H ₂₈ O ₂	240	21.333
C14:0	<i>n</i> -Tetradecanoate	Mysistate	C ₁₅ H ₃₀ O ₂	242	25.997
C16:0	<i>n</i> -Hexadecanoate	Palmitate	C ₁₇ H ₃₄ O ₂	270	41.329
C18:1	8-Octadecenoate	Oleate	C ₁₉ H ₃₆ O ₂	296	1.999
C18:0	<i>n</i> -Octadecanoate	Stearate	C ₁₉ H ₃₈ O ₂	298	1.333
-	Unidentified	-	-	274	7.999
Fraction C					99.99
C8:1	7-Octynoate	-	C ₉ H ₁₄ O ₂	154	1.923
C13:1	9-Tridecenoate	Decylacrylate	C ₁₄ H ₂₆ O ₂	226	59.613
C14:1	13-Tetradecynoate	-	C ₁₅ H ₂₆ O ₂	238	1.923
C15:4	Pentadecatetraenoate	-	C ₁₆ H ₂₄ O ₂	248	1.923
C16:1	9-Hexadecenoate	Palmitoleate	C ₁₇ H ₃₂ O ₂	268	17.307
C18:5	Octadecapentaenoate	-	C ₁₉ H ₂₈ O ₂	288	3.846
C18:2	5,8-Octadecadienoate	-	C ₁₉ H ₃₄ O ₂	294	2.884
C19:1	10-Nonadecenoate	Nonadecylenate	C ₂₀ H ₃₈ O ₂	310	1.923
-	Unidentified	-	-	190	4.807
-	Unidentified	-	-	318	3.846
Fraction D					99.99
C13:1	9-Tridecenoate	Decylacrylate	C ₁₄ H ₂₆ O ₂	226	14.691
C14:0	<i>n</i> -Tetradecanoate	Myristate	C ₁₅ H ₃₀ O ₂	242	29.855
C15:3	3,7,11-Trimethyl-2,6,10-dodecatrienoate	-	C ₁₆ H ₂₆ O ₂	250	12.795
C15:0	<i>n</i> -Pentadecanoate	Pentadecylate	C ₁₆ H ₃₂ O ₂	256	3.791
C16:0	<i>n</i> -Hexadecanoate	Palmitate	C ₁₇ H ₃₄ O ₂	270	23.695
C18:5	Octadecapentaenoate	-	C ₁₉ H ₂₈ O ₂	288	4.739
C18:2	5,8-Octadecadienoate	-	C ₁₉ H ₃₄ O ₂	294	3.791
C18:0	<i>n</i> -Octadecanoate	Stearate	C ₁₉ H ₃₈ O ₂	298	3.317
C22:4	6, 9,12,15-Docosatetraenoate	-	C ₂₃ H ₃₈ O ₂	346	3.317

Mol. wt. = Molecular weight, Rel. % age = Relative percentage.

Table 2. Total fatty acids analysed as methyl esters present in the methanol extracts of *Chara contraria* and *Nitella flexilis*.

Acid type	Systematic name	Molecular formula	Mol. wt.	Rel. % age	
				C.C.	N.F.
I. Saturated fatty acids				43.829	34.005
C11:0	<i>n</i> -Undecanoate	C12H24O2	200	-	0.167
C14:0	<i>n</i> -Tetradecanoate	C15H30O2	242	0.815	13.963
C15:0	<i>n</i> -Pentadecanoate	C16H32O2	256	0.509	2.457
C16:0	<i>n</i> -Hexadecanoate	C17H34O2	270	2.037	16.256
C17:0	<i>n</i> -Heptadecanoate	C18H36O2	284	3.266	-
C18:0	9-Methyl-heptadecanoate	C19H38O2	298	6.477	1.162
C21:0	<i>n</i> -Heneicosanoate	C22H44O2	340	7.949	-
C22:0	<i>n</i> -Docosanoate	C23H46O2	354	1.018	-
C24:0	<i>n</i> -Tetracosanoate	C25H50O2	382	5.972	-
C29:0	<i>n</i> -Nonacosanoate	C30H60O2	452	15.786	-
II. Monounsaturated fatty acids				15.063	30.677
a. Monoenoic fatty acids				15.063	29.711
C13:1	9-Tridecenoate	C14H26O2	226	-	18.57
C14:1	9-Tetradecenoate	C15H28O2	240	-	5.836
C16:1	9-Hexadecenoate	C17H32O2	268	-	4.326
C18:1	8-Octadecenoate	C19H36O2	296	1.833	0.499
C19:1	10-Nonadecenoate	C20H38O2	310	3.006	0.480
C20:1	9-Eicosenoate	C21H40O2	324	4.111	-
C22:1	11-Docosenoate	C23H44O2	352	4.834	-
C23:1	Tricosenoate	C24H46O2	366	1.279	-
b. Monoynoic fatty acids				-	0.96
C8:1	7-Octynoate	C9H14O2	1.54	-	0.480
C14:1	13-Tetradecynoate	C15H26O2	238	-	0.480
III. Diunsaturated fatty acids				-	13.076
C17:2	9,12-Heptadecadienoate	C18H32O2	280	-	10.569
C18:2	5,8-Octadecadienoate	C19H34O2	294	-	2.507
IV. Triunsaturated fatty acids				33.364	13.768
C15:3	3,7,11-Trimethyl-2,6,10-dodecatrienoate	C16H26O2	250	-	5.715
C16:3	Hexadecatrienoate	C17H28O2	264	-	8.053
C17:3	5,8,11-Heptadecatrienoate	C18H30O2	278	12.941	-
C29:3	Nonacosatrienoate	C30H54O2	446	20.423	-
V. Polyunsaturated fatty acids				-	4.294
C15:4	Pentadecatetraenoate	C16H24O2	248	-	0.480
C18:5	Octadecapentaenoate	C19H28O2	288	-	2.985
C22:4	6,9,12,15-Docosatetraenoate	C23H38O2	346	-	0.829
IV. Unidentified fatty acids				7.437	4.161
-	-	-	190	-	1.201
-	-	-	274	-	1.999
-	-	-	318	-	0.961
-	-	-	-	7.437	-

Mol. wt. = Molecular weight, Rel. % age = Relative percentage, C.C. = *Chara contraria*, N.F. = *Nitella flexilis*.

Table 3. Comparison of phycochemistry and bioactivity of stonewort algae.

Compounds/ Organisms	<i>Chara contraria</i>	<i>Nitella flexilis</i>
I. Natural products	%	%
B-Sitosterol	-	+
<i>Trans</i> -phytol	-	-
II. Antibacterial activity	mm	mm
<i>Bacillus cereus</i>	-	6
<i>Corynebacterium diphtheriae</i>	-	†6.5
<i>Escherichia coli</i>	-	-
<i>Klebsiella pneumoniae</i>	-	-
<i>Proteus mirabilis</i>	-	-
<i>Pseudomonas aeruginosa</i>	-	-
<i>Salmonella typhi</i>	-	-
<i>Shigella boydii</i>	-	†8
<i>Staphylococcus aureus</i>	-	-
<i>Streptococcus pyogenes</i>	-	-
III. Antifungal activity	%	%
<i>Alternaria alternata</i>	-	-
<i>Curvularia lunata</i>	2.9	3.0
<i>Drechslera australiensis</i>	-	-
<i>Fusarium solani</i>	-	-
<i>Fusarium sporotrichoids</i>	3.2	3.5
<i>Fusarium proliferatum</i>	-	-
<i>Macrophomina phaseolina</i>	3.5	3.7
<i>Rhizoctonia solani</i>	3.5	3.6
<i>Sclerotium rolfsii</i>	3.6	3.4
<i>Trichoderma harzianum</i>	3.3	3.0
IV. Phytotoxicity	%	%
500 µg/mL concentration	6.66	X
V. Brine shrimp bioassay		
1000 µg/mL LD ₅₀	-	-
VI. Insecticidal activity		
<i>Tribolium castaneum</i>	x	-
VII. Antitumour activity	%	%
100 µg	31	x

+ = Present, - = Absent /no activity, x = Not tested, †= Decrease in bacterial population/ unit area.

Discussion

The methanol extract of *C. contraria* revealed the presence of 9 saturated and 7 unsaturated FAs, while 1 FA could not be identified (Table 2). Nonacosatrienoic acid (C29:3) was found to be in largest amount (20.423 %) as compared to all the other FAs, so much so that it appeared to be the most dominant acid, while pentadecylic acid (C15:0) was present in the smallest amount (0.51 %). Other prevalent acids were nonacosanoic (C29:0, 15.786 %) and 5, 8, 11-heptadecatrienoic (C17:3, 12.94 %) acids. Unsaturated FAs were found to be in slightly larger quantity (48.427 %) than saturated FAs (43.829 %), while a proportion of 7.437 % remained unidentified. In a previous study a total number of 23 FAs were detected in *C. corallina* var. *wallichii* including a wide variety of saturated (36.5 %) and unsaturated (63.5 %) FAs and hexadecadienoic acid occurred in greatest amount (23.7 %), while pentadecylic (10.61 %) and palmitic (17.69 %) acids were the major SFAs (Khaliq-uz-Zaman *et al.*, 1998). In this way the two species of *Chara* differed from each other.

β -Sitosterol was isolated from *C. contraria*. It has also been obtained from the MeOH extract of *C. wallichii* in a dominant proportion (76.8 %) apart from three other sterols *i.e.* cholesterol, clerosterol and stigmasterol which occurred in small quantities (Shameel *et al.*, 1999). Sitosterol and fucosterol were detected as principal sterols in *N. opaca* (Heilbron, 1942); while clionasterol, an isomer of sitosterol and 28-isofucosterol were found to be the major sterols of *C. vulgaris* and *N. flexilis* (Patterson, 1972). In *C. australis* and *C. buckelii* also sitosterol was observed to be present along with several other sterols (Patterson *et al.*, 1991). It appears that β -sitosterol is the major sterol of *Chara*.

The extract of *C. contraria* exhibited a variable amount of bioactivity against a variety of tests (Table 3). A strong antibacterial, antifungal and phytotoxic activities were also noted in the MeOH, EtOAc-soluble part of MeOH extract and 4 isolated sterols from *C. wallichii* against a variety of test organisms (Khaliq-uz-Zaman *et al.*, 2001). Functional derivatives of 2-(methylthio) propane-1,3-dithiol, extracted from *C. globularis*, were found to possess insecticidal properties (Jacobsen & Pedersen, 1983). Subsequently *C. aspera* was observed to accumulate benzo[a]pyrene, a carcinogenic compound in its thallus (Isha *et al.*, 1987).

The methanol extract of *N. flexilis* revealed the presence of 5 saturated and 14 unsaturated FAs (Table 2). The former were present in smaller amount (34.005 %) than the latter (61.82 %). Similar observations were made on *C. corallina* var. *wallichii* (Khaliq-uz-Zaman *et al.*, 1998) and several other freshwater green algae of Sindh (Ghazala *et al.*, 2004a-d). Tridecenoic acid (C13:1) was found to be in largest quantity (18.576 %) and undecanoic acid (C11:0) occurred in the smallest amount (0.167 %). The unsaturated acids included 7 mono-, 2 di-, 2 tri- and 3 poly-unsaturated FAs. The monounsaturated acids also included 2 monoyenoic acids (C8:1 and C14:1) having one triple bond. The polyunsaturated acids were having 4-5 double bonds. Such acids were not detected in the previous studies on *Chara* (Khaliq-uz-Zaman *et al.*, 1998).

From sterol and terpene fractions of the methanol extract of *N. flexilis* no natural product could be detected. A variety of sterols have been isolated previously from *C. corallina* var. *wallichii* (Khaliq-uz-Zaman *et al.*, 1998). The crude extract of *N. flexilis* showed a strong antifungal activity and a slight antibacterial activity. It did not display any significant cytotoxic and insecticidal activity (Table 3). It was observed that the

activity of thiol containing enzymes of *Nitella* is influenced by *p*-benzoquinone and pyrocatechol (Ivanova & Stom, 1980). *Nitella* spp., have the enzymatic machinery to decompose pyrocatechol, hydroquinone, phenol, *p*-cresol, guaiacol and other benzene derivatives (Stom *et al.*, 1979). The MeOH extract, its EtOAc-soluble part and four sterols isolated from *C. wallichii* showed bioactivity against 10 species of bacteria and 10 of fungi and its phytotoxicity against *Lemna minor* (Khaliq-uz-Zaman *et al.*, 2001).

References

- Aliya, R. and M. Shameel. 1999. Phycochemical evaluation of four coenocytic green seaweeds from the coast of Karachi. *Pak. J. Mar. Biol.*, 5: 65-76.
- Aliya, R. and M. Shameel. 2003. Marine natural products of *Caulerpa* (Siphonocladophyceae). *Pak. J. Bot.*, 35: 659-669.
- Aliya, R., M. Shameel, S. Perveen, M. S. Ali, K. Usmanghani and V. U. Ahmad. 1994. Acyclic diterpene alcohols isolated from four algae of Bryopsidophyceae and their toxicity. *Pak. J. Mar. Sci.*, 3: 15-24.
- Aliya, R., M. Shameel, K. Usmanghani, S. Sabiha and V.U. Ahmad. 1995. Fatty acid composition of two siphonaceous green algae from the coast of Karachi. *Pak. J. Pharm. Sci.*, 8: 47-54.
- Atta-ur-Rahman, M. A. Khan, M. Shabbir, M. Abid, M. I. Chaudhary, A. Nasreen, M. A. Maqbool, M. Shameel and R. Sualeh. 1997. Nematocidal study of marine organisms. *Pak. J. Nematol.*, 15: 95-100.
- Ghazala, B., M. Shameel, M.I. Choudhary, S. Shahzad and S.M. Leghari. 2003. Phycochemistry and bioactivity of certain freshwater green algae of Sindh. *Pak. J. Bot.*, 35:695-704.
- Ghazala, B., M. Shameel, M.I. Choudhary, S. Shahzad and S.M. Leghari. 2004a. Phycochemistry and bioactivity of *Tetraspora* (Volvocophyta) from Sindh. *Pak. J. Bot.*, 36: 511-527.
- Ghazala, B., M. Shameel, M.I. Choudhary, S. Shahzad and S.M. Leghari. 2004b. Phycochemistry and bioactivity of *Zgnema* (Zygnemaceae) from Sindh. *Int. J. Biol. Biotech.*, 1: 335-342.
- Ghazala, B., M. Shameel, M.I. Choudhary, S. Shahzad and S.M. Leghari. 2004c. Phycochemistry and bioactivity of two microalgae (Volvocophyta) from Sindh. *Int. J. Biol. Biotech.*, 1: 343-350.
- Ghazala, B., M. Shameel, M.I. Choudhary, S. Shahzad and S.M. Leghari. 2004d. Phycochemistry and bioactivity of three green algae in comparison with seaweeds of Sindh coast. *Pak. J. Mar. Sci.*, 13: (in press).
- Heilbron, I.M. 1942. Some aspects of algal chemistry. *J. Chem. Soc.*, 79: 78-89.
- Isha, P., U. Kirso and H. Kuk. 1987. Experimental investigation on the accumulation of carcinogens by macrophytes of Baltic Sea. *Eksp. Vodn. Toksikol.*, 12: 182-190.
- Ivanova, G.G. and D.I. Stom. 1980. Changes in the activity of thiol containing enzymes of aquatic plants under the effect of phenolic compounds. *Tsitologiya.*, 22:91-95.
- Jacobsen, N. and L.E.K. Pedersen. 1983. Synthesis and insecticidal properties of derivatives of propane-1,3-dithiol (analogs of the insecticidal derivatives of dithiolane and trithiane from the alga *Chara globularis* Thuillier). *Pestic. Sci.*, 14: 90-97.
- Khaliq-uz-Zaman, S.M., S. Shameel, M. Shameel, S.M. Leghari and V.U. Ahmad. 1998. Bioactive compounds in *Chara corallina* var. *wallichii* (A. Br.) R.D. Wood (Charophyta). *Pak. J. Bot.*, 30: 19-31.
- Khaliq-uz-Zaman, S.M., K. Simin and M. Shameel. 2001. Antimicrobial activity and phytotoxicity of sterols from *Chara wallichii* A. Br. (Charophyta). *Pak. J. Sci. Ind. Res.*, 44: 301-304.
- Naqvi, S.B. S., D. Sheikh, K. Usmanghani, M. Shameel and R. Sheikh. 1992. Screening of marine algae of Karachi for haemagglutinin activity. *Pak. J. Pharm. Sci.*, 5: 129-138.
- Parveen, Z. 1997. Use of potato disc assay for the screening of antitumor agents. In: Amtul, Z. (ed.): *Workshop on the Use of Bioassay Techniques for New Drug Development*. HEJ Res. Inst. Chem., Kar. Univ., p. 1-2.
- Patterson, G.W. 1972. Sterols of *Nitella flexilis* and *Chara vulgaris*. *Phytochem.*, 11: 3481-3483.

- Patterson, G.W., K.C. McKenna, W.R. Lusby and M.A. Bisson 1991. Sterols of the Charophyceae. *J. Nat. Prod.*, 54: 1141-1143.
- Qasim, R. 1986. Studies on fatty acid composition of eighteen species of seaweeds from the Karachi coast. *J. Chem. Soc. Pak.*, 8: 223-230.
- Rizvi, M.A. and M. Shameel. 2003. Biological activity and elementology of benthic algae from Karachi coast. *Pak. J. Bot.*, 35: 717-729.
- Saygideger, S. 1997: Bioaccumulation and toxicity of lead in the green alga *Spirogyra fluvialis* Hilse and *Zygnema pectinatum* (Zygnemataceae). *Turk. J. Biol.* 21: 343-352.
- Shameel, M. 1987. Studies on the fatty acids from seaweeds of Karachi. In: I. Ilahi and F. Hussain (Eds.): *Modern Trends of Plant Science Research in Pakistan. Proc. Nat. Conf. Plant Sci.*, 3: 183-186.
- Shameel, M. 1990. Phycochemical studies on fatty acids from certain seaweeds. *Bot. Mar.* 33: 429-432.
- Shameel, M. 1993. Phycochemical studies on fatty acid composition of twelve littoral green seaweeds of Karachi coast. In: *Proceedings of the National Seminar on Study and Management in Coastal Zones in Pakistan*. N. M. Tirmizi & Q. B. Kazmi (Eds.): Pak. Nat. Commis. UNESCO, Karachi, p. 17-25.
- Shameel, S., S.M. Khaliq-uz-Zaman, M. Shameel, S.M. Leghari and V.U. Ahmad. 1999. Phycochemical investigations on *Chara wallichii* A. Braun. In: M. Ahmad & G.H. Razwani (Eds.): *Herb Medicines and Therapeutics*. Dept. Pharmacog., Univ. Karachi., p. 156-174.
- Stom, D.I., S.S. Trinofeeva and L.I. Belykh. 1979. Decomposition of benzene and some phenols by aquatic plants. *Rast. Khim. Kantserogeny.*, 157-158.
- Usmanghani, K., M. Shameel, M. Sualeh, K. Khan and Z.A. Mahmood. 1984. Antibacterial and antifungal activities of marine algae from Karachi seashore of Pakistan. *Fitotrap.*, 55:73-77.
- Usmanghani, K. and M. Shameel. 1986. Studies on the antimicrobial activity of certain seaweeds from Karachi coast. In: R. Ahmad & A. San Pietro (Eds.): *Prospects for Biosaline Research*. Proc. US-Pak. Biosal. Res. Worksh., Karachi, p. 519-526.

(Received for publication 22 June 2004)