

ECO-MORPHIC RESPONSE TO SALT STRESS IN TWO HALOPHYTIC GRASSES FROM THE CHOLISTAN DESERT, PAKISTAN

NARGIS NAZ, MANSOOR HAMEED* AND MUHAMMAD ASHRAF†

Department of Botany, University of Agriculture, Faisalabad 38040, Pakistan

†Second affiliation: King Saud University, Riyadh, Saudi Arabia

*Corresponding author's E-mail: hameedmansoor@yahoo.com

Abstract

A study was conducted to evaluate the ecotypic difference in salinity tolerance in two halophytic grass species *Aeluropus lagopoides* and *Sporobolus ioclados* from the Cholistan Desert, Pakistan in a hydroponic system. Three populations of each of the two grass species were collected from ecologically different habitats, i.e., lesser salt-affected Derawar Fort (DF), moderately salt-affected Bailahwala Dahar (BD), and highly salt-affected Ladam Sir (LS). Genetic variation for salinity tolerance was found in all populations of both *A. lagopoides* and *S. ioclados*, its magnitude corresponded to the selection pressure of the habitats. The lesser saline DF populations of both grass species were the least tolerant to salt stress in relation to most of the agro-morphological characteristic measured. The moderately saline BD populations were relatively less affected due to salt stress and 100 mM NaCl stimulated their growth measured in terms of plant height, root length and fresh and dry weights of shoots and root. The highly saline LS populations were the most tolerant among all populations of both grasses and optimal growth was recorded at 200 and 300 mM NaCl of the growth medium. Of the two grasses, *A. lagopoides* was superior to *S. ioclados* in salinity tolerance because its all three populations were relatively more tolerant as compared to those of the latter grass species.

Introduction

The importance of variation among ecotypes within a plant species has been documented since long (Bradshaw, 1984; Kuiper, 1990; Ashraf & Ahmad, 1995; Ashraf *et al.*, 2003; Hameed *et al.*, 2008; Naz *et al.*, 2009a). Such variation is of vital important in providing suitable genetic stock capable of coping with a multitude of environmental conditions (Pezeshki & DeLaune, 1997). Generally, the genetic heterogeneity among populations of a particular species is associated with evolutionary processes like interactions among natural selection, selection pressure, genetic drift and recombination, and mutation (Ashraf, 2004; Fjellheim & Rognli, 2005). Selection pressure of variable habitats during an evolutionary history of a species can evolve into quite diverse populations, which are genetically distinct and adapted to such specialized habitats (Eleuterius, 1989).

Genetically based variation has been reported in many salt tolerant and halophytic species e.g., in *Imperata cylindrica* (Cheng & Chou, 1997; Hameed *et al.*, 2009), *Cenchrus ciliaris* (Hameed *et al.*, 2002), *Cynodon dactylon* (Hameed & Ashraf, 2008), *Aeluropus lagopoides* and *Ochthochloa compressa* (Naz *et al.*, 2009b), *Dactylis spicata* (Brewer & Bertness, 1996), *Panicum antidotale* (Hameed & Ashraf, 2008), and *Spartina alterniflora* and *Spartina patens* (Seliskar, 1995).

Saline flats of the Cholistan desert support a number of halophytic or salt tolerant plant species, among including two considerably important grasses *Aeluropus lagopoides*

and *Sporobolus ioclados*. However, these two grasses were selected for the present study. *Aeluropus* is a genus of 6 species and represented by 3 species in Pakistan, whereas *Sporobolus* includes 250 species and represented by 10 species in Pakistan (Cope, 1982; Chaudhry, 1989). *Aeluropus littoralis* can tolerate up to 1,100 mM NaCl and *A. lagopoides* 1,500 mM NaCl (Bodla *et al.*, 1995; Gulzar *et al.*, 2003a&b). Many species in genus *Sporobolus* are highly salt tolerant and halophytic (Ashour & Mekki, 2006). Examples are *S. virginicus* (Ashour *et al.*, 1997; Bell & O'Leary, 2003), *S. airoides* (Marcum, 1999; Grieve *et al.*, 2004), *S. ioclados* (Gulzar *et al.*, 2005), *S. madraspatanus* (Joshi *et al.*, 2005), *S. spicatus* (Ramadan, 2001), *S. pyramidatus* (Tóth *et al.*, 1997), and *S. arabicus* (Khan & Ungar, 2001).

Identification of natural populations tolerant to high salinities is extremely vital in developing strategies for re-vegetation and reclamation of salt-affected areas. Salt tolerant ecotypes can provide exceptional means for investigating the adaptative mechanisms to tolerate environmental stresses (Ashraf, 2003), and therefore, can be used as a model plants (Hameed *et al.*, 2008; Pessarakli & Kopec, 2009). The present study was focused on genetic variations among the ecologically different populations and the morphological response to salt stress in two grasses (*A. lagopoides* and *S. ioclados*) to elevated levels of salinity in a hydroponic culture solution under the Faisalabad environmental conditions.

Materials and Methods

Ecotypic difference in salinity tolerance of three populations was investigated of each of each of two halophytic grass species *Aeluropus lagopoides* (L.) Trin. ex Thw., and *Sporobolus ioclados* (Nees ex. Trin.) Nees from the Cholistan Desert, Pakistan. The experiment was conducted hydroponically in half-strength Hoagland's nutrient culture solution in the Botanical Garden, University of Agriculture, Faisalabad. Three populations of each of the two grass species were collected in the Cholistan Desert from ecologically different habitats, i.e., lesser salt-affected Derawar Fort (DF; coordinates 71° 27' 32.73" E, 29° 24' 31.65" N; ECe 15.21 dS m⁻¹; Na⁺ 3246.46 mg L⁻¹; Cl⁻ 1483.21 mg L⁻¹), moderately salt-affected Bailahwala Dahar (BD; coordinates 70° 93' 23.34" E, 29° 38' 18.27" N; ECe 26.7 dS m⁻¹; Na⁺ 4382.45 mg L⁻¹; Cl⁻ 2446.62 mg L⁻¹), and highly salt-affected Ladam Sir (LS; coordinates 72° 64' 25.11" E, 30° 53' 26.42" N; ECe 49.28 dS m⁻¹; Na⁺ 5149.33 mg L⁻¹; Cl⁻ 2647.76 mg L⁻¹). Electrical conductivity of saturated soil extract (ECe) of the rhizospheric soil of all the populations of these grasses was measured using a electrical conductivity meter (WTW series InoLab pH/Cond 720), soil Na⁺ by a flame photometer (Model Sherwood 410, USA) and soil Cl⁻ by a chloride meter (Model 926, Sherwood Scientific Ltd., Cambridge).

Populations of both grasses from all three habitats were established in loamy-clay soil. Ramets were detached from each population and grown in half-strength Hoagland's nutrient solution (Hoagland & Arnon, 1950). Air pumps were used for about 12 h daily for the aeration of the hydroponic system.

The experiment was conducted in a 3-factor completely randomized design with 3 populations (DF, BD, and LS), 2 grasses (*A. lagopoides* and *S. ioclados*), and 4 salinity levels (0, 100, 200 and 300 mM NaCl) with 10 replications. Agro-morphological characteristics studied during the experiment were shoot and root length, top internodal length, number of tillers and leaves per plant, and fresh and dry weights of plant.

Results

All populations of both *Aeluropus lagopoides* and *Sporobolus ioclados* showed differential response under increasing salt levels. Shoot length in population of *A. lagopoides* from the least saline Derawar Fort (DF) consistently decreased with increase in salt level of the medium (Fig. 1). In contrast, the populations from moderately saline Bailahwala Dahar (BD) and highly saline Ladam Sir (LS) showed a slight increase up to 200 mM NaCl of external medium and thereafter a slight decrease was observed at 300 mM NaCl. Shoot length in all populations of *S. ioclados* were only affected by the highest salt level (300 mM NaCl), however, the population from the highest saline LS showed a consistent increase in shoot length up to 200 mM NaCl.

A similar response of root length was recorded in all populations of *A. lagopoides*, which depicted a gradual decrease with increasing salt levels of the hydroponic culture solution. The highly saline LS population, however, was relatively less affected in terms of root length as compared to the other populations by increasing salt levels. The *S. ioclados* population from lesser saline DF showed a considerable decrease in root length, while those from moderately saline BD and highly saline LS showed increased root length under elevated salt levels.

Top internodal length showed quite a variable behaviour under salt stress (Fig. 1). It was promoted by moderate salt level (100 mM NaCl) in all three populations of *S. ioclados*. Top internodal length in *A. lagopoides*, nevertheless, in the DF population consistently decreased, in BD population it increased up to 200 mM salt level, and in LS population, it consistently increased with increase in salt levels.

Number of tillers per plant and number of leaves per plant showed almost a similar response to elevated salt levels. The DF population of both *A. lagopoides* and *S. ioclados* were adversely affected in relation to these parameters by increasing salt levels. In contrast, BD population showed stability in these parameters and LS populations showed an increase in number of tillers and leaves with increase in salt levels.

Total leaf area was negatively affected in the populations of both grasses from lesser saline LS at higher salt levels. The leaf area of LS population of *A. lagopoides* was consistently increased with increase in salt levels, while that of the LS population of *S. ioclados* was slightly affected at the highest 300 mM NaCl level (Fig. 1).

A similar response in terms of root fresh and dry weights was recorded in all populations of both grass species (Fig. 2). *Aeluropus lagopoides* from DF showed the maximum of these parameters at 100 mM NaCl, whereas that from BD showed the maximum value at 200 mM and that from LS at 300 mM NaCl. Fresh and dry weights of DF and BD populations of *S. ioclados*, on the other hand, were slightly increased at 100 mM NaCl, and these parameters decreased gradually with further increase in the salt levels. The LS population, however, was adversely affected only at 300 mM NaCl in relation to these parameters.

A significant decrease was recorded in shoot fresh and dry weight in the DF population of both *A. lagopoides* and *S. ioclados* by increasing salt levels. These parameters were adversely affected at 300 mM NaCl in the BD populations of both grasses, whereas, in the LS population, these parameters were consistently increased with increase in salt levels (Fig. 2).




Fig. 1. Some agro-morphological characteristics recorded in ecologically different populations of two grass species under salt stress ($n = 10$, means $\pm$ SE, DF = Derawar Fort, BD = Bailahwala Dahar, LS = Ladam Sir).
0, 100, 200 and 300 mM NaCl.




Fig. 2. Fresh and dry weights of roots and shoots recorded in ecologically different populations of two grass species under salt stress ($n = 10$, means $\pm$ SE, DF = Derawar Fort, BD = Bailahwala Dahar, LS = Ladam Sir).

0, 100, 200 and 300 mM NaCl.

Discussion

Cholistan desert of Pakistan is a unique biome subjected to an interplay of several environmental stresses including drought, salinity, temperature and nutrient stresses (Arshad *et al.*, 2008). Environmental heterogeneity and long evolutionary history of this desert is enough to impose significant genetic variations within the native species, making them perfectly adaptative to the harsh environmental conditions of the desert (Weber, 2009). Inter-dunal saline patches in the Cholistan desert have a significant amount of variation in soil physico-chemical characteristics (Naz *et al.*, 2010), and this results in differential response of different ecotypes of the inhabiting species in diverse habitats. Such genetic variability within individual plant species has been reported by several authors in response to environmental variability (Toth *et al.*, 1997; Ramadan, 2001; Bell & O'Leary, 2003; Hameed *et al.*, 2009).

Genetic variation was found in all populations of both *A. lagopoides* and *S. ioclados*, which was the basis of their differential response to high salt stress (Naz *et al.*, 2009b; Hameed *et al.*, 2009). The lesser saline DF populations of both grass species were the least tolerant to salt stress in relation to most of the agro-morphological characteristics. All agro-morphological characteristics of both grasses showed a significant decrease due to increasing salinity. Such decrease in less tolerant ecotypes due to salt stress has been reported by several authors (Ashraf & Ahmad, 1995; Ashraf *et al.*, 2006; Hameed & Ashraf, 2008).

The moderately saline BD populations were relatively less affected by increasing salinity. However, moderate salt level (100 mM NaCl) resulted in increased growth in terms of plant height, root length and fresh and dry weights of shoots and root. There are only few reports on salt-induced growth stimulation in monocots, especially in grasses (Munns & Tester, 2008; Flowers & Colmer, 2008). However, optimal growth of salt tolerant or halophytic grasses has been reported by several authors at moderate salinities such as in *Puccinellia peisonis* (Stelzer & L  uchli, 1977), *Leymus arenarius* (Greipsson & Davy, 1996), and *Imperata cylindrica* (Hameed *et al.*, 2009).

The highly saline LS populations were the most tolerant among all populations of both grasses with respect to agro-morphological characteristics measured in the present study. Optimal growth of the LS population of both grasses was recorded at high salinity levels, mainly at 200 mM NaCl of the growth medium. In particular, the LS population of *A. lagopoides* showed the maximum growth in terms of fresh and dry biomass at the highest salt level i.e., 300 mM NaCl. Similar results have been reported by Marcum & Murdoch (1994) in *Stenotaphrum secundatum* and Naz *et al.*, (2009b) in *Ochthochloa compressa* at high salinities.

Ecological factors strongly affect the genetic variation within populations of specific species (Kolliker *et al.*, 1998; Hsao & Lee, 1999), which in turn, is responsible for the development of an ecotype adapted to such specific environments (Gunter *et al.*, 1996). From the present study it was evident that the variation in both grasses due to habitat differences is genetically based and this may be the basic requirement for the improvement of salinity tolerance in other plant species through conventional breeding and advanced molecular/genetic engineering techniques (Al-Khatib *et al.*, 1993; Munns & Tester, 2008). Varying degree of salinity tolerance in a variety of species due to ecotypic differences has been reported by several researchers e.g., in *Lolium multiflorum* (Hannaway *et al.*, 1999), *Leymus arenarius* (Greipsson & Davy, 1996), *Cymbopogon*

jawarancusa (Arshad *et al.*, 1995), *Sporobolus ioclados* (Arshad *et al.*, 1999) and *Cynodon dactylon* (Hameed & Ashraf, 2008).

Overall all populations (ecotypes) of both *A. lagopoides* and *S. ioclados* from highly saline LS were more tolerant as compared to those from moderately saline BD and lesser saline DF, as they showed optimal growth at moderate to high salinities in terms of various growth parameters studied.

Acknowledgement

This work is a part of Ph.D. thesis of Miss Nargis Naz, Regd. No. 2003-ag-468.

References

- Al-Khatib, M.M., T. McNeilly and J.C. Collins. 1993. The potential of selection and breeding for improved salt tolerance in Lucerne (*Medicago sativa* L.). *Euphytica*, 65: 43-51.
- Arshad, M., A.R. Rao, G. Akbar and M. Akhtar. 1999. Eco-physiological adaptability potential of *Sporobolus ioclados* in Cholistan Desert. *Pak. J. Biol. Sci.*, 2: 1604-1607.
- Arshad, M., A.R. Rao, M. Akhtar and G. Akbar. 1995. Potential genetic adaptability of *Cymbopogon jwarancusa* (Jones) Schult, from Cholistan Desert, Pakistan. *Sarhad J. Agri.*, 11: 63-69.
- Arshad, M., Anwar-ul-Hussan, M.Y. Ashraf, S. Noureen and M. Moazzam. 2008. Edaphic factors and distribution of vegetation in the Cholistan Desert, Pakistan. *Pak. J. Bot.*, 40: 1923-1931.
- Ashour, N.I. and B.B. Mekki. 2006. Halophytic forage plants in response to irrigation with diluted seawater. *World J. Agric. Sci.*, 2: 169-173.
- Ashour, N.I., M.S. Serage, A.K. Abd El-Halim and B.B. Mekki. 1997. Forage production from three grass species under saline irrigation in Egypt. *J. Arid. Environ.*, 37: 299-307.
- Ashraf, M. 2003. Relationships between leaf gas exchange characteristics and growth of differently adapted populations of Blue panicgrass (*Panicum antidotale* Retz.) under salinity or waterlogging. *Plant Sci.*, 165: 69-75.
- Ashraf, M. 2004. Some important physiological selection criteria for salt tolerance in plants. *Flora*, 199: 361-376.
- Ashraf, M. and H. Ahmad. 1995. Response of three arid zone grasses to salt and waterlogging. *Arid Soil Res. Rehabil.*, 9: 137-154.
- Ashraf, M., M. Hameed, M. Arshad, Y. Ashraf and K. Akhtar. 2006. Salt tolerance of some potential forage grasses from Cholistan Desert of Pakistan. In: *Ecophysiology of High Salinity Tolerant Plants*. (Eds.): M.A. Khan and D.J. Weber. Springer, Netherlands.
- Ashraf, M., R. Zafar and M.Y. Ashraf. 2003. Time-course changes in the inorganic and organic components of germinating sunflower achenes under salt (NaCl) stress. *Flora*, 198: 26-36.
- Bell, H.L. and J.W. O'Leary. 2003. Effects of salinity on growth and cation accumulation of *Sporobolus virginicus* (Poaceae). *Am. J. Bot.*, 90: 1416-1424.
- Bodla, M.A., M.R. Chaudhry, S.R.A. Shamsi and M.S. Baig. 1995. Salt tolerance in some dominant grasses of Punjab. In: *Biology of Salt Tolerant Plants*. (Eds.): M.A. Khan and I.A. Ungar. Book Crafters, Michigan, USA, p. 190-198.
- Bradshaw, A.D. 1984. The importance of evolutionary ideas in ecology and vice versa. In: *Evolutionary Ecology*. (Ed.): B. Sharrocks. Blackwell, Oxford, UK, pp: 1-25.
- Brewer, S.D. and M.D. Bertness. 1996. Natural disturbance and the genetic structure of marsh plant populations. *Oikos*, 77: 107-116.
- Chaudhary, S.A. 1989. Grasses of Saudi Arabia, National Herbarium, National Agriculture and Water Research Center, Ministry of Agriculture and Water, Riyadh, Kingdom of Saudi Arabia.
- Cheng, K.T. and C.H. Chou. 1997. Ecotypic variation of *Imperata cylindrica* populations in Taiwan: I. Morphological and molecular evidences. *Bot. Bull. Acad Sinica*, 38: 215-223.
- Cope, T.A. 1982. Poaceae, No. 143. In: *Flora of Pakistan*. (Eds.): E. Nasir and S.I. Ali. National Herbarium, Pakistan Agricultural Research Council, Islamabad.

- Eleuterius, L.N. 1989. Natural selection and genetic adaptation to hypersalinity in *Juncus roemerianus* Scheele. *Aquatic Bot.*, 36: 45-53.
- Fjellheim, S. and O.A. Rognli. 2005. Genetic diversity within and among Nordic meadow fescue (*Festuca pratensis* Huds.) cultivars determined on the basis of AFLP markers. *Crop Sci.*, 45: 2081-2086.
- Flowers, T.J. and T.D. Colmer. 2008. Salinity tolerance in halophytes. *New Phytol.*, 179: 945-963
- Greipsson, S. and A.J. Davy. 1996. Sand accretion and salinity as constraints on the establishment of *Leymus arenarius* for land reclamation in Iceland. *Ann. Bot.*, 78: 611-618.
- Grieve, C.M., J.A. Poss, S.R. Grattan, D.L. Suarez, S.E. Benes and P.H. Robinson. 2004. Evaluation of salt-tolerant forages for sequential water reuse systems. II. Plant ion relations. *Agric. Water Manag.*, 70: 121-135.
- Gulzar, S., M.A. Khan and I.A. Ungar. 2003a. Effects of salinity on growth, ionic content, and plant-water status of *Aeluropus lagopoides*. *Comm. Soil Sci. Plant Anal.*, 34: 1657-1668.
- Gulzar, S., M.A. Khan and I.A. Ungar. 2003b. Salt tolerance of a coastal salt marsh grass. *Comm. Soil Sci. Plant Anal.*, 34: 2595-2605.
- Gulzar, S., M.A. Khan, I.A. Ungar and X. Liu. 2005. Influence of salinity on growth and osmotic relations of *Sporobolus ioclados*. *Pak. J. Bot.*, 37: 119-129.
- Gunter, L.E., G.A. Tuskan and S.D. Wulschler. 1996. Diversity of populations of switchgrass based on RAPD markers. *Crop Sci.*, 36: 1017-1022.
- Hameed, M. and M. Ashraf. 2008. Physiological and biochemical adaptations of *Cynodon dactylon* (L.) Pers. from the Salt Range (Pakistan) to salinity stress. *Flora*, 203: 683-694.
- Hameed, M., A.A. Chaudhry, M.A. Man and A.H. Gill. 2002. Diversity of plant species in Lal Suhanra National Park, Bahawalpur, Pakistan. *J. Biol. Sci.*, 2: 267-274.
- Hameed, M., M. Ashraf and N. Naz. 2009. Anatomical adaptations to salinity in cogon grass [*Imperata cylindrica* (L.) Raeuschel] from the Salt Range, Pakistan. *Plant Soil*, 322: 229-238.
- Hameed, M., N. Naz, M.S.A. Ahmad, Islam-ud-Din and A. Riaz. 2008. Morphological adaptations of some grasses from the salt range, Pakistan. *Pak. J. Bot.*, 40: 1571-1578.
- Hannaway, D.B., S. Fransen and J. Cropper. 1999. Tall fescue. PNW Ext. Ser. Circ. 504. Oregon State Univ., Corvallis, OR.
- Hoagland D.R. and D.I. Arnon. 1950. The water culture method for growing plants without soil. In: Circular No. 347, Univ. Calif. Agric. Exp. Stn., Berkeley, CA, pp. 1-39.
- Hsao, J.Y. and M. Lee, 1999. Genetic diversity and microgeographic differentiation of Yushan cane (*Yushina nittakayamensis*; Poaceae) in Taiwan. *Mol. Ecol.*, 8: 263-270.
- Joshi, A.J., B.S. Mali and H. Harsha. 2005. Salt tolerance and germination and early growth of two forage grasses growing in marshy habitats. *Environ. Exp. Bot.*, 54: 267-274.
- Khan, M.A. and I.A. Ungar. 2001. Alleviation of salinity stress and the response to temperature in two seed morphs of *Halopyrum mucronatum* (Poaceae). *Aust. J. Bot.*, 49: 777-783.
- Kolliker, R., F.J. Stadelmann, B. Reidy and J. Nosberger. 1998. Fertilization and defoliation frequency affect genetic diversity of *Festuca pratensis* Huds. In permanent grasslands. *Mol. Ecol.*, 7: 1557-1567.
- Kuiper, P.J.C. 1990. Analysis of phenotypic responses of plants to changes in the environment in terms of stress and adaptation. *Acta Bot. Neerl.*, 39: 217-227.
- Marcum, K.B. 1999. Salinity tolerance mechanisms of grasses in the subfamily Chloridoideae. *Crop Sci.*, 39: 1153-1160.
- Marcum, K.B. and C.L. Murdoch. 1994. Salinity tolerance mechanisms of six C4 turfgrasses. *J. Am. Soc. Hortic. Sci.*, 119: 779-784.
- Munns, R. and M. Tester. 2008. Mechanisms of salinity tolerance. *Annu. Rev. Plant Biol.*, 59: 651-681.
- Naz, N., M. Hameed, A. Wahid, M. Arshad and M.S.A. Ahmad. 2009a. Patterns of ion excretion and survival in two stoloniferous arid zone grasses. *Physiol. Plant.*, 135: 185-195.
- Naz, N., M. Hameed, M. Ashraf, R. Ahmad and M. Arshad. 2009b. Eco-morphic variation for salt tolerance in some grasses from Cholistan Desert, Pakistan. *Pak. J. Bot.*, 41: 1707-1714.

- Naz, N., M. Hameed, M.S.A. Ahmad, M. Ashraf and M. Arshad. 2010. Is soil salinity one of the major determinants of community structure under arid environments? *Commun. Ecol.*, (in press).
- Pessarakli, M. and D.M. Kopec. 2009. Screening various ryegrass cultivars for salt stress tolerance. *J. Food Agri. Environ.*, 7: 739-743.
- Pezeshki, S.R. and R.D. DeLaune. 1997. Population differentiation in *Spartina patens*: Responses of photosynthesis and biomass partitioning to elevated salinity. *Bot. Bull. Acad. Sin.*, 115-120.
- Ramadan, T. 2001. Dynamics of salt secretion by *Sporobolus spicatus* (Vahl) Kunth from sites of differing salinity. *Ann. Bot.*, 87: 259-266.
- Seliskar, D.M. 1995 Exploiting plant genotypic diversity for coastal salt marsh creation and restoration. In: *Biology of salt-tolerant plants*. (Eds.): M.A. Khan and I.A. Ungar. Department of Botany, University of Karachi, Karachi, Pakistan, pp. 407-416.
- Stelzer, R. and A. Lauchli. 1977. Salt- and flooding tolerance of *Puccinellia peisonis*: I. The effect of NaCl- and KCl-salinity on growth at varied oxygen supply to the root. *Z. Pflanzenphysiol*, 83: 35-42.
- Tóth, T., M. Kertesz, L.C. Guerrat, J.L. Labrada, B.P. Machado, P.C. Fonseca and M.N. Martinez. 1997. Plant composition of a pasture as a predictor of soil salinity. *Rev. Biol. Trop.*, 45: 1385-1393.
- Weber, D.J. 2009. Adaptive mechanisms of halophytes in desert regions. In: *Salinity and water stress*. (Eds.): M. Ashraf, M. Ozturk and H.R. Athar. Springer-Verlag, pp: 179-186.

(Received for publication 14 September 2009)