

## SALINITY TOLERANCE OF THREE RANGE GRASSES AT GERMINATION AND EARLY GROWTH STAGES

PARVEEN AKHTAR<sup>1</sup> AND FARRUKH HUSSAIN<sup>2</sup>

<sup>1</sup>Jinnah College for Women, University of Peshawar

<sup>2</sup>Department of Botany, University of Peshawar, Peshawar, Pakistan.

### Abstract

Three range grasses viz., *Bothriochloa pertusa* (L.) A. Camus, *Dichanthium annulatum* Forssk and *Panicum antidotale* Retz were subjected to 5, 10 and 15 dSm<sup>-1</sup> and control (0.21 dSm<sup>-1</sup>) salinity levels in the laboratory. Germination percent, early seedling growth rate and moisture contents were determined. Salinity reduced germination in *Panicum antidotale* more than *Bothriochloa pertusa* and *Dichanthium annulatum* at all levels of applied salinity. The fresh weight of all the three grasses significantly declined specially in high concentration. The reduction in dry weight was obvious only at 15dSm<sup>-1</sup> in all the tested grasses.

### Introduction

Pakistan is the ninth populous country in the world with a population of more than 160 million. Unfortunately about 6.5 million hectare of land in Pakistan has got different degree of salinity and alkalinity hazard (Alam, 2002). The need for fodder and shelter for increasing population demands the use of unproductive salted land through the cultivation of salt tolerant plants as one of the possible approaches. Utilizing saline habitat for the production of fodder crops, besides providing vegetation cover, might also improve the health, milk and meat production of livestock. Studies made on the salinity tolerance of fodder plants have shown differential response to varying degrees of salinity (Jan *et al.*, 2000, Badger & Unger, 1998, White & Gartner, 1997; Tobe *et al.*, 2002; Li, 2007). Song *et al.*, (2006) worked on the effect of salinity and temperature on three salt resistant euhalophytes. Qu *et al.*, (2007) investigated the effects of salinity on germination and radicle growth of forage plants. It has been observed that salinity affects the germination and early seedling growth of susceptible species (Munns & James, 2003; Hanselin & Eggen, 2005; Khan & Gul, 2006). Sabir & Ashraf (2007) screened *Panicum maliaceum* accession for salinity tolerance at seedling stage. Similarly, Zehra & Khan (2007) worked on the effect of NaCl salinity on germination of *Phragmites*. As no such information is available on these three range grasses therefore the present study was conducted to see the tolerance of these three range grasses at germination and early seedling growth stage. The findings would help in screening range grasses suitable for cultivation on salted land for improving the socio economic uplift of people depending upon the rangelands.

### Materials and Methods

Germination and seedling growth studies were conducted under Laboratory conditions. Certified seeds of *Bothriochloa pertusa* (L.) A. Camus, *Dichanthium annulatum* Forssk and *Panicum antidotale* Retz were obtained from the Range Management Branch, Pakistan Forest Institute, Peshawar. Viability of the seeds was tested by tetrazolium test. The glassware was thoroughly washed and sterilized at 170°C for 4 hours prior to use. Ten viable seeds without covering of the above mentioned grasses were sown in 5, 10 and 15 dSm<sup>-1</sup> NaCl solution along with distilled water control (0.21 dSm<sup>-1</sup>) on twice folded Whatman No. 1 filter paper per seedbeds in Petri dishes. Each treatment was replicated 10

times. The Petri dishes were incubated at  $25\pm 1^{\circ}\text{C}$ . Germination percentage was recorded daily for upto 120 hours. Fresh weights of the 20 randomly selected seedlings from each treatment were determined and then dried at  $65\pm 1^{\circ}\text{C}$  for 72 hours to get dry weights. The moisture contents of the seedlings were determined on oven dry weight basis. The results were subjected to ANOVA (Steel & Torie, 1980).

### Results and Discussion

**Effect on % germination:** The germination of all the three tested grasses significantly declined even at  $5\text{ dSm}^{-1}$  level of treatment that further decreased at  $10\text{ dSm}^{-1}$  and  $15\text{ dSm}^{-1}$  (Table 1). *Panicum* was inhibited more than *Bothriochloa* and *Dichanthium* at all levels of applied salinity. Reduction in germination is in conformity with the reports of other workers (El-Nathlawy & El-Fawal, 1989; Hanselin & Eggen, 2005; Sabir & Ashraf, 2007; Khan & Gul, 2006), who also found reduced germination due to salinity. Priano & Platti (1989) also observed that *Chloris gayana*, *Elytrigia elongata* and *Elytrigia scabrifolia* seeds had reduced germination under saline condition. Similar findings regarding reduced germination of other species under salinity stress have also been reported by Shadded & Zaidan (1989), Hussain *et al.*, (1993); Ashraf *et al.*, (1989) and Zehra & Khan (2007).

**Effect on seedling growth:** Salinity affects both the establishment and growth of seedlings by retarding plumule and radicle growth. In the present case, except *Dichanthium* at  $5\text{ dSm}^{-1}$  salinity severely reduced the plumule and radicle growth of all the tested grasses in all the treatments especially at  $15\text{ dSm}^{-1}$  (Table 2). The findings are in conformity with other studies (Ali *et al.*, 1992; Ilahi & Begum, 1981; Lyra *et al.*, 1992; Zehra & Khan, 2007) who also observed decreased germination percentage and seedling growth with increasing salinity.

**Effect on biomass and moisture contents:** There has been considerable variation in the behavior of biomass and moisture contents under test condition among the grasses. However, it is seen that dry mass generally reduces while fresh biomass may increase or decrease under saline condition. The fresh weight of all the three grasses significantly declined especially at  $10\text{ dSm}^{-1}$  (Table 3), while the reduction in dry weight was obvious only at  $15\text{ dSm}^{-1}$  in all the tested grasses. The present findings are in conformity with those of Lyra *et al.*, (1992) who observed poor germination percentage, fresh weight and dry mass in *Sesamum* under salinity. Naqvi & Mahmood (1994) also reported that fresh dry weight and moisture contents of *Sesbania sesban* seedlings decreased with increasing levels of NaCl salinity. Likewise, Shadded & Zaidan (1989) concluded that NaCl salinity reduced the germination and seedling growth of *Raphanus sativus* and *Trigonella foenum-graecum*. The moisture contents showed inconsistent increase or decrease under saline conditions in the present study. The moisture contents of *Bothriochloa* in all the test condition and that of *Panicum* decreased at  $10\text{ dSm}^{-1}$  only. The reduction in moisture contents agrees with Shadded & Zaidan (1989) who reported reduced seedlings moisture contents under saline conditions. In most cases seedling exhibited higher moisture contents than control. This might be an adaptive mechanism to overcome salinity stress by becoming succulent by retaining more moisture. The present study suggests that the three tested grasses might germinate and survive under mild salinity levels on marginal salted soils. However, further study is needed in the field to see their germination and growth behavior in soil. The preliminary study just points at the possibility of further testing these grasses under more natural condition.

**Table 1. Effect of various levels of NaCl (dSm<sup>-1</sup>) salinization on the average germination % of three grasses under laboratory condition. Each value is a mean of 10 replicates, each with 10 seeds.**

Species	Control	Salinity (dSm <sup>-1</sup> ) levels						Average
		5	% of control	10	% of control	15	% of control	
<i>Bothriochloa pertusa</i>	65	41	63.08	31	47.69	15	23.08	29.00
<i>Dichanthium annulatum</i>	60	48	80.00	31	51.67	13	21.67	30.66
<i>Panicum antidotale</i>	92	41	44.56	37	40.22	17	18.48	31.66

**Table 2. Effect of various levels of NaCl (dSm<sup>-1</sup>) salinization on the radicle and plumule length in three grasses under laboratory condition. Each value is a mean of 10 replicates, each with 10 seeds**

Species	Control	Salinity (dSm <sup>-1</sup> ) levels					
		5	% of Control	10	% of Control	15	% of Control
<b>Radicle Growth (mm)</b>							
<i>Bothriochloa pertusa</i>	35	31	88.57	26	74.29	19	54.29
<i>Dichanthium annulatum</i>	26	24	92.31	17	65.38	14	53.85
<i>Panicum antidotale</i>	35	19	54.29	11	31.43	5	14.29
<b>Plumule Growth (mm)</b>							
<i>Bothriochloa pertusa</i>	33	26	78.79	19	57.58	10	30.30
<i>Dichanthium annulatum</i>	21	18	85.71	10	47.62	8	30.10
<i>Panicum antidotale</i>	36	25	69.44	9	25.00	8	22.22

**Table 3. Effect of various levels of NaCl (dSm<sup>-1</sup>) salinization on the fresh weight, dry weight (mg) and % moisture of seedlings of three grasses.**

Salinity dSm <sup>-1</sup>	Biomass (mg)					
	Fresh weight (mg)		Dry weight (mg)		% Moisture	
	Test	% of Control	Test	% of Control	Test	% of Control
<i>Bothriochloa pertusa</i>						
Control	45.50 a	100.00	19.98 a	100	127.73 a	100
5	39.90 b	87.69	19.00 a	95.10	110.00 b	86.12
10	28.90 c	62.96	18.99 a	95.05	52.18 c	40.85
15	19.90 d	43.74	10.00 b	50.05	99.00 b	77.51
<i>Dichanthium annulatum</i>						
Control	53.6 a	100	19.00 a	10	182.11 a	100
5	46.5 b	86.75	15.50 b	81.58	200.00 a	109.82
10	35.0 c	65.30	10.00 c	52.63	250 b	137.28
15	29.90 d	55.78	9.90 c	52.10	202.02 a	110.93
<i>Panicum antidotale</i>						
Control	47.2 a	100	11.00 a	100	229.89 a	100
5	37.70 b	79.87	10.00 a	90.91	277.00 b	120.49
10	21.00 c	44.49	10.00 a	90.91	110.00 c	47.85
15	20.0 c	42.37	5.00 b	45.45	300.00 d	130.50

Means followed by the same letter within a column for the species are not significantly different from each other.

## References

- Alam, S.M. 2002. Salinity poses distinct physiological threats to plants utilization of saline lands. Environment. The Daily Pakistan and Gulf Economist.
- Ali, R.M., Z.A. Ahmad and R. Abde-Basset. 1992. Effect of NaCl salinity on seed germination and endogenous levels of some metabolites and respiration of *Trigonella foenum-graecum* L., radicles. *Bull. Faculty Agric. Univ. Cairo*, 43(2): 817-828.
- Ashraf, M., T. McNeilly and A.D. Bradshaw. 1989. The potential for evaluation of tolerance to Sodium chloride, Calcium chloride, Magnesium chloride and seawater in four grass species. *New Phytologist*, 112(2): 245-254.
- Badger, K.S. and I. Unger. 1998. The effect of salinity on the germination of inland halophytes. *Can. J. Bot.*, 67(4): 442-445.
- El-Nathlawy, F.S. and M.A. El-Fawal. 1989. Tolerance of five oil crops to salinity and temperature stresses during germination. *Acta Agronomica Hungarica*, 38 (1-2): 59-65.
- Hanselin, M.H and T. Eggen. 2005. Salinity tolerance during germination of seashore halophytes and salt tolerance grass cultivars. *Seed Sci. Res.*, 15: 43-50
- Hussain, F., K. Janus and I. Ihsan. 1993. Effects of induced salinity on the germination and seedling growth of two cultivars of *Cicer arietinum*. *Sarhad J. Agric.*, 13(2): 33-39.
- Ilahi, I. and M. Begum. 1981. The effects of Sodium chloride on imbibition, germination, growth and development of *Helianthus annuus* L. *J. Sci. Tech.*, 5(2): 126-131.
- Jan, K.A., R. Koenig and K. Boyed. 2000. Tolerance of some turf Grasses. Electronic Publishing, *Utah State University Extension*.
- Khan, M.A. and B. Gul. 2006. Halophytic seed germination. In: *Ecophysiology of high salinity tolerant plants*. (Eds.): M.A. Khan & D.J. Weber. *Springler Verloge Netherlands*, pp. 11-30.
- Li, L. and X. Zhang. 2007. Effect of temperature on germination of two *kalidium* species. *Chin. J. Appl. Environ. Biol.*, 13: 317-321.
- Lyra, J.R. M.E. Araugo, J.P. Dantas, V. de P. Queiroga and Ride La. Bruno. 1992. Quality of sesame (*Sesemum indicum* L.) seeds produced under conditions of salt stress. *Revista Brasileira de Sementes*, 14(2): 201-206.
- Munnz, R. and R.A. James. 2003. Screening methods for salt tolerance. A case study with tetraploid wheat. *Plant Soil*, 253: 239-250
- Naqvi, S.A.R. and A. Mahmood. 1994. Effect of NaCl salinity on growth, nodulation and nitrogen fixation in *Sesbania sesban* L. *International. Symp. on High Salinity Tolerant Plants. Department of Botany, University of Karachi*. December 12-16. pp. 69.
- Priano, L.J. and M.A. Platti. 1989. Salinity tolerance of cultivated forages. *Facultad de Agronomia Veterinaria Universidad Nacional Del Litoral Senta Argentina*, 7(12): 113-116.
- Qu, X., Z. Huang, J. Baskin and C. Baskin. 2007. Effect of temperature light and salinity on germination and radicle growth of the geographically widespread halophytes. *Ann. Bot.*, 10: 1093.
- Sabir, P. and M. Ashraf. 2007. Screening of local accession *Panicum mliaceum* for salt tolerance at the seedling stage using biomass production and ion accumulation as selection criteria. *Pak. J. Bot.*, 39: 1655-2661.
- Shadded, M.A. and M.A. Zaidan. 1989. Effect of NaCl salinity on the rate of germination, seedling growth and some metabolic changes in *Raphanus sativa* L. and *Trigonella foenum-graecum* L. *Beitrag Zur Tropischen Landwirtschaft und Veterinarmedizin*, 27(2): 187-194.

- Song, J.G. Feng and F. Zhang. 2006. Salinity and temperature effect on three salt resistant euhalophytes, *Halostachys capsica*, *Kalidium foliatum* and *Halocnemum strobilaceum*. *Plant & Sci.*, 279: 201-207.
- Steel, R.G.D. and J.H. Tori. 1980. *Principles and Procedures of Statistics*. Second Edition MC Graw Hill Book Co., New York, U.S.A.
- Tobe, K., X. Li and K. Omasa. 2002. Effects of temperature on seed germination and radicle survival of *Kalidium capsicum* (Chenopodiaceae). *Aust. J. Bot.*, 50: 163-169.
- White, E.M. and E.R. Gartner. 1997. Salinity effects on blue grama (*Bouteloua gracilis*) growth in the green house. *Proc. S. D. Acad. Sci.*, 66(4): 76-84.
- [www.pakistaneconomist.com/issue2002/issue37/i&e4.htm](http://www.pakistaneconomist.com/issue2002/issue37/i&e4.htm)
- Zehra and M.A. Khan. 2007. Comparative effect of NaCl and sea salt on germination of halophytic grasses *Phragmites karka* at different temperature regimes. *Pak. J. Bot.*, 39: 1681-1694

(Received for publication 15 February 2008)