

DIVERSITY OF THE GENERA OF PENNATE DIATOMS IN THE PUNJAB

SYED TARIQ-ALI¹, MASUD-UL-HASAN² AND MUSTAFA SHAMEEL³

¹*Department of Botany, Federal Urdu University of Arts, Science & Technology,
Gulshan-e-Iqbal Campus, Karachi-75300*

²*Department of Botany, Govt. College University, Kachehri Road, Lahore-54000*

³*Department of Botany, University of Karachi, Karachi-75270, Pakistan*

Abstract

Twenty nine genera of algae belonging to 15 families of the phylum Bacillariophycota were collected during December 2003 and July 2006 from different freshwater habitats in various districts of the Punjab Province of Pakistan. Among them family Pinnularaceae with 19 species was most commonly found (25.33 %), next to it were the families Cymbellaceae (13.33 %), Naviculaceae (13.33 %) and Nitzschaceae (12 %) with 9 or 10 species each. The genera *Cymbella* with 10 species (13.33 %), *Nitzschia* with 9 species (12 %), *Navicula* and *Pinnularia* with 8 species each (10.66 %) occurred most commonly. About 64 % of diatoms were found as epiphytes, 22.66 % as planktic and epiphytic both, and 13.33 % as pure planktic individuals. Greatest proportion of collections (89.33 %) were made from Lahore City and its neighbouring areas, which was followed by the districts of Gujranwala (24 %) and Kasur (20 %). Most of the diatoms were collected in winter (42.66 %), smallest number of them were found in autumn (10.66 %), while in rest of the two seasons they occurred in almost equal proportions (25.33 & 26.66 %).

Introduction

Carter (1926) described for the first time 49 new species of freshwater diatoms from the region now included in Pakistan. Later on Abdul-Majeed (1935) gave an account of 62 diatoms of the undivided Punjab. A detailed investigation on the taxonomy of 102 species belonging to 25 genera of freshwater diatoms of Peshawar Valley (N. W. F. P.) was carried out by Salim & Khan (1960). Apart from that several attempts were also made to describe marine diatoms from the coastal waters of Karachi (Salim, 1954, 1963; Salim & Iqbal, 1964; Saifullah & Moazzam, 1978) and other areas of Sindh (Ghazala *et al.*, 2006).

Later on some species of freshwater diatoms were further reported from Punjab (Masud-ul-Hasan & Batool, 1987; Masud-ul-Hasan & Yunus, 1989), Islamabad (Inam *et al.*, 1986; Leghari *et al.*, 2002, 2004), N. W. F. P. (Leghari *et al.*, 1991, 1995, 2003), Balochistan (Anjum & Hussain, 1983; Leghari *et al.*, 2006), Sindh (Jahangir *et al.*, 2000; Leghari *et al.*, 2001, 2005a-c) and Azad Kashmir (Masud-ul-Hasan & Zeb-un-Nisa, 1986). But no composite study was carried out on the diversity of Bacillariophycota from any area of the country. Therefore, a wide range research program was started in March 2003 to make a detailed investigation on the diversity of this phylum and a large scale collection of diatoms was carried out from freshwater habitats of various districts of the Punjab.

Materials and Methods

Algal collections were made during December 2003 and July 2006 from various areas of the districts of Gujranwala, Jhang, Kasur, Lahore, Sargodha, Sheikhpura and Sialkot in the province of the Punjab. Planktic and epiphytic algae were collected from various freshwater habitats *e.g.*, rain pools, road-side ponds, fountain-water tanks, permanent water reservoirs, rivers, rice fields, canals and irrigation channels during different seasons of the year. The specimens were collected using phytoplankton net of 5-10 μm mesh. Water samples obtained by Nansen bottle from 3-4 m depth were filtered to separate planktic individuals. Epiphytic individuals were obtained by crushing the twigs, leaves and other parts of the collected vegetation. The algal specimens collected from different sources were centrifuged and preserved in plastic bottles containing 3 % formalin solution and brought to Karachi.

In the laboratory at Karachi the specimens were picked up using a dropper from the samples collected from various sources, placed in 10 % glycerin mounts and studied under Zeiss Stereomicroscope. Identification of the specimens down to the species level was carried out with the help of authentic literature (Cleve, 1894, 1895; Østrup, 1908; Hustedt, 1930; Salim & Khan, 1960; Giffen, 1963, 1966, 1970; Starmach, 1964; Nizamuddin, 1984).

Results and Discussion

Twenty-nine genera containing 75 species of planktic and epiphytic algae with solitary or colonial organisations belonging to one phylum, one class, one order and 15 families have been collected from various freshwater habitats. The collected specimens were investigated cytologically and morphologically, and on the basis of such characters they were taxonomically determined up to species level, but most of them appeared to be new records from these areas when compared with the previous observations (Inam *et al.*, 1986, Sultana *et al.*, 1991, Leghari *et al.*, 1995, 2002, 2004, Husna *et al.*, 2006, 2007, Shahida *et al.*, 2006). The collected genera have been systematically arranged according to the newly proposed classification (Shameel, 2001) with new taxonomic nomenclature suggested by Shameel (2008) has been used (Tables 1-3).

1. Taxonomic diversity: The order Bacillariales was represented in this study by 15 families, among them Pinnularaceae with 19 species was most commonly found (25.33 %). Next to it were the families Cymbellaceae (13.33 %), Naviculaceae (13.33 %) and Nitzschaceae (12.0 %) with 9 or 10 species (Table 1). At other localities also these families were found to be more prevalent than others (Cleve, 1894, 1895; Huber-Pestalozzi, 1942; Giffen, 1966, 1970). The families Bacillariaceae, Gomphonemaceae, Swirellaceae, Tabellaraceae, Thallassiosiraceae, Epithemaceae and Eunotaceae with 1 or 2 species were most poorly represented. While at the generic level, *Cymbella* with 10 species (13.33 %), *Nitzschia* with 9 species (12.0 %), *Navicula* and *Pinnularia* with 8 species each (10.66 %) occurred most commonly. *Bacillaria*, *Anomoeoneis*, *Fragilaria*, *Fragilariforma*, *Frustulia*, *Luticola*, *Mastogloia*, *Neidium*, *Stauroneis*, *Tabellaria*, *Cyclotella*, *Epithemia*, *Rhopalodia* and *Eunotia* with one species each, were very poorly represented genera in the present collection. Usually they were observed to be of rare occurrence in other studies (Thomson, 1938; Whitford, 1943; Prescott & Vinyard, 1965).

Table 1. Habitat diversity exhibited by the algal genera collected from Punjab Province of Pakistan.

Algal Taxa	Collected species		Planktic		Epiphytic		Planktic & Epiphytic	
	No	%	No	%	No	%	No	%
Phylum Bacillariophycota	75	65.21	10	13.33	48	64	17	22.66
Class Bacillariophyceae	75	65.21	10	13.33	48	64	17	22.66
Order Bacillariales	75	65.21	10	13.33	48	64	17	22.66
Family Achnantheaceae	4	5.33	-	-	3	75	1	25
<i>Achnanthes</i> Bory de Saint-Vincent	2	2.66	-	-	1	50	1	50
<i>Cocconeis</i> Ehrenberg	2	2.66	-	-	2	100	-	-
Family Bacillariaceae	1	1.33	-	-	1	100	-	-
<i>Bacillaria</i> J. F. Gmelin in Linnaeus	1	1.33	-	-	1	100	-	-
Family Cymbellaceae	10	13.33	1	10	5	50	4	40
<i>Cymbella</i> C. A. Agardh	10	13.33	1	10	5	50	4	40
Family Diatomaceae	3	4.0	-	-	3	100	-	-
<i>Diatoma</i> Bory de Saint-Vincent	3	4.0	-	-	3	100	-	-
Family Fragilaraceae	5	6.66	-	-	4	80	1	20
<i>Anomoeoneis</i> Pfitzer	1	1.33	-	-	-	-	1	100
<i>Fragilaria</i> Lyngbye	1	1.33	-	-	1	100	-	-
<i>Fragilariforma</i> (J. Ralfs) D.M. Williams et F. E. Round	1	1.33	-	-	1	100	-	-
<i>Synedra</i> Ehrenberg	2	2.66	-	-	2	100	-	-
Family Gomphonemaceae	2	2.66	-	-	2	100	-	-
<i>Gomphonema</i> Ehrenberg	2	2.66	-	-	2	100	-	-
Family Naviculaceae	10	13.33	9	90	1	10	-	-
<i>Frustulia</i> Rabenhorst	1	1.33	1	100	-	-	-	-
<i>Luticola</i> D.G. Mann in F.E. Round, R.M. Crawford et D.G. Mann	1	1.33	-	-	1	100	-	-
<i>Navicula</i> Bory emend. Cleve	8	10.66	8	100	-	-	-	-
Family Nitzschaceae	9	12.0	-	-	5	55.55	4	44.44
<i>Nitzschia</i> Hassall	9	12.0	-	-	5	55.55	4	44.44

Table 1. (Cont'd).

Algal Taxa	Collected species		Planktic		Epiphytic		Planktic & Epiphytic	
	No	%	No	%	No	%	No	%
Family Pinnularaceae	19	25.33	-	-	16	84.21	3	15.78
<i>Amphora</i> Ehrenberg ex Kützing	2	2.66	-	-	2	100	-	-
<i>Caloneis</i> Cleve in Cleve et Grove	3	4.0	-	-	3	100	-	-
<i>Gyrosigma</i> Hassall	3	4.0	-	-	2	66.66	1	33.33
<i>Mastogloia</i> Thwaites ex W. Smith	1	1.33	-	-	1	100	-	-
<i>Neidium</i> Pfitzer	1	1.33	-	-	1	100	-	-
<i>Pinnularia</i> (Ehrenberg) Ehrenberg	8	10.66	-	-	6	75	2	25
<i>Stauroneis</i> Ehrenberg	1	1.33	-	-	1	100	-	-
Family Surirellaceae	5	6.66	-	-	2	40	3	60
<i>Colletonema</i> de Brébisson ex Kützing	3	4.0	-	-	-	-	3	100
<i>Surirella</i> Turpin	2	2.60	-	-	2	100	-	-
Family Swirellaceae	2	2.60	-	-	2	100	-	-
<i>Cymatopleura</i> W. Smith	2	2.60	-	-	2	100	-	-
Family Tabellaraceae	1	1.33	-	-	1	100	-	-
<i>Tabellaria</i> C. G. Ehrenberg ex F.T. Kützing	1	1.33	-	-	1	100	-	-
Family Thalassiosiraceae	1	1.33	-	-	1	100	-	-
<i>Cyclotella</i> (Kützing) de Brébisson	1	1.33	-	-	1	100	-	-
Family Epithemaceae	2	2.60	-	-	1	50	1	50
<i>Epithemia</i> de Brébisson	1	1.33	-	-	1	100	-	-
<i>Rhopalodia</i> O. F. Müller	1	1.33	-	-	-	-	1	100
Family Eunotaceae	1	1.33	-	-	1	100	-	-
<i>Eunotia</i> Ehrenberg	1	1.33	-	-	1	100	-	-
Grand Total	75	-	10	13.33	48	64.0	17	22.66

No = Number of species, % = percentage of occurrence.

Table 2. Representation of algal genera in different localities (1-7) of the Punjab Province of Pakistan.

Algal Taxa	Number of species						
	*1	2	3	4	5	6	7
Phylum Bacillariophycota	14	2	15	37	1	2	5
Class Bacillariophyceae	14	2	15	37	1	2	5
Order Bacillariales	14	2	15	37	1	2	5
Family Achnantheaceae	-	-	-	1	1	-	-
<i>Achnanthes</i> Bory de Saint-Vincent	-	-	-	-	-	-	-
<i>Cocconeis</i> Ehrenberg	-	-	-	1	1	-	-
Family Bacillareaceae	-	-	-	-	1	-	-
<i>Bacillaria</i> J. F. Gmelin in Linnaeus	-	-	-	-	1	-	-
Family Cymbellaceae	-	-	4	4	-	-	1
<i>Cymbella</i> C. A. Agardh	-	-	4	4	-	-	1
Family Diatomaceae	1	-	-	3	-	-	-
<i>Diatoma</i> Bory de Saint-Vincent	1	-	-	3	-	-	-
Family Fragilaraceae	1	-	1	5	-	-	3
<i>Anomoeoneis</i> Pfitzer	-	-	-	1	-	-	1
<i>Fragilaria</i> Lyngbye	1	-	-	1	-	-	1
<i>Fragilariforma</i> (J. Ralfs) D.M. Williams et F.E. Round	-	-	1	1	-	-	-
<i>Synedra</i> Ehrenberg	-	-	-	2	-	-	1
Family Gomphonemaceae	-	-	-	1	-	-	-
<i>Gomphonema</i> Ehrenberg	-	-	-	1	-	-	-
Family Naviculaceae	6	-	1	3	-	1	-
<i>Frustulia</i> Rabenhorst	-	-	-	-	-	-	-
<i>Luticola</i> D.G. Mann in F.E. Round, R.M. Crawford et D.G. Mann	-	-	-	-	-	1	-
<i>Navicula</i> Bory emend. Cleve	6	-	1	3	-	-	-
Family Nitzschaceae	-	-	3	5	-	1	-
<i>Nitzschia</i> Hassall	-	-	3	5	-	1	-

Table 2. (Cont'd).

Algal Taxa	Number of species						
	*1	2	3	4	5	6	7
Family Pinnularaceae	4	2	6	7	-	-	-
<i>Amphora</i> Ehrenberg ex Kützing	-	-	2	-	-	-	-
<i>Caloneis Cleve in Cleve et Grove</i>	1	-	-	2	-	-	-
<i>Gyrosigma</i> Hassall	2	1	1	1	-	-	-
<i>Mastogloia</i> Thwaites ex W. Smith	-	-	1	-	-	-	-
<i>Neidium</i> Pfitzer	-	-	-	1	-	-	-
<i>Pinnularia</i> (Ehrenberg) Ehrenberg	1	1	2	3	-	-	-
<i>Stauroneis</i> Ehrenberg	-	-	-	1	-	-	-
Family Surirellaceae	-	-	-	5	-	-	-
<i>Colletonema</i> de Brébisson ex Kützing	-	-	-	3	-	-	-
<i>Surirella</i> Turpin	-	-	-	2	-	-	-
Family Swirellaceae	2	-	-	-	-	-	-
<i>Cymatopleura</i> W. Smith	2	-	-	-	-	-	-
Family Tabellaraceae	-	-	-	1	-	-	-
<i>Tabellaria</i> C. G. Ehrenberg ex F. T. Kützing	-	-	-	1	-	-	-
Family Thalassiosiraceae	-	-	-	-	-	-	-
<i>Cyclotella</i> (Kützing) de Brébisson	-	-	-	-	-	-	-
Family Epithemaceae	-	-	-	1	-	-	1
<i>Epithemia</i> de Brébisson	-	-	-	1	-	-	-
<i>Rhopalodia</i> O. F. Müller	-	-	-	-	-	-	1
Family Eunotaceae	-	-	-	1	-	-	-
<i>Eunotia</i> Ehrenberg	-	-	-	1	-	-	-
Grand Total	14	2	15	37	1	2	5
Percentage out of 75 species	18.66	2.66	20	49.33	1.33	2.66	6.66

*1 = Gujranwala, 2 = Jhang, 3 = Kasur, 4 = Lahore, 5 = Sargodha, 6 = Sheikhupura, 7 = Sialkot.

Table 3. Seasonal distribution of the algal genera collected from Punjab Province of Pakistan.

Algal Taxa	Coll. Sp.	Spring		Summer		Autumn		Winter	
		No	%	No	%	No	%	No	%
Phylum Bacillariophycota	75	20	26.66	19	25.33	8	10.66	32	42.66
Class Bacillariophyceae	75	20	26.66	19	25.33	8	10.66	32	42.66
Order Bacillariales	75	20	26.66	19	25.33	8	10.66	32	42.66
Family Achnanthesaceae	4	1	25	1	25	1	25	1	25
<i>Achnanthes</i> Bory de Saint-Vincent	2	1	50	-	-	-	-	1	50
<i>Cocconeis</i> Ehrenberg	2	-	-	1	50	1	50	-	-
Family Bacillariaceae	1	-	-	-	-	1	100	-	-
<i>Bacillaria</i> J. F. Gmelin in Linnaeus	1	-	-	-	-	1	100	-	-
Family Cymbellaceae	10	5	50	1	10	1	10	4	40
<i>Cymbella</i> C. A. Agardh	10	5	50	1	10	1	10	4	40
Family Diatomaceae	3	-	-	2	66.66	-	-	1	33.33
<i>Diatoma</i> Bory de Saint-Vincent	3	-	-	2	66.66	-	-	1	33.33
Family Fragilaraceae	5	1	20	4	80	-	-	-	-
<i>Anomoeoneis</i> Pfitzer	1	-	-	1	100	-	-	-	-
<i>Fragilaria</i> Lyngbye	1	-	-	1	100	-	-	-	-
<i>Fragilariforma</i> (J. Ralfs) D. M. Williams et F. E. Round	1	-	-	1	100	-	-	-	-
<i>Synedra</i> Ehrenberg	2	1	50	1	50	-	-	-	-
Family Gomphonemaceae	2	1	50	1	50	-	-	1	50
<i>Gomphonema</i> Ehrenberg	2	1	50	1	50	-	-	1	50
Family Naviculaceae	10	1	10	2	20	2	20	6	60
<i>Frustulia</i> Rabenhorst	1	-	-	1	100	-	-	-	-
<i>Luticola</i> D. G. Mann in F. E. Round, R.M. Crawford et D. G. Mann	1	-	-	-	-	1	100	-	-
<i>Navicula</i> Bory emend. Cleve	8	1	12.5	1	12.5	1	12.5	6	75
Family Nitzschaceae	9	1	11.11	3	33.33	2	22.22	4	44.44
<i>Nitzschia</i> Hassall	9	1	11.11	3	33.33	2	22.22	4	44.44

Table 3. (Cont'd).

Algal Taxa	Coll. Sp.	Spring		Summer		Autumn		Winter	
		No	%	No	%	No	%	No	%
Family Pinnularaceae	19	4	21.05	3	15.78	1	5.26	11	57.89
<i>Amphora</i> Ehrenberg ex Kützing	2	-	-	-	-	-	-	2	100
<i>Caloneis</i> Cleve in Cleve et Grove	3	1	33.33	1	33.33	-	-	1	33.33
<i>Gyrosigma</i> Hassall	3	-	-	-	-	-	-	3	100
<i>Mastogloia</i> Thwaites ex W. Smith	1	-	-	-	-	-	-	1	100
<i>Neidium</i> Pfitzer	1	-	-	1	100	-	-	-	-
<i>Pinnularia</i> (Ehrenberg) Ehrenberg	8	3	37.5	1	12.5	-	-	4	50
<i>Stauroneis</i> Ehrenberg	1	-	-	-	-	1	100	-	-
Family Surirellaceae	5	3	60	2	40	-	-	-	-
<i>Colletonema</i> de Brébisson ex Kützing	3	1	33.33	2	66.66	-	-	-	-
<i>Surirella</i> Turpin	2	2	100	-	-	-	-	-	-
Family Swirellaceae	2	-	-	-	-	-	-	2	100
<i>Cymatopleura</i> W. Smith	2	-	-	-	-	-	-	2	100
Family Tabellaraceae	1	1	100	-	-	-	-	-	-
<i>Tabellaria</i> C. G. Ehrenberg ex F. T. Kützing	1	1	100	-	-	-	-	-	-
Family Thalassiosiraceae	1	-	-	-	-	-	-	1	100
<i>Cyclotella</i> (Kützing) de Brébisson	1	-	-	-	-	-	-	1	100
Family Epithemaceae	2	1	50	-	-	-	-	1	50
<i>Epithemia</i> de Brébisson	1	-	-	-	-	-	-	1	100
<i>Rhopalodia</i> O. F. Müller	1	1	100	-	-	-	-	-	-
Family Eunotaceae	1	1	100	-	-	-	-	-	-
<i>Eunotia</i> Ehrenberg	1	1	100	-	-	-	-	-	-
Grand Total	75	20	26.66	19	25.33	8	10.66	32	42.66

Coll. Sp. = Collected species, No = number of species, % = percentage of occurrence.

2. Habitat diversity: The collected genera were found to occur in three different ways of living (Table I): (1) only in planktic condition, (2) only in epiphytic habitat, (3) both in planktic and epiphytic conditions (sometimes free-floating, sometimes attached with other algae and water plants). No epilithic, epipsammic or epipellic alga was collected, none of them was found simultaneously in planktic and benthic conditions (Table 1). Almost 64 % of them occurred as epiphytes, 22.66 % as planktic and epiphytic both, and 13.33 % as pure planktic individuals. There are several families such as Achnantheaceae, Bacillariaceae, Diatomaceae, Fragilaraceae, Gomphonemaceae, Nitzschaceae, Pinnularaceae, Surirellaceae, Swirellaceae, Tabellaraceae, Thalassiosiraceae, Epithemaceae and Eunotaceae, in which no purely planktic individual was observed. As the diatoms are non-motile and non-flagellated individuals, they usually occur in attached conditions as epiphyte (Østrup, 1908; Hustedt, 1930; Giffen, 1963; Starmach, 1964; Hohn & Hellermann, 1966).

3. Diversity of localities: The algae were collected from 7 different localities, wherefrom the collected species exhibited variations in their diversities (Table 2). Greatest proportion of algal collections (89.33 %) were made from Lahore City and its neighbouring areas. Largest collections of blue-green and green algae were also made from this district during previous studies (Naz *et al.*, 2009, Zarina *et al.*, 2009). Probably, it may be due to the presence of innumerable water bodies such as rivers, canals, tubewells and fountain-displaying areas at Lahore (Husna *et al.*, 2006; Munir *et al.*, 2006). Other groups of freshwater algae were also found in abundance at Lahore (Masud-ul-Hasan, 1978a, b, 1980; Masud-ul-Hasan & Yunus, 1989). It was followed by the districts of Gujranwala (24 %) and Kasur (20 %), where appreciable collections were made. The poorest diversity was shown in the collections from the districts of Jhang and Sargodha (4.0 & 5.33 %). During previous studies, poor growth of algae was also recorded for the districts of Jhang and Sargodha (Faridi *et al.*, 1981; Masud-ul-Hasan & Batool, 1987; Naz *et al.*, 2009).

4. Seasonal diversity: The collected diatoms were found in all the four seasons of the year (Table 3). Most of the species were collected in winter (42.66 %), smallest number of them were found in autumn (10.66 %), while in rest of the two seasons they occurred in almost equal proportions (26.66 and 25.33 %). It was similarly observed in other localities of the world (Thomson, 1938; Gerloff & Lüdemann, 1966). Different genera behaved variably in their pattern of seasonal diversity. *Cymbella* occurred mainly (50 %) in spring, *Navicula* (75 %) and *Nitzschia* (44.44 %) were found mainly in winter, while they were collected in all the four seasons of the year. Species of *Surirella*, *Tabellaria*, *Rhopalodia* and *Eunotia* were only found in spring, while they were absent in other seasons. Species of *Anomoeoneis* and *Neidium* occurred only in summer, they were not found in other seasons. Species of *Bacillaria*, *Luticola* and *Stauroneis* were found only in autumn, being absent in other three seasons. Species of several genera such as *Amphora*, *Gyrosigma*, *Mastogloia*, *Cymatopleura*, *Cyclotella* and *Epithemia* occurred during winter season only and could not be collected in other seasons of the year. Certain genera were found in any two seasons and remained absent in the other two seasons. In this regard, genera of the naviculoid diatoms exhibited great variation in their seasonal diversity among themselves, as was observed quite earlier (Cleve, 1894, 1895).

References

- Abdul-Majeed, M. 1935. *Freshwater algae of the Punjab. I. Bacillariophyta (Diatomeae)*. Punj. Univ. Publ., Lahore, 45 pp.

- Anjum, G. and F. Hussain. 1983. The genera *Cymbella* and *Navicula* from Quetta-Pishin Valley, Balochistan. *J. Sci. Tech.*, 7: 1-7.
- Carter, N. 1926. Freshwater algae from India. *Rec. Bot. Surv. Ind.*, 9: 263-302.
- Cleve, P.T. 1894. Synopsis of the naviculoid diatoms-I. *Kongl. Svensk. Vet.-Akad. Handl.*, 26: 1-194.
- Cleve, P.T. 1895. Synopsis of the naviculoid diatoms-II. *Kongl. Svensk. Vet.-Akad. Handl.*, 27: 1-129.
- Faridi, M.A.F., A.T. Anjum and G. Anjum. 1981. Algae associated with alluvial gold of Indus at Attock. *Geol. Bull. Peshawar Univ.*, 14: 151-157.
- Gerloff, J. and D. Lüdemann. 1966. *Leitfaden der Trink- und Brauch-wasserbiologie*. Gust. Fisch. Verlag, Stuttgart, 360 pp.
- Ghazala, B., R. Ormond and F. Hannah. 2006. Phytoplankton communities of Pakistan: I. Dinophyta and Bacillariophyta from the coast of Sindh. *Int. J. Phycol. Phycochem.*, 2: 183-196.
- Giffen, M.H. 1963. Contribution to the diatom flora of South Africa. I. Diatoms of the estuaries of the Eastren Cape Province. *Hydrobiol.*, 21: 201-265.
- Giffen, M.H. 1966. Contribution to the diatom flora of South Africa. III. Diatoms of the marine littoral regions at Kidd's Beach near East London, Cape Province, South Africa. *Nova Hedw.*, 13: 245-292+5 pls.
- Giffen, M.H. 1970. New and interesting marine and littoral diatoms from Sea Point, near Cape Town, South Africa. *Bot. Mar.*, 13: 87-99.
- Hohn, M.H. and J. Hellermann. 1966. New diatoms from the Lewes-Rehoboth Canal, Delaware and Chesapeake Bay area of Baltimore, Maryland. *Trans. Amer. Microscop. Soc.*, 85: 115-130.
- Huber-Pestalozzi, G. 1942. Das Phytoplankton des Süßwassers. II. Diatomeen. In: *Die Binnengewässer*. 2, (Ed.): Thienemann, p. 1-549.
- Husna, R., A. Zarina, Masud-ul-Hasan and M. Shameel. 2006. Taxonomic study of some microalgae from Lahore, Pakistan. *Int. J. Phycol. Phycochem.*, 2: 157-164.
- Husna, R., A. Zarina, Masud-ul-Hasan and M. Shameel. 2007. Taxonomic study of some diatoms from Lahore, Pakistan. *Int. J. Phycol. Phycochem.*, 3: 55-64.
- Hustedt, F. 1930. Bacillariophyta. In: *Süßwasserflora*. 10, (Ed.): Pascher, p. 1-466.
- Inam, B., K. Nazir and M.K. Leghari. 1986. Some diatoms from Islamabad-I. *J. Sci. Technol.*, 10: 1-3.
- Jahangir, T.M., M.Y. Khuhawar, S.M. Leghari, W.A. Balouch, A.A. Leghari and A. Leghari. 2000. Some studies on water quality and biological life at Kinjhar and Haleji lakes of district Thatta, Sindh, Pakistan. *Pak. J. Biol. Sci.*, 3: 1965-1972.
- Leghari, M.K., K. Sultana and T. Bando. 1991. Taxonomic studies of order Naviculales Malka Parbat: Kaghan (Part I). *Biologia*, 37: 9-12.
- Leghari, M.K., K. Sultana and M. Haga. 1995. Diatoms from unexplored Diامر face of Nanga Parbat Part I. *Biologia*, 41: 11-12.
- Leghari, M.K., M.Y. Leghari and M. Shah. 2002. Ecological study of algal flora of Korang Nalla of Rawal Dam, Islamabad. *Biologia*, 48: 65-80.
- Leghari, M.K., M.Y. Leghari and M. Shah and S.N. Arbani. 2003. Ecological study of algal flora of Wah Garden, district Attock, Pakistan. *Pak. J. Bot.*, 35: 705-716.
- Leghari, M.K., M.Y. Leghari and S.M. Leghari. 2004. Water chemistry and its relation with algae of Rawal Dam, Islamabad and Wah Garden, district Attock. *Sindh Univ. Res. J. (Sci. Ser.)*, 36: 29-48.
- Leghari, S.M., T.M. Jahangir, M.Y. Khuhawar and A. Leghari. 2001. Physico-chemical and biological study of Dhabeji springs, Malir, Karachi, Sindh, Pakistan. *Online J. Biol. Sci.*, 1: 623-627.
- Leghari, S.M., M.Y. Khuhawar, T.M. Jahangir and A. Abdullah. 2005a. Limnological study of natural springs at Gharo Creek, district Thatta, Sindh, Pakistan. *Int. J. Phycol. Phycochem.*, 1: 37-42.
- Leghari, S.M., M.Y. Khuhawar, T.M. Jahangir and A. Abdullah. 2005b. Limnological study of Pir Bukhari (Karsaz) and Manghopir warm springs, Karachi, Sindh, Pakistan. *Int. J. Phycol. Phycochem.*, 1: 151-158.

- Leghari, S.M., T.M. Jahangir, M.Y. Khuhawar and A. Leghari. 2005c. Some studies on Toungh natural springs of Thana Bula Khan, district Dadu, Sindh, Pakistan. *Int. J. Phycol. Phycochem.*, 1: 167-172.
- Leghari, S.M., M.Y. Leghari, M.A. Mahar, M.Y. Khuhawar and T.M. Jahangir. 2006. Limnological study of Hanna Lake and Urak Spring of Quetta, Balochistan, Pakistan. *Int. J. Phycol. Phycochem.*, 2: 33-38.
- Masud-ul-Hasan. 1978a. A contribution to the freshwater algae of the Punjab-I. *Biologia*, 24: 25-30.
- Masud-ul-Hasan. 1978b. A contribution to the freshwater algae of the Punjab-II. *Biologia*, 24: 81-96.
- Masud-ul-Hasan. 1980. A contribution to the freshwater algae of the Punjab-III. *Biologia*, 26: 71-79.
- Masud-ul-Hasan and I. Batool. 1987. A taxonomic study of some freshwater algae from Attock and Sargodha District. *Biologia*, 33: 345-366.
- Masud-ul-Hasan and A. Yunus. 1989. An addition to the algal flora of Lahore. *Biologia*, 35: 99-131.
- Masud-ul-Hasan and Zeb-un-Nisa. 1986. Taxonomic studies of some freshwater algae from Azad Jammu and Kashmir. *Biologia*, 32: 229-256.
- Munir, M.A., Masud-ul-Hasan and M. Shameel. 2006. Seasonal variation of algal genera in a freshwater pond near Lahore and its physicochemical properties. *Int. J. Phycol. Phycochem.*, 2: 197-202.
- Naz, S., A. Zarina, Masud-ul-Hasan and M. Shameel. 2009. Diversity of Cyanophycota in the north-eastern areas of Pakistan. *Proc. Pak. Acad. Sci.*, 46: 29-40.
- Nizamuddin, M. 1984. *Diatoms of Libya*. Dept. of Botany, Univ. Al-Fateh, Tripoli, 144 pp.
- Østrup, E. 1908. Freshwater diatoms. In: *Botany of the Faeröes Based Upon Danish Investigations*. (Ed.): E. Warming, Gyldendalske Boghandel, Nordisk Forlag, Copenhagen, p. 260-290.
- Prescott, G.W. and W.C. Vinyard. 1965. Ecology of Alaskan freshwater algae: V. Limnology and flora Malikpuk Lake. *Trans. Amer. Microscop. Soc.*, 84: 427-478.
- Saifullah, S.M. and M. Moazzam. 1978. Species composition and seasonal occurrence of centric diatoms in a polluted marine environment. *Pak. J. Bot.*, 10: 53-64.
- Salim, K.M. 1954. Some plankton diatoms from the Karachi Coast. *Pak. J. Sci.*, 6.
- Salim, K.M. 1963. A systematic account of some marine diatoms from the Karachi Coast. *Pesh. Univ. J.*, 8: 19-52.
- Salim, K.M. and M.M. Iqbal. 1964. Distribution of diatoms in the intertidal zone, rocky ledge of Manora. *Pak. J. Sci.*, 16.
- Salim, K.M. and M.H. Khan. 1960. *The Diatomales: The Fresh Water Diatoms of Peshawar Valley*. Dept. Botany, Peshawar Univ., Peshawar, 66 pp. +11 pls.
- Shahida, B., A. Zarina, Masud-ul-Hasan and M. Shameel. 2006. Taxonomic study of some Volvocophyta and Bacillariophyta from Rabwah and Sargodha, Pakistan. *Int. J. Phycol. Phycochem.*, 2: 173-182.
- Shameel, M. 2001. An approach to the classification of algae in the new millennium. *Pak. J. Mar. Biol.*, 7: 233-250.
- Shameel, M. 2008. Change of divisional nomenclature in the Shameelian Classification of algae. *Int. J. Phycol. Phycochem.*, 4: 225-232.
- Starmach, K. 1964. *Flora Slodkowodna Polski*, 6. *Chrysophyta, II. Bacillariophyceae-Okrzemki*. Panstwowe Wydawnictwo Naukowe, 610 pp.
- Thomson, R. 1938. A preliminary survey of the freshwater algae of eastern Kansas. *Univ. Kansas. Sci. Bull.*, 25: 5-83.
- Whitford, L.A. 1943. *The Freshwater Algae of North Carolina*. Dept. Bot., Ohio State Univ. 445:131-171.
- Zarina, A., Masud-ul-Hasan and M. Shameel. 2009. Diversity of freshwater green macro-algae in the Punjab and neighbouring areas of Pakistan. *Pak. J. Bot.*, 41: 277-291.