

## INTERACTIVE EFFECT OF HUMIC ACID AND FARMYARD MANURE ON SOIL HEALTH AND MICROBIAL ACTIVITY IN CALCAREOUS SOIL

SHAHZAD AFZAL<sup>1,2</sup>, DOST MUHAMMAD<sup>1</sup>, RAFI ULLAH<sup>3</sup>, MUHAMMAD ADNAN<sup>3\*</sup>,  
BEENA SAEED<sup>3</sup>, RASHA M. ALZAYED<sup>4</sup>, SONDOS A. ALHAJOUJ<sup>4</sup>, MEAAD F. ALAIDA<sup>4</sup>,  
MANZOOR AHMAD<sup>5</sup>, A. ALTALHI<sup>6</sup> AND MADEHA O. I. GHOBASHY<sup>7,8\*</sup>

<sup>1</sup>Department of Soil and Environmental Sciences, The University of Agriculture Peshawar, 25120, Peshawar-Khyber Pakhtunkhwa-Pakistan

<sup>2</sup>Shenzhen Key Laboratory of Marine Microbiome Engineering, Institute for Advanced Study, Shenzhen University, Shenzhen 518060, China

<sup>3</sup>Department of Agriculture, University of Swabi, Anbar-23561, Swabi-Khyber Pakhtunkhwa-Pakistan

<sup>4</sup>Biology Department, College of Science, Jouf University, Sakaka 41412, Saudi Arabia

<sup>5</sup>Department of Agriculture, Bacha Khan University Charsada, Khyber Pakhtunkhwa-Pakistan

<sup>6</sup>Biology, College of Science, Taif University, P.O. Box 11099, Taif 21944, Saudi Arabia

<sup>7</sup>Department of Biology, Faculty of Science, University of Tabuk, Tabuk, 71491, Saudi Arabia

<sup>8</sup>Biodiversity Genomics Unit, Faculty of Science, University of Tabuk, Tabuk, 71491, Saudi Arabia

\*Corresponding author's email: [madnan@uoswabi.edu.pk](mailto:madnan@uoswabi.edu.pk) ; [mghobashy@ut.edu.sa](mailto:mghobashy@ut.edu.sa)

### Abstract

Humic acid (HA) improves soil solids surface chemistry and thereby optimizes biological activity and fertility of soil. However, its interactive effect with farmyard manure (FYM) under calcareous soil is not fully explored. Therefore, a laboratory incubation experiment (1<sup>st</sup>) was conducted to investigate effect of various levels of lignitic coal derived humic acid (0, 100 and 200 mg kg<sup>-1</sup> soil) applied alone or in conjunction with 10 g FYM kg<sup>-1</sup> on soil microbial activity and nutrients availability for 79 days. In the 2<sup>nd</sup> experiment, the same soil was remoistened and treated with lower levels of HA (0, 2, 4, 6, 8 and 10 mg kg<sup>-1</sup>) for 14 days to further verify its effect on soil microbial activity. The combine application of HA and FYM significantly increased CO<sub>2</sub> release except at 2<sup>nd</sup> day, however, application of 100 and 200 mg HA produced non-consistent supremacy over each other at different incubation intervals. Combine use of HA and FYM consistently accelerated the rate of CO<sub>2</sub> production over all incubation intervals as compared to the sole application of HA or control. Cumulative release of CO<sub>2</sub> also showed that application of HA either as 100 or 200 mg produced higher (1014 and 1035 mg CO<sub>2</sub> kg<sup>-1</sup> respectively) as compared to control (918 mg kg<sup>-1</sup>) at 79<sup>th</sup> day. Integrated use of HA+FYM further amplified cumulative CO<sub>2</sub> releases as compared to control or alone HA at both levels. In the second experiment, application of HA increased CO<sub>2</sub> release with increasing doses up to 6 mg HA kg<sup>-1</sup> only at day 4 of incubation while its effect was non-significant for the rest of incubation interval. The integrated application of HA and FYM significantly improved post incubated soil pH, Organic matter and NPK content compared to their sole application. Therefore, it can be concluded that the application of FYM further amplify the positive effect of humic acid on soil health and shall be adopted.

**Key words:** Humic acid, Farmyard manure, Cumulative CO<sub>2</sub> releases, Incubated soil, Microbial activity.

### Introduction

Calcareous soils contain large amount of CaCO<sub>3</sub> and cover more than 30% of the Earth's surface (Bolan *et al.*, 2023). High CaCO<sub>3</sub> adversely affect the physical and chemical properties of these soils (FAO, 2016). Low water-holding capacity, high pH and infiltration rate, low CEC, low organic matter (OM) and clay contents, poor structure, nutrient loss by leaching or deep percolation, surface crusting and cracking, and N fertilizer loss are only a few of the difficulties associated with cultivating these soils, nutritional imbalances, especially P and micronutrients (El-Hady & Abo-Sedera, 2006). However, organic matter/substances and the availability of N can mitigate the high carbonate content related adversities in calcareous soils.

Organic substances release essential nutrients upon decomposition (Andriamananjara *et al.*, 2019). Farmyard manure (FYM) has a beneficial effect on soil biota, microbial activity and nutrient transformation, chemical properties like CEC, pH, nutrient chelation, solubility and physical properties like porosity, bulk density, structure and movement of states and energy (Melero *et al.*, 2009). It enhances the bio-availability of nutrients, reduce losses,

accelerate microbial activities and release such enzymes and hormones that ultimately improve crop yield and quality (Adnan *et al.*, 2018).

Humic acid (HA) has been used as a soil conditioner and fertilizer in agriculture on a small scale. Humic compounds have been shown to have important effects on plant development and soil structure. Their application in right quantities can help plants and roots to develop faster (Minhas *et al.*, 2024). Commercial nutrient-containing products known as humic substances (HS) enhance soil fertility, boost nutrient availability, lessen the adverse effects of chemical fertilizers, and eliminate harmful NO<sub>2</sub> and NO<sub>3</sub> ions from the soil, all of which improve plant performance (Osman & Rady, 2012; Burton *et al.*, 2003). According to Sangeetha *et al.*, (2006), they make up the majority of the soil's organic matter. By altering the soil's chemical and biological composition, they restore soil fertility. They also enhance the performance and biomass of plants. By improving soil structure and qualities like aggregation, aeration, permeability, water-holding capacity, and the availability and movement of micronutrients, HS can indirectly raise soil fertility (Cimrin & Yilmaz, 2005). They have the direct ability to influence the mechanisms involved

in their absorption and movement into plant tissues (Nardi *et al.*, 2002). Furthermore, by forming complex forms or chelating agents with metallic cations, HS can alter the solubility of numerous nutrients (Ouni *et al.*, 2014). According to Jamal *et al.*, (2023), HS can interact with P in the soil to boost plant uptake and decrease fixation. Under conditions of salinity and high carbonate content, the HA can enhance plant performance and the uptake of macro and micronutrients by plants (Katkot *et al.*, 2009). By providing nutrients to developing plants, HS can improve the soil's fertility and yield (Osman & Rady, 2012; Brye *et al.*, 2006). Barron and Humic acid increases the activity of microbes, cation exchange capacity and forms chelates with other nutrients, buffers soil pH (Julie & Bugbee, 2006). Sarlak *et al.*, (2024) also reported that humic acid serves as catalyst in increasing microorganism activities in soil.

Almost 90% of Pakistan's soils are calcareous in nature, alkaline in reaction, low in organic matter and deficient in several essential nutrients. The main practice to correct the nutrients deficiency in Pakistan is applying the chemical fertilizers. The price of chemical fertilizers is increasing with exponential rate posing the main obstacle in balanced and economical fertilization in Pakistan. Huge amount is spent on the import of chemical fertilizer every year. Along with higher input costs, continuous use of inorganic fertilizers may not be sustainable and can cause soil degradation and hazardous effects of environment (Sarlaki *et al.*, 2024). Hence chemical fertilizer needs to be supplemented with organic manures to produce on sustainable basis for ever increasing population and keep soil and environment from further decay. We hypothesized that, the combine application of humic acid with FYM has preservative effect in promoting the availability of nutrients, production and yield of different crops. However, their performance depend on soil and climatic conditions. Therefore, in the present study, influence of various levels of HA applied alone or in mixture with organic manure on microbial activity and soil fertility was investigated under calcareous conditions to verify their beneficial role in crop production.

## Material and Method

**Soil sampling:** The fresh composite soil sample was collected from top surface (0-15 cm deep) from the new developmental farm the University of Agriculture Peshawar, Khyber Pukhtoonkhwa, Pakistan. The fresh soil sample was shade dried, and sieved (mesh size < 2mm). The moisture content in soil was determined for the purpose of correction of data on dry mass basis. The experimental soil was non saline ( $0.36 \text{ dSm}^{-1}$ ) alkaline (7.6 pH), low in organic matter (1.14%), N ( $55.1 \text{ mg kg}^{-1}$ ) and P ( $3.20 \text{ mg kg}^{-1}$ ) while, was sufficient in K ( $180.6 \text{ mg kg}^{-1}$ ). The experimental site is semi-arid (300– 500 mm rainfall year<sup>-1</sup>) in nature where more than 60% rainfall occurs in summer season (Adnan *et al.*, 2018).

**Experimentation:** In 2016, two laboratory incubation experiments were carried out in the Department of Soil and Environmental Sciences' Soil Microbiology laboratory at the University of Agriculture, Peshawar, Pakistan. A basic completely randomized design (CRD) with three

replications was used for both trials. In order to confirm their advantageous role in crop production, the purpose of these experiments was to assess the effects of different concentrations of HA applied either alone or in combination with organic manure on microbial activity and soil fertility under calcareous conditions. Therefore, in 1<sup>st</sup> experiment, a 50 g air dried collected non-saline, calcareous soil (already collected and processed) was taken in a 500 ml conical flask and treated with HA at the rate of 0, 100, 200  $\text{mg kg}^{-1}$  with and without FYM ( $20 \text{ t ha}^{-1}$  or  $10 \text{ g kg}^{-1}$ ). Basal dose of N and K was applied to all treatments at the rate of 60 mg N and 30  $\text{mg K}_2\text{O kg}^{-1}$  as urea and sulfate of potash. Prepared flasks along with three blanks were further processed for  $\text{CO}_2$  evaluation according to the procedure of Horwath & Paul (1994) and incubated at  $28 \pm 2^\circ\text{C}$  for 79 days. Microbial activity as index for  $\text{CO}_2$  release from soil was measured on 2, 5, 11, 21, 36, 52, 79 day of incubation. Similarly, another experiment was conducted with lower levels of humic acid (0, 2, 4, 6, 8 and 10  $\text{mg HA kg}^{-1}$ ) for a period of 14 days to see effect of lower levels of HA on soil microbial activity at day 4, 8, 11 and 14 of incubation. The distilled water was spared on soil on the day of reading to balance the moisture loss by maintaining the initial weight of flask containing soil.

**Laboratory analysis:** The conical flask with a capacity of 500 mL were used, to take a 50 g HA and FYM treated soil as per proposed treatment structure. A 5 ml of 0.3 N NaOH were taken in a vial and properly sealed in a conical flask, and incubated at  $28^\circ\text{C}$  for 0, 3, 5, 11, 21, 36, 52, and 79 days for the first experiment and 4, 8, 11 and 14 days for the second experiment. The vial was taken out at each incubation period and titrated against 0.1 N HCl in the presence of 10 ml 1 M  $\text{BaCl}_2$  solution using phenolphthalein as indicator. The amount of HCl consumed in titration was used to calculate the amount of  $\text{CO}_2$  in each flask. The  $\text{CO}_2$  produced was calculated by difference between sample and blank (without soil) readings by the following formula (Frioni, 1990):

$$\text{CO}_2 \text{ release (mg kg}^{-1} \text{ CO}_2 - 1 = \frac{(\text{Blank} - \text{Sample}) \times E}{W \times \text{Days}}$$

where blank and samples are is average HCl volume used in the titration of the blanks and samples, respectively, W is weight of dry soil (g), and E is  $\text{CO}_2$  equivalent.

Changes in soil pH were measured in 1:5 suspensions (McClean, 1996), organic matter by using  $\text{K}_2\text{Cr}_2\text{O}_7$  as an oxidizing agent as decried by (Nelson & Sommer, 1996) AB-DTPA extractable P and K by the method (Soltanpour & Schwab, 1977) and mineral N by Kjeldahl distillation method (Mulvaney, 1996). These parameters were determined in the soil at the end of incubation to see the effect of humic acid and farmyard manure.

## Statistical analysis

Analysis of variance (ANOVA) was carried out according to complete randomized design using MSTATC package (Russel, 1989). Means were also compared using the least significant difference (LSD) test (Steel & Torrie, 1980).

## Results

**Rate of CO<sub>2</sub> release (mg kg<sup>-1</sup>) from incubated soil:** The release of CO<sub>2</sub> showed significant ( $p < 0.05$ ) increase with application of humic acid and FYM except on day 2<sup>nd</sup> of incubation where its effect was non-significant (Table 1). The CO<sub>2</sub> release tended to increase non-significantly with 100 and 200 mg Kg<sup>-1</sup>. However, the conjunctive use of HA+FYM consistently accelerated the rate of CO<sub>2</sub> production over all incubation intervals as compared to sole supplementation of HA or control but supremacy over sole application of FYM was variable at different incubation intervals. At some incubation intervals sole application of FYM produced higher CO<sub>2</sub> release as compared to HA+FYM that might be associated with higher doses of HA. On 5<sup>th</sup> day of incubation, the conjunctive use of 100 mg HA+FYM and 200 mg HA+FYM resulted in higher CO<sub>2</sub> releases of 35.82 and 35.62 as compared to 32.82 mg CO<sub>2</sub> kg<sup>-1</sup> d<sup>-1</sup> with application of FYM alone. Sole application of HA also produced higher rate of CO<sub>2</sub> as compared to control but this was less than FYM or HA+FYM. On day 11<sup>th</sup>, HA applied alone or in combination with FYM produced higher CO<sub>2</sub> as compared to control but less than sole FYM. Though conjunctive use of HA+FYM produced less CO<sub>2</sub> as compared to sole FYM but the difference was non-significant. On day 21, similar trends were observed. One day 36<sup>th</sup>, the conjunctive use of HA+FYM resulted significantly ( $p < 0.05$ ) higher releases of CO<sub>2</sub> as compared to control and sole application of HA but non-significant to sole FYM. On day 52 and 79 similar trends were observed but this time HA at 200 mg kg<sup>-1</sup> + FYM produced lower CO<sub>2</sub> as compared to alone FYM or 100 mg HA+FYM. Comparing changes of CO<sub>2</sub> release over incubation intervals, it was observed that initially the rate of CO<sub>2</sub> releases was increased up to day 21 for control FYM and day 36 for 10 g FYM kg<sup>-1</sup> soil and then declined with time up to 79<sup>th</sup> day of incubation.

**Cumulative CO<sub>2</sub> release (mg kg<sup>-1</sup>):** The results of cumulative release of CO<sub>2</sub> showed discrete differences among various treatments at the end of 79<sup>th</sup> day of incubation period. Application of HA either as 100 or 200 mg kg<sup>-1</sup> soil produced higher CO<sub>2</sub> release of 1014 and 1035 mg kg<sup>-1</sup>, respectively as compared to 918 mg kg<sup>-1</sup> in case of no-HA (control) (Fig. 1). Conjunctive use of HA+FYM also showed remarkable increase in CO<sub>2</sub> releases as compared to control (0 HA and 0 FYM) or alone HA at both levels, however, these releases were less as compared to sole application of FYM. The highest cumulative CO<sub>2</sub> release (mg kg<sup>-1</sup>) was observed in sole FYM treated soil while the lowest was observed in control. Similarly, under

the 2<sup>nd</sup> incubation study the highest cumulative Co2 was recorded under 6 mg HA ha<sup>-1</sup> while the lowest were recorded under control as shown in Fig. 2.

**Release of CO<sub>2</sub> from incubated soil treated with lower doses of HA:** In the 2<sup>nd</sup> incubation study where CO<sub>2</sub> release was evaluated with lower doses as shown in table 2. The results showed that application of HA consistently increased CO<sub>2</sub> release from soil with increasing doses up to 6 mg HA kg<sup>-1</sup> at day 4<sup>th</sup> of the incubation while on day 8, 11 and 14 its effect was non-significant. Similarly, the highest cumulative release of CO<sub>2</sub> as 594 mg kg<sup>-1</sup> was recorded at 6.0 mg HA kg<sup>-1</sup> as compared to control (502 mg kg<sup>-1</sup>) at the end of 14 days of incubation. These results confirmed the influential effect of HA on microbial activates and suggested that lower doses of HA upto 6 mg kg<sup>-1</sup> soil equivalent to 12 kg ha<sup>-1</sup> was the most appropriate doses of HA for improving the microbial activity in soil.

**Changes in soil pH, SOM, mineral N and AB-DTPA extractable P and K:** The analysis of variance indicated that integrated application of HA and FYM significantly affected soil pH, organic matter, mineral N and ABDTPA extractable P and K as compared to control (0 HA and 0 FYM) (Table 3). The sole application of HA at the rate of 200 mg kg<sup>-1</sup> soil significantly lowered soil pH from 7.72 in control to 7.51 while there was no effect of HA on soil pH where FYM was applied at the rate of 10 g kg<sup>-1</sup> soil. There organic matter content didn't show significant response to HA application under the same level of organic matter, however under the both control and 10 g FYM kg<sup>-1</sup> soil organic matter content was increased with increasing level of HA. Application of HA at the rate of 200 mg and 100 mg per kg soil along with + 10g FYM kg<sup>-1</sup> soil resulted higher soil organic matter of 1.57 and 1.52% respectively as compared to 1.41% in sole FYM. Application of HA at 100 and 200 mg kg<sup>-1</sup> soil produced OM of 1.15 and 1.14% which is similar to 1.12% of the control. The conjunctive use of HA+FYM produced higher organic matter content than alone application of FYM and HA treatments. The higher mineral N and ABDTPA extractable P and K of 136.2, 12.84 and 194 mg kg<sup>-1</sup> respectively were observed in treatments receiving 200 mg HA + FYM as compared to 104.7, 8.87 and 183 mg kg<sup>-1</sup> soil in control respectively. The P concentration showed little changes with HA applied. The highest P as 14.10 mg kg<sup>-1</sup> was recorded in sole FYM treatment which was statistically at par to 100 and 200 mg HA kg<sup>-1</sup> + 10 g FYM kg<sup>-1</sup> soil while significantly higher than 8.87 mg kg<sup>-1</sup> in control. Combined application of HA+ FYM significantly improved soil mineral N, ABDTPA extractable K and organic matter while didn't affect ABDTPA compared to sole application of FYM.

**Table 1. Effect of HA and FYM on rate of CO<sub>2</sub> at given intervals released from soil incubated for 80 days at 28±2°C.**

Treatment (kg <sup>-1</sup> soil)		Incubation time (days)						
		2	5	11	21	36	52	79
HA (mg)	FYM (g)	CO <sub>2</sub> Release (mg kg <sup>-1</sup> d <sup>-1</sup> )						
0	0	63.50	20.61 b	16.81 b	9.91 c	6.95 b	12.57 b	8.29 c
100	0	60.37	23.02 b	18.70 b	10.38 c	8.86 b	12.33 b	10.30 bc
200	0	56.93	25.22 b	17.33 b	11.94 c	8.52 b	13.72 b	10.17 bc
0	10	68.19	32.82 a	24.37 a	17.57 a	15.13 a	17.99 a	13.61 a
100	10	66.32	35.82 a	23.42 a	16.29 ab	17.38 a	17.93 a	11.87 ab
200	10	66.00	35.62 a	22.58 a	13.63 bc	17.44 a	15.82 ab	12.27 ab

Means with different letter(s) are significantly different ( $p < 0.05$ ). HA, FYM, OM, N, P and K represent humic acid, farmyard manure, organic matter, nitrogen, phosphorus and potassium, respectively




Fig. 1. Cumulative CO<sub>2</sub> released from soil treated with HA and FYM during incubation at room temperature (28±2°C). HA and FYM represents humic acid and farmyard manure, respectively.




Fig. 2. Cumulative CO<sub>2</sub> (mg kg<sup>-1</sup>) released from soil treated with lower doses of HA during incubation at room temperature (28±2°C). HA stands for humic acid.

Table 2. Release of CO<sub>2</sub> from soil treated with lower doses of HA.

HA (mg kg <sup>-1</sup> )	Incubation interval (days)			
	4	8	11	14
	CO <sub>2</sub> (mg kg <sup>-1</sup> d <sup>-1</sup> )			
0	30.39 b	34.38	46.75	34.09 a
2	33.52 b	33.46	44.46	34.55 a
4	38.43 b	34.83	44.00	33.63 a
6	49.63 a	37.58	46.29	35.47a
8	31.73 b	31.63	46.29	32.70 a
10	33.52 b	33.00	47.67	34.09 a

Means with different letter(s) are significantly different ( $p < 0.05$ ). HA, FYM, OM, N, P and K represent humic acid, farmyard manure, organic matter, nitrogen, phosphorus and potassium, respectively

## Discussion

The soil microbial respiration is the first and the most often used index that reflects the value of emitted CO<sub>2</sub> during the respiratory process of the microorganisms in the soil and shows their number as well as their metabolic activity (Parastesh *et al.*, 2019; Smitha *et al.*, 2019). Our finding indicated that humic acid improved the biological properties of soil as indicated by the enhanced releases of CO<sub>2</sub> during incubation. The supplementary experiment using lower doses of HA showed that CO<sub>2</sub> release was increased with increasing doses of HA up to 6.0 mg HA ha<sup>-1</sup> at all incubation intervals. The combine application of HA+FYM produced higher rate and cumulative CO<sub>2</sub> releases as compared to sole application of HA or control (0 HA and 0 FYM) but supremacy over sole FYM was inconsistent at different incubation intervals. These results are in accordance with Parastesh *et al.*, (2019) Sharif, (2002), and Dost & Khattak (2008) who reported increases in the activities of microbes with humic acid supplementation. Ampong *et al.*, (2022) reported that the biotic properties of soil got improved with humic acid application. Though a very little amount of humic acid (about up to 2-4 kg HA ha<sup>-1</sup>) is usually recommended as compared to the dose of 100 and 200 mg HA kg<sup>-1</sup> which are equivalent to 200 and 400 kg HA ha<sup>-1</sup>, respectively. The decrease in microbial activity with HA+FYM as compared to sole FYM as evident from cumulative release of CO<sub>2</sub> (Figs. 1 & 2) was further investigated taking lower levels

of HA as 0, 2, 4, 6, 8 and 10 mg HA kg<sup>-1</sup>. Adnan *et al.* 2018 conducted an incubated study and reported that emission of CO<sub>2</sub> was more pronounced in soils treated with organic sources (PM and FYM) of phosphorus as compared with mineral sources (SSP and RP) of phosphorus. Yang *et al.* (2021) observed a very high level of CO<sub>2</sub> emissions during week 1 of incubation. Decreased soil respiration over time may be due to a reduction in digestive processes and a lack of easily decomposing nutrients in the soil that support a limited number of soil organisms (Adnan *et al.*, 2018; Bhuma *et al.*, 2003).

HA application with FYM declined the efficiency of sole FYM which might be associated to higher levels of humic acid as 100 mg HA kg<sup>-1</sup> equivalent to 200 kg HA ha<sup>-1</sup> while HA is usually applied to soil in 2 to 4 kg ha<sup>-1</sup> (Sharif, 2002, Sharif *et al.*, 2003, Khattak & Muhammd, 2008; Chatterjee *et al.*, 2011). Janssens & Luyssaert (2009) discussed that soil fertility has a profound effect on carbon dioxide (C) in the global ecosystem. Sathiya *et al.* (2003) reported a linear trend in the release of N up to 20 and P and K up to 40 kg HA ha<sup>-1</sup>. Similarly, Sarir, (1998) and Sharif *et al.*, (2002) also advocated for use of lower doses of HA.

Soil pH showed a non-significant declining trend with HA application as compared to control or sole application of FYM. However, with application of FYM the soil pH was significantly reduced regardless of HA application. Likewise, Jamal *et al.*, (2024) found that the addition of FYM decreased soil pH in calcareous sandy soil. The soil pH could have been reduced due to the chemical oxidation and microbiological decomposition of FYM in soil, which produced acidic compounds that help reduce soil pH. The production of organic acids (amino acid, glycine, cysteine, and humic acid) during the mineralization (ammonization and ammonification) of organic materials by heterotrophs and nitrification by autotrophs can also cause a decrease in soil pH (Kumar *et al.*, 2020).

In our case, the addition of FYM also increased SOM. Other researchers also found that FYM not only reduced the oxidation stability of SOM but also improved the SOM content of the soil up to 1.2–2.9 kg ha<sup>-1</sup> (Li *et al.*, 2017; Ding *et al.*, 2020). The increase in SOM with

FYM treatment may be partially due to the input of organic matter found in the FYM (Li *et al.*, 2017; Rehim *et al.*, 2020), although the increase in SOM. We generally observed a increase in plant available N, P (AB-DTPA) and P with the addition of humic acid and FYM. In general, FYM application has appreciable and dynamic impacts on the chemical fractions of P because P from FYM gradually turns into available forms over time (Ma *et al.*, 2020). The increase in available NP and K in our study might be due to the release of significant quantities of CO<sub>2</sub> during FYM decomposition (Andriamananjara *et*

*al.*, 2019) and the complexing of cations such as Ca<sup>2+</sup>, thus reducing their fixation in calcareous soils (Fixen & Bruulsema, 2014; Jamal *et al.*, 2018; McMullen *et al.*, 2015). Furthermore, the FYM contains organic acids, which are known to increase nutrients solubility (Hopkins, 2015). HA being a poly functional molecule form chelates with nutrients and keep them in solution (Davies *et al.*, 2001). HA treated soils produce additional CO<sub>2</sub> that should be linked to improve the dissolution of Ca-apatite by carbonic acid formed due to respiration of microbial activities (Adnan *et al.*, 2019).

**Table 3. Changes in soil pH, organic matter, mineral N and AB-DTPA extractable P and K as influenced by HA and FYM at day 79 of incubation.**

HA ( mg kg <sup>-1</sup> )	FYM (g kg <sup>-1</sup> )	pH	OM (%)	N	P	K
				mg kg <sup>-1</sup>		
0	0	7.72 a	1.12 a	104.70 c	8.87 b	183 b
100	0	7.54 ab	1.15 a	109.15 c	9.17 b	187 b
200	0	7.51 b	1.14 a	119.93 b	8.99 b	181 b
0	10	7.70 ab	1.41 b	128.60 a	14.10 a	187 ab
100	10	7.67 ab	1.52 b	130.33 ab	13.49 a	187 ab
200	10	7.67 ab	1.57 b	136.20 a	12.84 a	194 a

Means with different letter(s) are significantly different (p<0.05). HA, FYM, OM, N, P and K represent humic acid, farmyard manure, organic matter, nitrogen, phosphorus and potassium, respectively

## Conclusions

The integrated application of Humic acid along with farmyard manure improved the fertility and microbial activity of soil. The supplementary experiment using lower doses of HA didn't show significant effect over CO<sub>2</sub> release except at day 4<sup>th</sup> of incubation where significantly higher CO<sub>2</sub> was released at 6.0 mg HA ha<sup>-1</sup>. The conjunctive application of HA+FYM produced higher rate and cumulative CO<sub>2</sub> releases as compared to sole application of HA or control but supremacy over sole FYM was inconsistent at different incubation intervals. Soil pH decreased, SOM along with mineral N, and AB-DTPA extractable P and K increased with HA at the end of 79 d of 1<sup>st</sup> incubation experiment. Conjunctive use of HA+FYM showed increases in N and K but declining trend in AB-DTPA extractable P and K was shown. Further experiments on conjunctive use of lower levels of HA with FYM are suggested for various soils to confirm the results.

## Acknowledgements

This article is derived from a research grant funded by the Research, Development, and Innovation Authority (RDIA) - Kingdom of Saudi Arabia with grant number (13445-Tabuk-2023-UT-R-3-1-SE).

## References

- Adnan, M., S. Fahad, I.A. Khan, M. Saeed. S. Saud, M.Z. Ihsan, M. Raiz, D. Wang and C. Wu. 2019. Integration of poultry manure and phosphate solubilizing bacteria improved availability of Ca bound P in calcareous soils. *3Biotech.*, 9(10): 1-10.
- Adnan, M., Z. Shah, M. Sharif and H. Rahman. 2018. Liming induces carbon dioxide (CO<sub>2</sub>) emission in PSB inoculated alkaline soil supplemented with different phosphorus sources. *Environ. Sci. Polut. Res.*, 25(10): 9501-9509.
- Ampong, K., M.S. Thilakarathna and L.Y. Gorim. 2022. Understanding the role of humic acids on crop performance and soil health. *Front. Agrobiol.*, 4: 848621.
- Andriamananjara, A., T. Rakotoson, T. Razafimbelo, L. Rabeharisoa, P. Razafimanantsoa and D. Masse. 2019. Farmyard manure improves phosphorus use efficiency in weathered p deficient soil. *Nutr. Cycl. Agroecosys.*, 115: 407-425.
- Bhuma, M. and G. Selvakumari. 2003. Studies on the effect of green gram to potassium humate on soil fertility. *Madras Agric. J.*, 90(7-9): 444-449.
- Bolan, N., P. Srivastava, C.S. Rao, P.V. Satyanaraya, G.C. Anderson, S. Bolan and M.B. Kirkham. 2023. Distribution, characteristics and management of calcareous soils. *Adv. Agron.*, 182: 81-130.
- Brye, K.R., B. Golden and N.A. Slaton. 2006. Poultry litter decomposition as affected by litter form and rate before flooding for rice production. *Soil Sci. Soc. Amer. J.*, 70(4): 1155-1167.
- Burton, D.L., L. Xh and C.A. Grant. 2008. Influence of fertilizer nitrogen source and management practice on N<sub>2</sub>O emissions from two Black Chernozemic soils. *Can. J. Soil Sci.*, 88(2): 219-227.
- Chatterjee, A. and G.D. Jenerette. 2011. Changes in soil respiration Q<sub>10</sub> during drying–rewetting along a semi-arid elevation gradient. *Geoderma*, 163(3-4): 171-177.
- Cimrin, K.M. and I. Yilmaz. 2005. Humic acid applications to lettuce do not improve yield but do improve phosphorus availability. *Acta Agriculturae Scandinavica*, 55(1): 58-63.
- Davies, G., E.A. Ghabbour and C. Steelink. 2001. Humic acids: Marvelous products of soil chemistry. *J. Chem. Edu.*, 78(12): 1609.
- Ding, Z., A.M. Kheir, M.G. Ali, O.A. Ali, A.I. Abdelaal, X.E. Li and Z. He. 2020. The integrated effect of salinity, organic amendments, phosphorus fertilizers, and deficit irrigation on soil properties, phosphorus fractionation and wheat productivity. *Sci. Rep.*, 10(1): 2736.
- El-Hady, O.A. and S.A. Abo-Sedera. 2006. Conditioning effect of composts and acrylamide hydrogels on a sandy calcareous soil. II-Physico-Bio-Chemical Properties of the Soil, 876-884.

- Fixen, P.E. and T.W. Bruulsema. 2014. Potato management challenges created by phosphorus chemistry and plant roots. *Amer. J. Potato Res.*, 91: 121-131.
- Horwath, W.R. and E.A. Paul. 1994. Microbial biomass. In: *Methods of soil analysis, part 2. Chemical and microbiological properties*. (Eds.): R.W. Weaver, J.S. Angle and P.S. Bottomley. *SSSA Book Series*, 5:753-771.
- Jamal, A., D. Muhammad, M.U. Rahman and H. Jamal. 2018. Application of adsorption isotherms in evaluating the influence of humic acid and farmyard manure on phosphorous adsorption and desorption capacity of calcareous soil. *World Sci. News*, (107): 136-149.
- Jamal, A., M.F. Saeed, A. Mihoub, B.G. Hopkins, I. Ahmad and A. Naem. 2023. Integrated use of phosphorus fertilizer and farmyard manure improves wheat productivity by improving soil quality and P availability in calcareous soil under subhumid conditions. *Fron. Plant Sci.*, 14: 1034421.
- Jamal, S.M., R.K. Yadav and J. Dayal. 2024. Innovative soil management technologies for a changing climate. In: *Transforming Agricultural Management for a Sustainable Future: Climate Change and Machine Learning Perspectives*, Cham: Springer Nature Switzerland, 203-213.
- Julie, C. and B. Bugbee. 2006. The use of humic acid to ameliorate iron deficiency stress. *Biol. Biochem.*, 2: 67-71.
- Katkat, A.V., H. Çelik, M.A. Turan and B.B. Asik 2009. Effects of soil and foliar applications of humic substances on dry weight and mineral nutrients uptake of wheat under calcareous soil conditions. *Aust. J. Basic App. Sci.*, 3(2): 1266-1273.
- Khattak, R.A. and D. Muhammad. 2008. Increasing crop production through humic acid in salt affected soils in Kohat division (NWFP). *Final Technical Progress Report. Pak-US Collaborative Research Endeavor, ALP project, PARC, Islamabad*.
- Kumar, V., P.K. Sharma, H.S. Jatav, S.K. Singh, A. Rai, S. Kant and A. Kumar. 2020. Organic amendments application increases yield and nutrient uptake of mustard (*Brassica juncea*) grown in chromium-contaminated soils. *Comm. Soil Sci. Plant Anal.*, 51(1): 149-159.
- Li, S., J. Li, B. Zhang, D. Li, G. Li and Y. Li. 2017. Effect of different organic fertilizers application on growth and environmental risk of nitrate under a vegetable field. *Sci. Rep.*, 7(1): 17020.
- Ma, Q., Y. Wen, J. Ma, A. Macdonald, P. Hill, Chadwick and D.L. Jone. 2020. Long-term farmyard manure application affects soil organic phosphorus cycling: A combined metagenomic and 33P/14C labelling study. *Soil Biol. Bioch.*, 149: 107959.
- McMullen, R.L., K.R. Brye and E.E. Gbur. 2015. Soil respiration as affected by long-term broiler litter application to a Uduft in the Ozark Highlands. *J. Environ. Qual.*, 44(1): 115-126.
- Melero, S., R. López-Garrido, E. Madejón, J.M. Murillo, K. Vanderlinden, R. Ordóñez and F. Moreno. 2009. Long-term effects of conservation tillage on organic fractions in two soils in southwest of Spain. *Agri. Ecosys. Environ.*, 133(1-2): 68-74.
- Minhas, A., M.S. Saeed, A. Ehsan, G. Murtaza, M. Akram, M. Mehran and R.K. Syed. 2024. Response of integrated use of humic acid and chemical fertilizer on growth and yield of rice crop (*Oryza sativa* L.) in calcareous soil. *Pak. J. Bot.*, 56(3): 879-887.
- Mulvaney, R.L. 1996. Nitrogen-Inorganic forms. In: (Ed.): D.L. Sparks. *Methods of soil analysis. Part 3. Amer. Soc. Agron.*, 38: 1123-1184.
- Nardi, S., D. Pizzeghello, A. Muscolo and A. Vianello. 2002. Physiological effects of humic substances on higher plants. *Soil Biol. Biochem.*, 34(11): 1527-1536.
- Nelson, D.W. and L.E. Sommer. 1996. Total C, organic C and organic matter. In: (Ed.): D.L. Sparks. *Method Soil Analysis part 3. Amer. Soc. Agron.*, 34: 961-1010.
- Osman, A.S. and M.M. Rady. 2012. Ameliorative effects of sulphur and humic acid on the growth, anti-oxidant levels, and yields of pea (*Pisum sativum* L.) plants grown in reclaimed saline soil. *The J. Hort. Sci. Biotech.*, 87(6): 626-632.
- Ouni, Y., T. Ghnaya, F. Montemurro, C. Abdelly and A. Lakhdar. 2014. The role of humic substances in mitigating the harmful effects of soil salinity and improve plant productivity. *Int. J. Plant Prod.*, 8(3): 353-374.
- Parastesh, F., H.A. Alikhani and H. Etesami. 2019. Vermicompost enriched with phosphate-solubilizing bacteria provides plant with enough phosphorus in a sequential cropping under calcareous soil conditions. *J. Cleaner Prod.*, 221: 27-37.
- Russel, D.F. 1989. MSTATC, version 2, Director Crops and Soil Sciences Department, Michigan State University. *Knowledge Dynamics corporation: Canyon Lake, Texas*.
- Sangeetha, M., P. Singaram and R.D. Devi. 2006. Effect of lignite humic acid and fertilizers on the yield of onion and nutrient availability. In: *Proceedings of 18th World Congress of Soil Science July*. pp. 9-15.
- Sarir, M.S. 1998. Utilization of natural resources for increasing crop production. *National seminar on sustainable management of natural resources in Pakistan. University of Peshawar*.
- Sarlaki, E.M., H. Kianmehr, N. Marzban, A. Shafizadeh, S.A. Tajuddin, S. Hu and M. Aghbashl. 2024. Advances and challenges in humic acid production technologies from natural carbonaceous material wastes. *Chem. Eng. J.*, 155521.
- Sharif, M., R.A. Khattak and M.S. Sarir. 2002b. Wheat yield and nutrients accumulation as affected by humic acid and chemical fertilizers. *Sarhad J. Agri.*, 18(3): 323-329.
- Sharif, M., R.A. Khattak and M.S. Sarir. 2002 a. Effect of different levels of lignitic coal derived humic acid on growth of maize plants. *Comm. Soil Sci. Plants Anal.*, 33: 19-20.
- Sharif, M., R.A. Khattak and M.S. Sarir. 2003. Residual effect of humic acid and chemical fertilizers on maize yield and nutrient accumulation. *Sarhad J. Agric.*, 19(4): 543-550.
- Smitha, G.R., B. Basak, B.V. Thondaiman and A. Saha. 2019. Nutrient management through organics, bio-fertilizers and crop residues improves growth, yield and quality of sacred basil (*Ocimum sanctum* Linn). *Indus. Crops Prod.*, 128: 599-606.
- Smitha, G.R., B.B. Basak, V. Thondaiman and A. Saha. 2019. Nutrient management through organics, bio-fertilizers and crop residues improves growth, yield and quality of sacred basil (*Ocimum sanctum* Linn) *Ind. Crop. Prod.*, 128: 599-606.
- Soltanpour, P.N. and A.P. Schwab. 1977. A new soil test for simultaneous extraction of macro and micro-nutrients in alkaline soils. *Comm. Soil Sci. Plant Anal.*, 8: 195-267.
- Steel, R.G.D. and J.H. Torrie. 1980. Principles and procedures of statistics. A Biometrical Approach. *McGraw-Hill, New York*.
- Yang, F., C. Tang and M. Antonietti. 2021. Natural and artificial humic substances to manage minerals, ions, water, and soil microorganisms. *Chem. Soc. Rev.*, 50: 6221-6239.

(Received for publication 21 April 2024)